

(19)  Canadian  
Intellectual Property  
Office

An Agency of  
Industry Canada

Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

(11) **CA 2 493 364**

(13) **A1**

(40) 12.02.2004

(43) 12.02.2004

(12)

(21) **2 493 364**

(22) **18.07.2003**

(51) Int. Cl. 7: **C12N 15/82, A01H 5/00,  
C12N 15/11, C12N 9/12,  
C12N 15/54**

(85) **21.01.2005**

(86) **PCT/EP03/007877**

(87) **WO04/013333**

(30) **102 34 287.3 DE 26.07.2002**

(71) **BASF PLANT SCIENCE GMBH,  
67056, LUDWIGSHAFEN, XX (DE).**

(72) **KOCK, MICHAEL (DE).  
FRANK, MARKUS (DE).  
BADUR, RALF (DE).**

(74) **ROBIC**

(54) **NOUVEAUX PROCÉDES DE SÉLECTION**

(54) **INVERSION OF THE NEGATIVE-SELECTIVE EFFECT OF NEGATIVE MARKER PROTEINS USING  
SELECTION METHODS**

(57)

The invention relates to methods for producing transformed plant cells or organisms by transforming a population of plant cells comprising at least one marker protein having a directly or indirectly toxic effect therefor, by means of at least one nucleic acid sequence to be inserted, said sequence being combined with at least one compound preferably a DNA construct which is able to reduce the expression, quantity, activity and/or function of the marker protein. The transformed plant cells have a growth advantage in relation to the non-transformed cells as a result of the action of said compound.

BEST AVAILABLE COPY



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2493364 A1 2004/02/12

(21) **2 493 364**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2003/07/18  
(87) Date publication PCT/PCT Publication Date: 2004/02/12  
(85) Entrée phase nationale/National Entry: 2005/01/21  
(86) N° demande PCT/PCT Application No.: EP 2003/007877  
(87) N° publication PCT/PCT Publication No.: 2004/013333  
(30) Priorité/Priority: 2002/07/26 (102 34 287.3) DE

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C12N 15/82, A01H 5/00, C12N 15/11,  
C12N 15/54, C12N 9/12  
(71) Demandeur/Applicant:  
BASF PLANT SCIENCE GMBH, DE  
(72) Inventeurs/Inventors:  
KOCK, MICHAEL, DE;  
FRANK, MARKUS, DE;  
BADUR, RALF, DE  
(74) Agent: ROBIC

(54) Titre : NOUVEAUX PROCEDES DE SELECTION

(54) Title: INVERSION OF THE NEGATIVE-SELECTIVE EFFECT OF NEGATIVE MARKER PROTEINS USING  
SELECTION METHODS

(57) Abrégé/Abstract:

The invention relates to methods for producing transformed plant cells or organisms by transforming a population of plant cells comprising at least one marker protein having a directly or indirectly toxic effect therefor, by means of at least one nucleic acid sequence to be inserted, said sequence being combined with at least one compound preferably a DNA construct which is able to reduce the expression, quantity, activity and/or function of the marker protein. The transformed plant cells have a growth advantage in relation to the non-transformed cells as a result of the action of said compound.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPO 191

OPIC



CIPO

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES  
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG(19) Weltorganisation für geistiges Eigentum  
Internationales Büro(43) Internationales Veröffentlichungsdatum  
12. Februar 2004 (12.02.2004)

PCT

(10) Internationale Veröffentlichungsnummer  
**WO 2004/013333 A3**(51) Internationale Patentklassifikation: **C12N 15/82**,  
9/12, 15/54, 15/11, A01H 5/00

(21) Internationales Aktenzeichen: PCT/EP2003/007877

(22) Internationales Anmeldedatum:  
18. Juli 2003 (18.07.2003)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:  
102 34 287.3 26. Juli 2002 (26.07.2002) DE(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme  
von US): BASF PLANT SCIENCE GMBH [DE/DE];  
67056 Ludwigshafen (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): KOCK, Michael  
[DE/DE]; Am Leutbusch 12, 67105 Schifferstadt  
(DE). FRANK, Markus [DE/DE]; Rheindammstr.  
30, 68163 Mannheim (DE). BADUR, Ralf [DE/DE];  
Theodor-Storm-Str. 7B, 67117 Limburgerhof (DE).(74) Anwalt: GOLDSCHIED, Bettina; c/o BASF Aktiengesellschaft,  
67056 Ludwigshafen (DE).(81) Bestimmungsstaaten (national): AE, AG, AL, AM, AT,  
AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR,  
CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KI, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,  
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,  
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.(84) Bestimmungsstaaten (regional): ARIPO Patent (GI,  
GM, KI, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW),  
eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,  
TM), europäisches Patent (AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL,  
PT, RO, SE, SI, SK, TR), OAPI Patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Veröffentlicht:

— mit internationalem Recherchenbericht

(88) Veröffentlichungsdatum des internationalen  
Recherchenberichts: 15. Juli 2004Zur Erklärung der Zweibuchstaben-Codes und der anderen Ab-  
kürzungen wird auf die Erklärungen ("Guidance Notes on Co-  
des and Abbreviations") am Anfang jeder regulären Ausgabe der  
PCT-Gazette verwiesen.(54) Title: INVERSION OF THE NEGATIVE-SELECTIVE EFFECT OF NEGATIVE MARKER PROTEINS USING SELEC-  
TION METHODS(54) Bezeichnung: REVERTIERUNG DER NEGATIV-SELEKTIVEN WIRKUNG VON NEGATIVEN MARKERPROTEINEN  
ALS SELEKTIONSVERFAHREN(57) Abstract: The invention relates to methods for producing transformed plant cells or organisms by transforming a population  
of plant cells comprising at least one marker protein having a directly or indirectly toxic effect therefor, by means of at least one  
nucleic acid sequence to be inserted, said sequence being combined with at least one compound preferably a DNA construct which  
is able to reduce the expression, quantity, activity and/or function of the marker protein. The transformed plant cells have a growth  
advantage in relation to the non-transformed cells as a result of the action of said compound.(57) Zusammenfassung: Die vorliegende Erfindung betrifft Verfahren zur Herstellung transformierter pflanzlicher Zellen oder Or-  
ganismen durch Transformation einer Population pflanzlicher Zellen, die mindestens ein Markerprotein mit einem für diese direkt  
oder indirekt toxischen Effekt umfasst, mit mindestens einer zu insertierenden Nukleinsäuresequenz in Kombination mit mindes-  
tens einer Verbindung - bevorzugt einem DNA-Konstrukt - befähigt zur Verminderung der Expression, Menge, Aktivität und/oder  
Funktion des Markerproteins, wobei die transformierten pflanzlichen Zellen infolge der Wirkung besagter Verbindung gegenüber  
nicht-transformierten Zellen einen Wachstumsvorteil haben.

WO 2004/013333 A3

**INVERSION OF THE NEGATIVE-SELECTIVE EFFECT OF NEGATIVE  
MARKER PROTEINS USING SELECTION METHODS**

Description


The present invention relates to processes for preparing transformed plant cells or organisms by transforming a population of plant cells which comprises at least one marker protein having a direct or indirect toxic effect for said population, with at least one nucleic acid sequence to be inserted in combination  
10 with at least one compound, preferably a DNA construct, capable of reducing the expression, amount, activity and/or function of the marker protein, with the transformed plant cells having a growth advantage over nontransformed cells, due to the action of said compound.

Genetic material is successfully introduced usually only into a very limited number of target cells of a population. This necessitates the distinction and isolation of successfully transformed from nontransformed cells, a process which is referred to as selection. Traditionally, the selection is carried out by way of a "positive" selection, wherein the transformed cell is enabled to grow and to survive, whereas the untransformed cell is inhibited in its growth or destroyed (McCormick et al. (1986) Plant  
20 Cell Reports 5:81-84). A positive selection of this kind is usually implemented by genes which code for a resistance to a biocide (e.g. a herbicide such as phosphinothricin, glyphosate or bromoxynil, a metabolism inhibitor such as 2-deoxyglucose 6-phosphate (WO 98/45456) or an antibiotic such as tetracycline, ampicillin, kanamycin, G 418, neomycin, bleomycin or hygromycin). Such genes are also referred to as positive selection markers. The positive selection marker is coupled (physically or by means of cotransformation) to the nucleic acid sequence to be introduced into the cell genome and is then introduced into the cell. Subsequently, the cells are cultured on a medium under the appropriate selection pressure (for example in the presence of an appropriate antibiotic or herbicide), whereby the transformed cells, owing to the required resistance to said selection pressure, have a growth/survival advantage and can thus be selected.  
30 Positive selection markers which may be mentioned by way of example are:



1a

- phosphinothricin acetyltransferases (PAT) (also: Bialophos® resistance; bar) acetylate the free amino group of the glutamine synthase inhibitor phosphinothricin (PPT) and thus achieve a detoxification (de Block et al. (1987) EMBO J



PF 53790

2

6:2513-2518; Vickers JE et al. (1996) Plant Mol Biol Reporter 14:363-368; Thompson CJ et al. (1987) EMBO J 6:2519-2523).

- 5 - 5-enolpyruvylshikimate 3-phosphate synthases (EPSPS) impart a resistance to the unselective herbicide Glyphosat® (N-(phosphonomethyl)glycine; Steinrucken HC et al. (1980) Biochem Biophys Res Commun 94:1207-1212; Levin JG and Sprinson DB (1964) J Biol Chem 239:1142-1150; Cole DJ (1985) Mode of action of glyphosate; A literature analysis, p. 48-74. In: Grossbard E and Atkinson D (eds.) The herbicide glyphosate. Buttersworths, Boston.). Glyphosate-tolerant EPSPS variants for use as selection markers have been described (Padgett SR et al. (1996). New weed control opportunities: development of soybeans with a Roundup Ready™ gene. In: Herbicide Resistant Crops (Duke SO, ed.), pp. 53-84. CRC Press, Boca Raton, FL; Saroha MK and Malik VS (1998) J Plant Biochemistry and Biotechnology 7:65-72; Padgett SR et al. (1995) Crop Science 35(5):1451-1461; US 5,510,471; US 5,776,760; US 5,864,425; US 5,633,435; US 5,627,061; US 5,463,175; EP-A 0 218 571).
- 10 -
- 15 - neomycin phosphotransferases constantly impart a resistance to aminoglycoside antibiotics such as neomycin, G418, hygromycin, paromomycin or kanamycin by reducing the inhibiting action thereof by means of a phosphorylation reaction (Beck et al. (1982) Gene 19:327-336).
- 20 -
- 25 - 2-deoxyglucose 6-phosphate phosphatases impart a resistance to 2-deoxyglucose (EP-A 0 807 836; Randez-Gil et al. (1995) Yeast 11:1233-1240; Sanz et al. (1994) Yeast 10:1195-1202).
- 30 -
- 35 - acetolactate synthases impart a resistance to imidazolinone/sulfonylurea herbicides (e.g. imazzamox, imazapyr, imazaquin, imazethapyr, amidosulfuron, azimsulfuron, chlorimuron ethyl, chlorsulfuron; Sathasivan K et al. (1990) Nucleic Acids Res 18(8):2188).

40 In addition, resistance genes to the antibiotics hygromycin (hygromycin phosphotransferases), chloramphenicol (chloramphenicol acetyltransferase), tetracycline, streptomycin, zeocine and ampicillin ( $\beta$ -lactamase gene; Datta N, Richmond MH. (1966) Biochem J 98(1):204-9) have been described.

45 Genes such as isopentenyl transferase (ipt) from Agrobacterium tumefaciens (strain:PO22) (GenBank Acc. No.: AB025109) may likewise be used as selection markers. The ipt gene is a key enzyme of cytokine biosynthesis. Its overexpression facilitates the re-

## 3

generation of plants (e.g. selection on cytokine-free medium) (Ebinuma H et al. (2000) Proc Natl Acad Sci USA 94:2117-2121; Ebinuma H et al. (2000) Selection of Marker-free transgenic plants using the oncogenes (*ipt*, *rol* A, B, C) of *Agrobacterium* as  
5 selectable markers, In Molecular Biology of Woody Plants. Kluwer Academic Publishers). The disadvantages here are, firstly, the fact that the selection disadvantage is based on usually subtle differences in cell proliferation and, secondly, the fact that the plant acquires unwanted properties (gall tumor formation) due  
10 to transformation with an oncogene.

EP-A 0 601 092 describes various other positive selection markers. Examples which may be mentioned are:  $\beta$ -glucuronidase (in connection with, for example, cytokinin glucuronide), mannose  
15 6-phosphate isomerase (in connection with mannose), UDP-galactose 4-epimerase (in connection with galactose, for example).

Negative selection markers are used for selecting organisms in  
20 which marker sequences have been successfully deleted (Koprek T et al. (1999) Plant J 19(6):719-726). In the presence of a negative selection marker, the corresponding cell is destroyed or experiences a growth disadvantage. Negative selection involves, for example, the negative selection marker introduced into the plant  
25 converting a compound which otherwise has no action disadvantageous to the plant into a compound with a disadvantageous (i.e. toxic) action. Examples of negative selection markers include: thymidine kinase (TK), for example of Herpes simplex virus (Wigler et al. (1977) Cell 11:223), cellular adenine phosphoribosyl  
30 transferase (APRT) (Wigler et al. (1979) Proc Natl Acad Sci USA 76:1373), hypoxanthine phosphoribosyl transferase (HPRT) (Jolly et al. (1983) Proc Natl Acad Sci USA 80:477), diphtheria toxin A fragment (DT-A), the bacterial xanthine-guanine  
phosphoribosyl transferase (*gpt*; Besnard et al. (1987) Mol. Cell. Biol. 7:4139; Mzoz and Moolten (1993) Human Gene Therapy  
35 4:589-595), the *codA* gene product coding for a cytosine deaminase (Gleave AP et al. (1999) Plant Mol Biol. 40(2):223-35; Perera RJ et al. (1993) Plant Mol Biol 23(4): 793-799; Stougaard J; (1993) Plant J 3:755-761; EP-A1 595 873), the cytochrome P450 gene (Koprek et al. (1999) Plant J 16:719-726), genes coding for a haloalkane dehalogenase (Naested H (1999) Plant J 18:571-576), the  
40 *iaaH* gene (Sundaresan V et al. (1995) Genes & Development 9:1797-1810) or the *tms2* gene (Fedoroff NV & Smith DL (1993) Plant J 3: 273-289). The negative selection markers are usually  
45 employed in combination with "prodrugs" or "pro-toxins", compounds which are converted into toxins by the activity of the selection marker.

## 4

5-Methylthioribose (MTR) kinase is an enzyme whose enzymic activity in plants, bacteria and protozoa, but not in mammals, has been described. The enzyme may convert an MTR analog (5-(triomethyl)thioribose) as a "subversive substrate" of the methionine salvage pathway via an unstable intermediate to give the toxic compound carbothionyl difluoride.

Said selection systems have various disadvantages. The introduced selection marker (e.g. resistance to antibiotics) is justified only during transformation and selection but is later a usually unnecessary and often also undesired protein product. This may be disadvantageous for reasons of consumer acceptance and/or approval as a food and/or feed product. Another disadvantage in this connection is the fact that the selection marker used for selection is usually genetically coupled to the nucleic acid sequence to be inserted into the genome and cannot be decoupled by segregation during propagation or crossing. Usually, deletion of the marker sequence is required, making additional steps necessary. In addition, biotechnological studies require in numerous cases multiple transformation with various gene constructs. Here, each transformation step requires a new selection marker unless the previously used marker is to be laboriously deleted first. This, however, necessitates a broad palette of well-functioning selection markers which are not available for most plant organisms.

Consequently, it was the object of the invention to provide novel selection processes for selecting transformed plant cells and organisms, which, if possible, no longer have the disadvantages of the available systems. This object is achieved by the present invention.

The invention firstly relates to a process for preparing transformed plant cells or organisms, which process comprises the following steps:

- a) transforming a population of plant cells, with the cells of said population containing at least one marker protein capable of causing directly or indirectly a toxic effect for said population, with at least one nucleic acid sequence to be inserted in combination with at least one compound capable of reducing the expression, amount, activity and/or function of at least one marker protein, and
- b) selecting transformed plant cells whose genome contains said nucleic acid sequence and which have a growth advantage

## 5

tage over nontransformed cells, due to the action of said compound, from said population of plant cells, the selection being carried out under conditions under which the marker protein can exert its toxic effect on the non-transformed cells.

In a preferred embodiment, the marker protein is a protein capable of converting directly or indirectly a substance X which is nontoxic for said population of plant cells into a substance Y which is toxic for said population. In this case, the process of the invention preferably comprises the following steps:

- a) transforming the population of plant cells with at least one nucleic acid sequence to be inserted in combination with at least one compound capable of reducing the expression, amount, activity and/or function of at least one marker protein, and
- b) treating said population of plant cells with the substance X at a concentration which causes a toxic effect for nontransformed cells, due to the conversion by the marker protein, and
- c) selecting transformed plant cells whose genome contains said inserted nucleic acid sequence and which have a growth advantage over nontransformed cells, due to the action of said compound, from said population of plant cells, the selection being carried out under conditions under which the marker protein can exert its toxic effect on the nontransformed cells.

The nontoxic substance X is preferably a substance which does not naturally occur in plant cells or organisms or occurs naturally therein only at a concentration which can essentially not cause any toxic effect. In the scope of the process of the invention, preference is given to applying the nontoxic substance X exogenously, for example via the medium or the growth substrate.

The term "compound capable of reducing the expression, amount, activity and/or function of at least one marker protein" is to be understood broadly and generally means any compounds which cause, directly or indirectly, alone or in cooperation with other factors, a reduction in the amount of protein, amount of RNA, gene activity, protein activity or protein function of at least one marker protein. Said compounds are also referred to under the generic term "anti-marker protein" compounds. The term "anti-marker

## 6

protein" compound includes in particular, but is not limited to, the nucleic acid sequences, ribonucleic acid sequences, double-stranded ribonucleic acid sequences, antisense ribonucleic acid sequences, expression cassettes, peptides, proteins or other factors used in the preferred embodiments within the scope of the process of the invention.

In a preferred embodiment, "anti-marker protein" compound means a DNA construct comprising

- a) at least one expression cassette suitable for expressing a ribonucleic acid sequence and/or, if appropriate, a protein, said nucleic acid sequence and/or protein being capable of reducing the expression, amount, activity and/or function of the marker protein, or
- b) at least one sequence which causes a partial or complete deletion or inversion of the sequence coding for said marker protein and thus enables the expression, amount, activity and/or function of the marker protein to be reduced, and also, if appropriate, further functional elements which facilitate and/or promote said deletion or inversion, or
- c) at least one sequence which causes an insertion into the sequence coding for said marker protein and thus enables the expression, amount, activity and/or function of the marker protein to be reduced, and also, if appropriate, further functional elements which facilitate and/or promote said insertion.

The process of the invention stops the negative-selective action of the marker protein. To this extent, an "anti-marker protein" compound acts directly (e.g. via inactivation by means of insertion into the gene coding for the marker protein) or indirectly (e.g. by means of the ribonucleic acid sequence expressed via the expression cassette and/or, where appropriate, of the protein translated therefrom) as a positive selection marker. Hence, the selection system of the invention is to be referred to as a "reverse selection system", since it "reverts" the negative-selective action of the marker protein.

The process of the invention means a drastic broadening of the repertoire of positive selection processes for selecting transformed plant cells.

Another advantage is the fact that in a particular, preferred embodiment (e.g. via the action of a double-stranded or antisense RNA), it is possible to implement the selection effect without expressing a foreign protein (see below).

5

It is also advantageous that the marker protein used indirectly for selection (e.g. the negative selection marker) is not coupled genetically to the nucleic acid sequence to be inserted into the genome. In contrast to the otherwise customary selection processes, the marker protein, if it is a transgene, may be removed by simple segregation in the course of subsequent propagation or crossing.

15 "Plant cell" means within the scope of the present invention any type of cell which has been derived from a plant organism or is present therein. In this context, the term includes by way of example protoplasts, callus or cell cultures, microspores, pollen, cells in the form of tissues such as leaves, meristem, flowers, 20 embryos, roots, etc. Included are, in particular, all of those cells and cell populations which are suitable as target tissues for a transformation.

In this context, "plant organism" comprises any organism capable 25 of photosynthesis and also the cells, tissues, parts or propagation material (such as seeds or fruits) derived therefrom. Included within the scope of the invention are all genera and species of higher and lower plants of the plant kingdom. Preference is given to annual, perennial, monocotyledonous and dicotyledonous 30 plants and also gymnosperms.

"Plant" means within the scope of the invention all genera and species of higher and lower plants of the plant kingdom. The term 35 includes the mature plants, seed, shoots and seedlings, and also parts, propagation material (for example tubers, seeds or fruits), plant organs, tissues, protoplasts, callus and other cultures, for example cell cultures, derived therefrom, and also any other types of groupings of plant cells to give functional or structural units. Mature plants means plants at any developmental 40 stage beyond that of the seedling. Seedling means a young immature plant at an early developmental stage. "Plant" comprises all annual and perennial monocotyledonous and dicotyledonous plants and includes by way of example but not by limitation those of the genera Cucurbita, Rosa, Vitis, Juglans, Fragaria, Lotus, Medicago, Onobrychis, Trifolium, Trigonella, Vigna, Citrus, Linum, Geranium, Manihot, Daucus, Arabidopsis, Brassica, Raphanus, Sinapis, 45 Atropa, Capsicum, Datura, Hyoscyamus, Lycopersicon,

## 8

Nicotiana, Solarium, Petunia, Digitalis, Majorana, Cichorium, Helianthus, Lactuca, Bromus, Asparagus, Antirrhinum, Heterocallis, Nemesis, Pelargonium, Panieum, Pennisetum, Ranunculus, Senecio, Salpiglossis, Cucumis, Browaalia, Glycine, Pisum, Phaseolus, Lolium, Oryza, Zea, Avena, Hordeum, Secale, Triticum, Sorghum, Picea and Populus.

Preference is given to plants of the following plant families:

10 Amaranthaceae, Asteraceae, Brassicaceae, Carophyllaceae, Chenopodiaceae, Compositae, Cruciferae, Cucurbitaceae, Labiatae, Leguminosae, Papilionoideae, Liliaceae, Linaceae, Malvaceae, Rosaceae, Rubiaceae, Saxifragaceae, Scrophulariaceae, Solanaceae, Sterculiaceae, Tetragoniaceae, Theaceae, Umbelliferae.

15 Preferred monocotyledonous plants are selected in particular from the monocotyledonous crop plants such as, for example, those in the family of Gramineae such as alfalfa, rice, corn, wheat or other cereal species such as barley, millet, rye, triticale or  
20 oats and also from sugar cane and all grass species.

Preferred dicotyledonous plants are selected in particular from the dicotyledonous crop plants such as, for example,

- Asteraceae, such as sunflower, tagetes or calendula and others,

25

- Compositae, in particular the genus Lactuca, very especially the species sativa (lettuce) and others,

30 - Cruciferae, especially the genus Brassica, very especially the species napus (oilseed rape), campestris (beet), oleracea cv Tastie (cabbage), oleracea cv Snowball Y (cauliflower) and oleracea cv Emperor (broccoli) and other cabbage species; and the genus Arabidopsis, very especially the species thaliana, and  
35 cress or canola and others,

- Cucurbitaceae, such as melon, pumpkin/squash or zucchini and others,

40 - Leguminosae, especially the genus Glycine, very especially the species max (soybean) and alfalfa, pea, bean plant or peanut, and others

45 - Rubiaceae, preferably the subclass Lamiidae, such as, for example, Coffea arabica or Coffea liberica (coffee bush) and others,



## 9

- Solanaceae, in particular the genus *Lycopersicon*, very especially the species *esculentum* (tomato), the genus *Solanum*, very especially the species *tuberosum* (potato) and *melongena* (eggplant), and the genus *Capsicum*, very especially the species *annuum* (pepper) and tobacco and others,

- Sterculiaceae, preferably the subclass Dilleniidae, such as, for example, *Theobroma cacao* (cacao tree) and others,

- Theaceae, preferably the subclass Dilleniidae, such as, for example, *Camellia sinensis* or *Thea sinensis* (tea shrub) and others,

- Umbelliferae, especially the genus *Daucus* (very especially the species *carota* (carrot)) and *Apium* (very especially the species *graveolens dulce* (celery)) and others,

- and linseed, cotton, hemp, flax, cucumber, spinach, carrot, sugar beet and the various tree, nut and grapevine species, in particular banana and kiwi.

- Plant organisms for the purposes of the invention are furthermore other photosynthetically active capable organisms such as, for example, algae, cyanobacteria and mosses. Preferred algae are green algae such as, for example, algae of the genus *Haematococcus*, *Phaedactylum tricornatum*, *Volvox* or *Dunaliella*. Particular preference is given to *Synechocystis*.

- Particular preference is given to the group of plants, consisting of wheat, oats, millet, barley, rye, corn, rice, buckwheat, sorghum, triticale, spelt, linseed, sugar cane, oilseed rape, cress, *Arabidopsis*, cabbage species, soybean, alfalfa, pea, bean plants, peanut, potato, tobacco, tomato, eggplant, paprika, sunflower, tagetes, lettuce, calendula, melon, pumpkin and zucchini.

Most preference is given to

- a) plants suitable for producing oil, such as, for example, oilseed rape, sunflower, sesame, safflower (*Carthamus tinctorius*), olive tree, soybean, corn, peanut, ricinus, oil palm, wheat, cacao tree or various nut species such as, for example, walnut, coconut or almond. Among these, particular preference is in turn given to dicotyledonous plants, in particular oilseed rape, soybean and sunflower.

## 10

- b) plants suitable for producing starch, such as corn, wheat or potato, for example.
- 5 c) plants which are utilized as food and/or feedstuff and/or as useful plants and in which a resistance to pathogens would be advantageous, such as barley, rye, rice, potato, cotton, flax or linseed, for example.
- 10 d) plants which may be suitable for producing fine chemicals such as, for example, vitamins and/or carotenoids, such as oilseed rape, for example.

"Population of plant cells" means any group of plant cells, which  
15 may be subjected within the scope of the present invention to a transformation and from which transgenic plant cells transformed by the process of the invention may be obtained and isolated. In this context, said population may also be, for example, a plant tissue, organ or a cell culture, etc. Said population may com-  
20 prise by way of example but not by limitation an isolated zygote, an isolated immature embryo, embryogenic callus, plant or else various flower tissues (both in vitro and in vivo).

"Genome" means the entirety of genetic information of a plant  
25 cell and comprises both genetic information of the nucleus and that of the plastids (e.g. chloroplasts) and mitochondria. However, genome preferably means the genetic information of the nucleus (for example of the nuclear chromosomes).

30 "Selection" means identifying and/or isolating successfully transformed plant cells from a population of nontransformed cells by using the process of the invention. This does not necessarily require that the selection be carried out directly with the transformed cells immediately after transformation. It is also  
35 possible to carry out the selection only at a later time, even with a later generation of the plant organisms (or cells, tissues, organs or propagation material derived therefrom) resulting from the transformation. Thus it is possible, for example, to transform Arabidopsis plants directly using, for example, the  
40 vacuum infiltration method (Clough S & Bent A (1998) Plant J 16(6):735-43; Bechtold N et al. (1993) CR Acad Sci Paris 1144(2):204-212), which subsequently produce transgenic seeds which may then be subjected to selection.

45 The fact that the nucleic acid sequence to be inserted is transformed "in combination with" the "anti-marker protein" compound (e.g. a DNA construct) is to be understood broadly and means that

## 11

- at least one nucleic acid sequence to be inserted and at least one "anti-marker protein" compound are functionally coupled to one another so that the presence of the "anti-marker protein" compound in the plant cell, and of the selection advantage related thereto, indicates the parallel presence of the inserted nucleic acid sequence as likely. The nucleic acid sequence to be inserted and the "anti-marker protein" compound (e.g. a DNA construct) here may be, preferably but not necessarily, part of a single nucleic acid construct (e.g. a transformation construct or transformation vector), i.e. be present physicochemically coupled via a covalent bond. However, they may also be jointly introduced separately, for example in the course of a cotransformation, and exert their function within the scope of the process of the invention also in this way. In the case of the "anti-marker protein compound" acting via expressing an RNA (e.g. an antisense RNA or double-stranded RNA) or being such an RNA, "in combination" may also include those embodiments in which said RNA and the RNA expressed by the nucleic acid sequence inserted into the genome form an RNA strand.
- 20 "Nontoxic substance X" generally means substances which, compared to their reaction product Y, under otherwise identical conditions, have a reduced, preferably an essentially lacking biological activity, preferably toxicity. In this context, the toxicity of substance Y is at least twice as high as that of substance X, preferably at least five times as high, particularly preferably at least ten times as high, very particularly preferably at least twenty times as high, most preferably at least one hundred times as high. "Identical conditions" here means that all conditions are kept the same, apart from the different substances X and Y.
- 30 Accordingly, identical molar concentrations of X and Y are used, with the medium, temperature, type of organism and density of organism, etc. being the same. The substance X may be converted to the substance Y in various ways, for example by hydrolysis, deamination, hydrolysis, dephosphorylation, phosphorylation, oxidation or any other type of activation, metabolization or conversion. The substance X may be, by way of example but not by limitation, the inactive precursor or derivative of a plant growth regulator or herbicide.
- 40 "Toxicity" or "toxic effect" means a measurable, negative influence on the physiology of the plant or of the plant cell and may comprise here symptoms such as, for example, but not limited thereto, a reduced or disrupted growth, a reduced or disrupted rate of photosynthesis, a reduced or disrupted cell division, a reduced or disrupted regeneration of a complete plant from cell culture or callus, etc.

## 12

The plant cells successfully transformed by means of the process of the invention may, to put it differently, have a growth advantage or selection advantage over the nontransformed cells of the same starting population under the influence of the substance

5 "X". Growth or selection advantage is to be understood here broadly and means, for example, the fact that said transformed plant cells are capable of forming shoots and/or can be regenerated to give complete plants, whereas the nontransformed cells can do this only with a marked delay, if at all.

10

The term of "marker protein" is to be understood broadly and generally means all of those proteins which are capable of

15 i) exerting per se a toxic effect on the plant or plant cell, or

ii) converting directly or indirectly a nontoxic substance X into a substance Y which is toxic for the plant or plant cell.

20

In this context, the marker protein may be a plant-intrinsic, endogenous gene or else a transgene from a different organism. Preferably, the marker protein itself has no essential function for the organism including the marker protein. If the marker protein

25 per se exerts a toxic effect, then it will preferably be expressed, for example, under an inducible promoter rather than constitutively.

30 Preferably, however, the marker protein converts directly or indirectly a nontoxic substance X into a substance Y which is toxic for the plant or plant cell. Particularly preferred marker proteins are the "negative selection markers" as are used, for example, in the course of targeted deletions from the genome.

35 Examples of marker proteins which may be mentioned but which are not limiting are:

(a) cytosine deaminases (Coda or CDase), with preference being given to using as the nontoxic substance X substances such as

40 5-fluorocytosine (5-FC). Cytosine deaminases catalyze the deamination of cytosine to give uracil (Kilstrup M et al. (1989) J Bacteriol 171:2124-2127; Anderson L et al. (1989) Arch Microbiol 152:115-118). Bacteria and fungi which have CDase activity convert 5-FC to the toxic metabolite ("Y")

45 5-fluorouracil (5-FU) (Polak A & Scholer HJ (1975) Chemotherapy (Basel) 21:113-130). 5-FC itself has low toxicity (Bennett JE, in Goodman and Gilman: the Pharmacological Basis

## 13

- of Therapeutics. 8th ed., eds. Gilman AG et al. (Pergamon Press, New York) pp. 1165-1181). However, 5-FU has a highly cytotoxic effect, since it is subsequently metabolized to fluoro-UTP (FUTP) and fluoro-dUMP (FdUMP) and thus inhibits RNA and DNA synthesis (Calabrisi P & Chabner BA in Goodman and Gilman: the Pharmacological Basis of Therapeutics. 8th ed., eds. Gilman AG et al. (Pergamon Press, New York) pp. 1209-1263); Damon LE et al. (1989) Pharmac Ther 43:155-189).
- Cells of higher plants and mammalian cells have no significant CDase activity and cannot deaminate 5-FC (Polak A et al. (1976) Chemotherapy 22:137-153; Koechlin BA et al. (1966) Biochemical Pharmacology 15:434-446). In this respect, the CDase is introduced as a transgene (e.g. in the form of a transgenic expression cassette) into plant organisms in the course of the process of the invention. Corresponding transgenic plant cells or organisms are then used as masterplants as starting material. Appropriate CDase sequences, transgenic plant organisms and the process of carrying out negative selection processes using, for example, 5-FC as nontoxic substance X, are known to the skilled worker (WO 93/01281; US 5,358,866; Gleave AP et al. (1999) Plant Mol Biol 40(2):223-35; Perera RJ et al. (1993) Plant Mol Biol 23(4):793-799; Stougaard J (1993) Plant J 3:755-761); EP-A1 595 837; Mullen CA et al. (1992) Proc Natl Acad Sci USA 89(1):33-37; Kobayashi T et al. (1995) Jpn J Genet 70(3):409-422; Schlaman HRM & Hooykaas PFF (1997) Plant J 11:1377-1385; Xiaohui Wang H et al. (2001) Gene 272(1-2): 249-255; Koprek T et al. (1999) Plant J 19(6):719-726; Gleave AP et al. (1999) Plant Mol Biol 40(2):223-235; Gallego ME (1999) Plant Mol Biol 39(1):83-93; Salomon S & Puchta H (1998) EMBO J 17(20):6086-6095; Thykjaer T et al. (1997) Plant Mol Biol 35(4):523-530; Serino G (1997) Plant J 12(3):697-701; Risseuw E (1997) Plant J 11(4):717-728; Blanc V et al. (1996) Biochimie 78(6):511-517; Corneille S et al. (2001) Plant J 27:171-178). Cytosine deaminases and the genes coding therefor may be obtained from a multiplicity of organisms, preferably microorganisms such as, for example, the fungi *Cryptococcus neoformans*, *Candida albicans*, *Torulopsis glabrata*, *Sporothrix schenckii*, *Aspergillus*, *Cladosporium* and *Phialophora* (JE Bennett, Chapter 50: Antifungal Agents, in Goodman and Gilman's the Pharmacological Basis of Therapeutics 8th ed., A.G. Gilman, ed., Pergamon Press, New York, 1990) and the bacteria *E.coli* and *Salmonella typhimurium* (Andersen L et al. (1989) Arch Microbiol 152:115-118).

## 14

The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

5 Particular preference is given to sequences according to Gen-Bank Acc. No: S56903, and to the modified *codA* sequences described in EP-A1 595 873, which make expression in eukaryotes possible. Preference is given here to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 2 or, preferably, 4, in particular the sequences according to SEQ ID NO: 10 1 or, preferably, 3.

(b) cytochrome P-450 enzymes, in particular the bacterial cytochrome P-450 *SU1* gene product (CYP105A1) from *Streptomyces griseolus* (strain ATCC 11796), with preference being given to using as nontoxic substance X substances such as the pro sulfonylurea herbicide R7402 (2-methylethyl-2-3-dihydro-N-[(4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]-1,2-benzothiazole-7-sulfonamide 1,1-dioxide). Corresponding sequences and the process of carrying out negative selection processes using, for example, R7402 as nontoxic substance X are known to the skilled worker (O'Keefe DP et al. (1994) *Plant Physiol* 105:473-482; Tissier AF et al. (1999) *Plant Cell* 11:1841-1852; Koprek T et al. (1999) *Plant J* 19(6):719-726; 25 O'Keefe DP (1991) *Biochemistry* 30(2):447-55). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

30 Particular preference is given to sequences according to Gen-Bank Acc. No: M32238. Preference is further given to nucleic acid sequences coding for the polypeptide according to SEQ ID NO: 6, in particular the sequence according to SEQ ID NO: 5.

35 (c) indoleacetic acid hydrolases such as, for example, *Agrobacterium tumefaciens*, *tms2* gene product, with preference being given to using as nontoxic substance X substances such as auxin amide compounds or naphthaleneacetamide (NAM) (with NAM being converted to naphthaleneacetic acid, a phytotoxic substance). Corresponding sequences and the process of carrying out negative selection processes using, for example, NAM as nontoxic substance X are known to the skilled worker (Fedoroff NV & Smith DL (1993) *Plant J* 3:273-289; Upadhyaya NM et al. (2000) *Plant Mol Biol Rep* 18:227-223; Depicker AG et al. (1988) *Plant Cell rep* 104:1067-1071; Karlin-Neumann GA et al. (1991) *Plant Cell* 3:573-582; Sundaresan V et al. 45 (1995) *Gene Develop* 9:1797-1810; Cecchini E et al. (1998) *Mutat Res* 401(1-2):199-206; Zubko E et al. (2000) *Nat Biotech-*

## 15

not 18:442-445). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

5 Particular preference is given to sequences according to Gen-Bank Acc. No: NC\_003308 (Protein\_id="NP\_536128.1), AE009419, AB016260 (Protein\_id="BAA87807.1) and NC002147. Preference is further given to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 8 or 10, in particular the sequences according to SEQ ID NO: 7 or 9.

15 (d) haloalkane dehalogenases (dhla gene product), for example from Xanthobacter autotrophicus GJ10. The dehalogenase hydrolyzes dihaloalkanes such as 1,2-dichloroethane (DCE) to give halogenated alcohols and inorganic halides (Naested H et al. (1999) Plant J 18(5):571-576; Janssen DB et al. (1994) Annu Rev Microbiol 48: 163-191; Janssen DB (1989) J Bacteriol 171(12):6791-9). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

25 Particular preference is given to sequences according to Gen-Bank Acc. No: M26950. Preference is further given to nucleic acid sequences coding for the polypeptide according to SEQ ID NO: 12, in particular the sequence according to SEQ ID NO: 11.

30 (e) thymidine kinases (TK), in particular viral TKs from viruses such as Herpes simplex virus, SV40, cytomegalovirus, Varicella zoster virus, in particular the TK of Herpes simplex virus type 1 (TK HSV-1), with preference being given to using as nontoxic substance X substances such as Acyclovir, Ganciclovir or 1,2-deoxy-2-fluoro- $\beta$ -D-arabinofuranosil-5-iodouracil (FIAU). Corresponding sequences and the process of carrying out negative selection processes using, for example, Acyclovir, Ganciclovir or FIAU as nontoxic substance X are known to the skilled worker (Czako M & Marton L (1994) Plant Physiol 104:1067-1071; Wigler M et al. (1977) Cell 11(1):223-232; McKnight SL et al. (1980) Nucl Acids Res 8(24):5949-5964; McKnight SL et al. (1980) Nucl Acids Res 8(24):5931-5948; Preston et al. (1981) J Virol 38(2):593-605; Wagner et al. (1981) Proc Natl Acad Sci USA 78(3):1441-1445; St. Clair et al. (1987) Antimicrob Agents Chemother 31(6):844-849). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

16

itly referred to.

5 Particular preference is given to sequences according to Gen-Bank Acc. No: J02224, V00470 and V00467. Preference is also given to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 14 or 16, in particular the sequences according to SEQ ID NO: 13 or 15.

10 (f) guanine phosphoribosyl transferases, hypoxanthine phosphoribosyl transferases or xanthine guanine phosphoribosyl transferases, with preference being given to using as nontoxic substance X substances such as 6-thioxanthine or allopurinol. Preference is given to guanine phosphoribosyl transferases  
15 (gpt), for example from E. Coli (Besnard et al. (1987) Mol Cell Biol 7:4139; Mzoz and Moolten (1993) Human Gene Therapy 4:589-595; Ono et al. (1997) Hum Gene Ther 8(17):2043-55), hypoxanthine phosphoribosyl transferases (HPRT; Jolly et al. (1983) Proc Natl Acad Sci USA 80:477; Fonwick "The HGPRT System", pp. 333-373, M. Gottesman (ed.), Molecular Cell Genetics, John Wiley and Sons, New York, 1985), xanthine guanine  
20 phosphoribosyl transferases, for example from Toxoplasma gondii (Knoll LJ et al. (1998) Mol Cell Biol 18(2):807-814; Donald RG et al. (1996) J Biol Chem 271(24):14010-14019). The sequences, materials and processes disclosed in the context  
25 of said publications are hereby explicitly referred to.

Particular preference is given to sequences according to Gen-Bank Acc. No: U10247 (Toxoplasma gondii HXGPRT), M13422  
30 (E. coli gpt) and X00221 (E. coli gpt). Preference is also given to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 18, 20 or 22, in particular the sequences according to SEQ ID NO: 17, 19 or 21.

35 (g) purine nucleoside phosphorylases (PNP; DeoD gene product), for example from E. coli, with preference being given to using as nontoxic substance X substances such as 6-methylpurine deoxyribonucleoside. Corresponding sequences and the process of carrying out negative selection processes using, for example,  
40 6-methylpurine deoxyribonucleoside as nontoxic substance X are known to the skilled worker (Sorscher EJ et al. (1994) Gene Therapy 1:233-238). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

45

Particular preference is given to sequences according to Gen-Bank Acc. No: M60917. Preference is also given to nucleic



## 17

acid sequences coding for the polypeptide according to SEQ ID NO: 24, in particular the sequence according to SEQ ID NO: 23.

- 5 h) phosphonate monoester hydrolases which convert inactive ester derivatives of the herbicide glyphosate (e.g. glycerylglyphosate) into the active form of the herbicide. Corresponding sequences and the process of carrying out negative selection processes using, for example, glycerylglyphosate are known to the skilled worker (US 5,254,801; Dotson SB et al. (1996) Plant J 10(2):383-392; Dotson SB et al. (1996) J Biol Chem 271(42): 25754-25761). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

Particular preference is given to sequences according to Gen-Bank Acc. No: U44852. Preference is also given to nucleic acid sequences coding for the polypeptide according to SEQ ID NO: 26, in particular the sequence according to SEQ ID NO: 25.

- (i) aux-1 and, preferably, aux-2 gene products, for example of the Ti plasmids of Agrobacterium strains such as A.rhizogenes or A.tumefaciens (Beclin C et al. (1993) Transgenics Res 2:4855); Gaudin V, Jouanin L. (1995) Plant Mol Biol. 28(1):123-36.

The activity of the two enzymes causes the plant cell to produce indoleacetamide (IAA). Aux-1 encodes an indoleacetamide synthase (IAMS) and converts tryptophan into indoleacetamide (VanOnckelen et al. (1986) FEBS Lett. 198: 357-360). Aux-2 encodes the enzyme indoleacetamide hydrolase (IAMH) and converts indoleacetamide, a substance without phytohormone activity, into the active auxin indoleacetic acid (Inze D et al. (1984) Mol Gen Genet 194:265-274; Tomashow et al. (1984) Proc Natl Acad Sci USA 81:5071-5075; Schroder et al. (1984) Eur J Biochem 138:387-391). The enzyme IAMH may also hydrolyze a number of indoleamide substrates such as, for example, naphthaleneacetamide, the latter being converted into the plant growth regulator naphthaleneacetic acid (NAA). The use of the IAMH gene as a negative selection marker is described, for example, in US 5,180,873. Corresponding enzymes have also been described in A. rhizogenes, A. vitis (Canaday J et al. (1992) Mol Gen Genet 235:292-303) and Pseudomonas savastanoi (Yamada et al. (1985) Proc Natl Acad Sci USA 82:6522-6526). The use as a negative selection marker for destroying partic-

## 18

ular cell tissues (e.g. pollen; US 5,426,041) or transgenic plants (US 5,180,873) has been described. Corresponding sequences and the process of carrying out negative selection processes using, for example, naphthaleneacetamide are known to the skilled worker (see above). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

Particular preference is given to sequences according to the GenBank Acc. No: M61151, AF039169 and AB025110. Preference is also given to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 28, 30, 32, 34 or 36, in particular the sequences according to SEQ ID NO: 27, 29, 31, 33 or 35.

(j) adenine phosphoribosyl transferases (APRT), with preference being given to using as nontoxic substance X substances such as 4-aminopyrazolopyrimidine. Corresponding sequences and the process of carrying out negative selection processes with use are known to the skilled worker (Wigler M et al. (1979) Proc Natl Acad Sci USA 76(3):1373-6; Taylor et al. "The APRT System", pp., 311-332, M. Gottesman (ed.), Molecular Cell Genetics, John Wiley and Sons, New York, 1985).

k) methoxinine dehydrogenases, with preference being given to using as nontoxic substance X substances such as 2-amino-4-methoxybutanoic acid (methoxinine) which is converted into the toxic methoxyvinyl glycine (Margraff R et al. (1980) Experimentia 36: 846).

l) rhizobitoxin synthases, with preference being given to using as nontoxic substance X substances such as 2-amino-4-methoxybutanoic acid (methoxinine) which is converted into the toxic 2-amino-4-[2-amino-3-hydroxypropyl]-trans-3-butanoic acid (rhizobitoxin) (Owens LD et al. (1973) Weed Science 21:63-66),

m) 5-methylthioribose (MTR) kinases, with preference being given to using as nontoxic substance X substances such as 5-(trifluoromethyl)thioribose (MTR analog, "subversive substrate") which is converted, via an unstable intermediate, into the toxic substance (Y) carbothionyl difluoride. The MTR kinase is a key enzyme of the methionine salvage pathway. Corresponding enzyme activities have been described in plants, bacteria and protozoa but not in mammals. MTR kinases of various species have been identified owing to defined sequence motifs (Sekowska A et al. (2001) BMC Microbiol 1:15;

19

<http://www.biomedcentral.com/1471-2180/1/15>). Corresponding sequences and the process of carrying out negative selection processes using, for example, 5-(trifluoromethyl)thioribose are known to the skilled worker and readily obtainable from the appropriate sequence database (e.g. GenBank) (Sekowska A et al. (2001) BMC Microbiol 1:15; Cornell KA et al. (1996) 317:285-290). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.

However, a plant MTR kinase has not yet been identified unambiguously and is provided within the scope of the process of the invention (SEQ ID NO: 39 and, respectively, 40). In addition, homologs from other plant species are provided, namely from corn (SEQ ID NO: 59 and, respectively, 60), oilseed rape (SEQ ID NO: 61, 63 and, respectively, 62, 64), rice (SEQ ID NO: 65 and, respectively, 66) and soybean (SEQ ID NO: 67 and, respectively, 68).

Accordingly, the invention further relates to amino acid sequences encoding a plant 5-methylthioribose kinase, wherein said amino acid sequence contains at least one sequence selected from the group consisting of SEQ ID NO: 60, 62, 64, 66 or 68.

Accordingly, the invention further relates to nucleic acid sequences encoding a plant 5-methylthioribose kinase, wherein said nucleic acid sequence contains at least one sequence selected from the group consisting of SEQ ID NO: 59, 61, 63, 65 or 67. Even if said sequences are in parts only fragments of complete cDNAs, their length is nevertheless more than sufficient in order to ensure a use and functionality as antisense RNA or double-stranded RNA. Preference is given to using as marker protein a plant endogenous MTR kinase. Further endogenous plant MTR kinases may readily be identified by means of screening databases or gene libraries using conserved, MTK kinase-typical motifs. Said motifs may be derived from Fig. 9a-b, for example. Such motifs may comprise, by way of example but not by limitation, the following sequences:

E(V/I)GDGN(L/I)N(L/Y/F)V(F/Y), preferably EVGDGNLN(Y/F)V(F/Y)  
KQALPY(V/I)RC  
SWPMT(R/K)ERAYF  
PEVYHFDRT  
GMRY(I/L)EPPHI  
CRLTEQVVFSDPY  
HGDLH(S/T)GS

Further suitable motifs may be derived from Fig. 9a-b without difficulty.

- 5 Particular preference is given to sequences according to Gen-Bank Acc. No: AF212863 or AC079674 (Protein\_ID=AAG51775.1). Preference is also given to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 38 or 40, in particular  
10 the sequences according to SEQ ID NO: 37 or 39.
- n) alcohol dehydrogenases (Adh), in particular plant Adh-1 gene products, with preference being given to using as nontoxic  
15 substance X substances such as allyl alcohol which is converted in this manner into the toxic substance (Y) acrolein. Corresponding sequences and the process of carrying out negative selection processes using, for example, allyl alcohol are known to the skilled worker and readily obtainable from the appropriate sequence database (e.g. GenBank) (Wisman E et  
20 al. (1991) Mol Gen Genet 226(1-2):120-8; Jacobs M et al. (1988) Biochem Genet 26(1-2):105-22; Schwartz D. (1981) Environ Health Perspect 37:75-7). The sequences, materials and processes disclosed in the context of said publications are hereby explicitly referred to.
- 25 Particular preference is given to sequences according to Gen-Bank Acc. No: X77943, M12196, AF172282, X04049 or AF253472. Preference is also given to nucleic acid sequences coding for polypeptides according to SEQ ID NO: 42, 44, 46 or 48, in  
30 particular the sequences according to SEQ ID NO: 41, 43, 45 or 47.
- (o) Further suitable negative selection markers are those sequences which exert per se a toxic action on plant cells,  
35 such as, for example, diphtheria toxin A, ribonucleases such as barnase and also ribosome-inhibiting proteins such as ricin. In this context, these proteins are preferably expressed in the plant cells inducibly rather than constitutively. The induction is preferably carried out chemically, it being possible,  
40 for example, to use the chemically inducible promoters mentioned below in order to ensure said chemically induced expression.
- 45 "Reduction" or "to reduce" is to be interpreted broadly in connection with a marker protein or with its amount, expression, activity and/or function and comprises the partial or essentially complete stopping or blocking, based on different cell-biological

## 21

mechanisms, of the functionality of a marker protein in a plant cell, plant or a part, tissue, organ, cells or seeds derived therefrom.

- 5 A reduction for the purpose of the invention also comprises a reduction of the amount of a marker protein down to an essentially complete lack of said marker protein (i.e. a lack of detectability of marker protein activity or marker protein function or a lack of immunological detectability of said marker protein). In  
10 this context, expression of a particular marker protein (or of its amount, expression, activity and/or function) in a cell or an organism is reduced preferably by more than 50%, particularly preferably by more than 80%, very particularly preferably by more than 90%, most preferably by more than 98%. Reduction means in  
15 particular also the complete lack of the marker protein (or of its amount, expression, activity and/or function). In this context, activity and/or function mean preferably the property of the marker protein of exerting a toxic effect on the plant cell or the plant organism and, respectively, the ability to convert  
20 the substance X into the substance Y. The toxic effect caused by the marker protein is reduced preferably by more than 50%, particularly preferably by more than 80%, very particularly preferably by more than 90%, most preferably by more than 98%. "Reduction" includes of course within the scope of the present  
25 invention also a complete, 100% reduction or removal of the marker protein (or of its amount, expression, activity and/or function) (for example by deleting the marker protein gene from the genome).
- 30 The invention comprises various strategies for reducing the expression, amount, activity and/or function of the marker protein. The skilled worker appreciates the fact that a number of various methods are available in order to influence the expression, amount, activity and/or function of a marker protein in the de-  
35 sired way. Examples which may be mentioned but which are not limiting are:
- a) introducing at least one marker protein double-stranded ribo-  
40 nucleic acid sequence (MP-dsRNA) or an expression cassette or expression cassettes ensuring expression thereof. Included are those processes in which the MP-dsRNA is directed against a marker protein gene (i.e. genomic DNA sequences such as promoter sequences) or a marker protein gene transcript (i.e. mRNA sequences).  
45

## 22

- 5 b) introducing at least one marker protein antisense ribonucleic acid sequence (MP-antisenseRNA) or an expression cassette ensuring expression thereof. Included are those processes in which the MP-antisenseRNA is directed against a marker protein gene (i.e. genomic DNA sequences) or a marker protein gene transcript (i.e. RNA sequences).  $\alpha$ -anomeric nucleic acid sequences are also included.
- 10 c) introducing at least one MP-antisenseRNA combined with a ribozyme or an expression cassette ensuring expression thereof
- 15 d) introducing at least one marker protein sense ribonucleic acid sequence (MP-senseRNA) for inducing a cosuppression or an expression cassette ensuring expression thereof
- 20 e) introducing at least one DNA- or protein-binding factor against a marker protein gene, marker protein RNA or marker protein or an expression cassette ensuring expression thereof
- 25 f) introducing at least one viral nucleic acid sequence causing degradation of the marker protein RNA or an expression cassette ensuring expression thereof
- 30 g) introducing at least one construct for generating a functional loss (e.g. generation of stop codons, shifts in the reading frame etc.) on a marker protein gene, for example by generating an insertion, deletion, inversion or mutation in a marker protein gene. Preferably, knockout mutants may be generated by means of targeted insertion into said marker protein gene via homologous recombination or by introducing sequence-specific nucleases against marker protein gene sequences.
- 35 It is known to the skilled worker that it is also possible to use other processes within the scope of the present invention in order to reduce a marker protein or its activity or function. For example, it may also be advantageous, depending on the type of the marker protein used, to introduce a dominant-negative variant of a marker protein or an expression cassette ensuring expression thereof. In this context, any single one of these processes may cause a reduction in the expression, amount, activity and/or function of a marker protein. A combined application is also conceivable. Further methods are known to the skilled worker and may
- 40 comprise hindering or stopping the processing of the marker protein, the transport of the marker protein or of its mRNA, the inhibition of ribosome attachment, the inhibition of RNA splicing,
- 45

## 23

the induction of an enzyme degrading marker protein RNA and/or the inhibition of translational elongation or termination.

The embodiments below will describe by way of example the individual preferred processes:

a) Introducing a double-stranded ribonucleic acid sequence of a marker protein (MP-dsRNA)

10 The process of gene regulation by means of double-stranded RNA ("double-stranded RNA interference"; dsRNAi) has been described many times for animal and plant organisms (e.g. Matzke MA et al. (2000) Plant Mol Biol 43:401-415; Fire A. et al (1998) Nature  
15 391:806-811; WO 99/32619; WO 99/53050; WO 00/68374; WO 00/44914; WO 00/44895; WO 00/49035; WO 00/63364). The processes and methods described in the references indicated are hereby explicitly referred to. dsRNAi processes are based on the phenomenon that simultaneously introducing the complementary strand and contour  
20 strand of a gene transcript suppresses expression of the corresponding gene in a highly efficient manner. Preferably, the phenotype caused is very similar to that of a corresponding knockout mutant (Waterhouse PM et al. (1998) Proc Natl Acad Sci USA 95:13959-64). The dsRNAi process has proved to be particularly  
25 efficient and advantageous in reducing marker protein expression.

Double-stranded RNA molecule means within the scope of the invention preferably one or more ribonucleic acid sequences which, owing to complementary sequences, are theoretically (e.g. according  
30 to the base pair rules by Watson and Crick) and/or actually (e.g. owing to hybridization experiments in vitro and/or in vivo) capable of forming double-stranded RNA structures. The skilled worker is aware of the fact that the formation of double-stranded RNA structures represents a state of equilibrium. Preferably, the ratio of double-stranded molecules to corresponding dissociated  
35 forms is at least 1 to 10, preferably 1:1, particularly preferably 5:1, most preferably 10:1.

The invention therefore further relates to double-stranded RNA  
40 molecules (dsRNA-Moleküle) which, when introduced into a plant organism (or into a cell, tissue, organ or propagation material derived therefrom) cause the reduction of at least one marker protein. The double-stranded RNA molecule for reducing expression of a marker protein (MP-dsRNA) here preferably comprises  
45

a) a "sense" RNA strand comprising at least one ribonucleotide sequence which is essentially identical to at least a part of

## 24

the "sense" RNA transcript of a nucleic acid sequence coding for a marker protein, and

- 5 b) an "antisense" RNA strand which is essentially, preferably fully, complementary to the RNA sense strand under a).

With respect to the dsRNA molecules, marker protein nucleic acid sequence preferably means a sequence according to SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45 or 47 or a functional equivalent thereof.

15 "Essentially identical" means that the dsRNA sequence may also have insertions, deletions and also individual point mutations in comparison with the marker protein target sequence and nevertheless causes an efficient reduction in expression. The homology (as defined hereinbelow) between the "sense" strand of an inhibitory dsRNA and at least one part of the "sense" RNA transcript of a nucleic acid sequence coding for a marker protein (or between the "antisense" strand of the complementary strand of a nucleic acid sequence coding for a marker protein) is preferably at least 20 75%, preferably at least 80%, very particularly preferably at least 90%, most preferably 100%.

25 A 100% sequence identity between dsRNA and a marker protein gene transcript is not absolutely necessary in order to cause an efficient reduction in marker protein expression. Consequently, the process is advantageously tolerant toward sequence deviations as may be present due to genetic mutations, polymorphisms or evolutionary divergences. Thus it is possible, for example, using the 30 dsRNA which has been generated starting from the marker protein sequence of the first organism, to suppress marker protein expression in a second organism. This is particularly advantageous when the marker protein used is a plant-intrinsic, endogenous marker protein (for example a 5-methylthioribose kinase or alcohol dehydrogenase). For this purpose, the dsRNA preferably includes 35 sequence regions of marker protein gene transcripts which correspond to conserved regions. Said conserved regions may be readily derived from sequence comparisons.

40 The length of the subsection is at least 10 bases, preferably at least 25 bases, particularly preferably at least 50 bases, very particularly preferably at least 100 bases, most preferably at least 200 bases or at least 300 bases.

45 Alternatively, an "essentially identical" dsRNA may also be defined as a nucleic acid sequence capable of hybridizing with part of a marker protein gene transcript (e.g. in 400 mM NaCl, 40 mM



## 25

PIPES pH 6.4, 1 mM EDTA at 50°C or 70°C for 12 to 16 h).

5 "Essentially complementary" means that the "antisense" RNA strand may also have insertions, deletions and also individual point mutations in comparison with the complement of this "sense" RNA strand. The homology between the "antisense" RNA strand and the complement of the "sense" RNA strand is preferably at least 80%, preferably at least 90%, very particularly preferably at least 95%, most preferably 100%.

10

15 "Part of the "sense" RNA transcript" of a nucleic acid sequence coding for a marker protein means fragments of an RNA or mRNA transcribed or transcribable from a nucleic acid sequence coding for a marker protein, preferably from a marker protein gene. In this context, the fragments have a sequence length of preferably at least 20 bases, preferably at least 50 bases, particularly preferably at least 100 bases, very particularly preferably at least 200 bases, most preferably at least 500 bases. The complete 20 transcribable RNA or mRNA is also included. Included are also sequences such as those which may be transcribed under artificial conditions from regions of a marker protein gene which are otherwise, under natural conditions, not transcribed, such as promoter regions, for example.

25

The dsRNA may consist of one or more strands of polyribonucleotides. Naturally, in order to achieve the same purpose, it is also possible to introduce a plurality of individual dsRNA molecules which comprise in each case one of the above-defined ribo- 30 nucleotide sequence sections into the cell or the organism. The double-stranded dsRNA structure may be formed starting from two complementary, separate RNA strands or, preferably, starting from a single, self-complementary RNA strand. In this case, the "sense" RNA strand and the "antisense" RNA strand are preferably 35 connected covalently to one another in the form of an inverted "repeat".

40 As described in WO 99/53050, for example, the dsRNA may also comprise a hairpin structure by connecting the "sense" and the "antisense" strands by a connecting sequence ("linker"; for example an intron). Preference is given to the self-complementary dsRNA structures, since they require only the expression of an RNA sequence and always comprise the complementary RNA strands in an 45 equimolar ratio. The connecting sequence may is preferably an intron (e.g. an intron of the potato ST-LS1 gene; Vancanneyt GF et al. (1990) Mol Gen Genet 220(2):245-250).

## 26

The nucleic acid sequence coding for a dsRNA may include further elements such as, for example, transcription termination signals or polyadenylation signals.

5 Bringing together, if intended, the two strands of the dsRNA in a cell or plant may be achieved by way of example in the following way:

- 10 a) transformation of the cell or plant with a vector comprising both expression cassettes,
- b) cotransformation of the cell or plant with two vectors, one of which comprises the expression cassettes containing the  
15 "sense" strand and the other one of which comprises the expression cassettes containing the "antisense" strand.

The formation of the RNA duplex may be initiated either outside or inside the cell.

20

The dsRNA may be synthesized either in vivo or in vitro. For this purpose, a DNA sequence coding for a dsRNA may be inserted into an expression cassette under the control of at least one genetic control element (such as a promoter, for example). A polyadenylation is not necessary and neither need any elements for initiating a translation be present. Preference is given to the expression cassette for the MP-dsRNA being present on the transformation construct or the transformation vector. For this purpose, the expression cassettes coding for the "antisense" strand and/or the "sense" strand of an MP-dsRNA or for the self-complementary strand of the dsRNA are preferably inserted into a transformation vector and introduced into the plant cell by using the processes described below. A stable insertion into the genome may be advantageous for the process of the invention but is not absolutely necessary. Since a dsRNA causes a long-term effect, transient expression is also sufficient in many cases. The dsRNA may also be part of the RNA to be expressed by the nucleic acid sequence to be inserted by fusing it, for example, to the 3'-untranslated part of said RNA.

40

The dsRNA may be introduced in an amount which makes possible at least one copy per cell. Higher amounts (e.g. at least 5, 10, 100, 500 or 1000 copies per cell) may, if appropriate, cause a more efficient reduction.

45

- b) Introducing an antisense ribonucleic acid sequence of a marker protein (MP-antisenseRNA)

## 27

Processes for reducing a particular protein by means of the "antisense" technique have been described multiple times, also in plants (Sheehy et al. (1988) Proc Natl Acad Sci USA 85: 8805-8809; US 4,801,340; Mol JN et al. (1990) FEBS Lett 268(2):427-430). The antisense nucleic acid molecule hybridizes or binds to the cellular mRNA and/or genomic DNA coding for the marker protein to be reduced, thereby suppressing transcription and/or translation of said marker protein. The hybridization may be produced in a conventional manner via the formation of a stable duplex or, in the case of genomic DNA, by binding of the antisense nucleic acid molecule to the duplex of the genomic DNA via specific interaction in the large groove of the DNA helix.

An MP-antisenseRNA may be derived using the nucleic acid sequence coding for this marker protein, for example the nucleic acid sequence according to SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45 or 47 according to the base pair rules by Watson and Crick. The MP-antisenseRNA may be complementary to the entire transcribed mRNA of the marker protein, may be limited to the coding region or may consist only of an oligonucleotide which is complementary to a part of the coding or noncoding sequence of the mRNA. Thus, for example, the oligonucleotide may be complementary to the region comprising the translation start site for the marker protein. The MP-antisenseRNA may be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length, but may also be longer and comprise at least 100, 200, 500, 1000, 2000 or 5000 nucleotides. MP-antisenseRNA are preferably expressed recombinantly in the target cell in the course of the process of the invention.

The MP-antisenseRNA may also be part of an RNA to be expressed by the nucleic acid sequence to be inserted by being fused, for example, to the 3'-untranslated part of said RNA.

The invention further relates to transgenic expression cassettes containing a nucleic acid sequence coding for at least part of a marker protein, with said nucleic acid sequence being functionally linked in antisense orientation to a promoter functional in plant organisms. Said expression cassettes may be part of a transformation construct or transformation vector or else may be introduced in the course of a cotransformation.

In a further preferred embodiment, expression of a marker protein may be inhibited by nucleotide sequences which are complementary to the regulatory region of a marker protein gene (e.g. a marker protein promoter and/or enhancer) and which form with the DNA double helix there triple-helical structures, thereby reducing transcription of the marker protein gene. Corresponding processes

have been described (Helene C (1991) Anticancer Drug Res 6(6):569-84; Helene C et al. (1992) Ann NY Acad Sci 660:27-36; Maher LJ (1992) Bioassays 14(12):807-815).

5 In a further embodiment, the MP-antisenseRNA may be an  $\alpha$ -anomeric nucleic acid. Such  $\alpha$ -anomeric nucleic acid molecules form with complementary RNA specific double-stranded hybrids in which, in contrast to the conventional  $\beta$ -nucleic acids, the two strands are oriented parallel to one another (Gautier C et al. (1987) Nucleic  
10 Acids Res 15:6625-6641).

c) Introducing an MP-antisenseRNA combined with a ribozyme

15 Advantageously, the above-described antisense strategy may be coupled to a ribozyme process. Catalytic RNA molecules or ribozymes may be adapted to any target RNA and cleave the phosphodiester backbone in specific positions, thereby functionally deactivating said target RNA (Tanner NK (1999) FEMS Microbiol Rev 23(3):257-275). In the process, the ribozyme is not modified it-  
20 self but is capable of cleaving in an analogous manner further target RNA molecules, thereby acquiring the properties of an enzyme. The incorporation of ribozyme sequences into "antisense" RNAs imparts specifically to these "antisense" RNAs this enzyme-like, RNA-cleaving property and thus increases their efficiency  
25 in inactivating the target RNA. The preparation and use of appropriate ribozyme "antisense" RNA molecules have been described (inter alia in Haseloff et al. (1988) Nature 334: 585-591); Haselhoff and Gerlach (1988) Nature 334:585-591; Steinecke P et al. (1992) EMBO J 11(4):1525- 1530; de Feyter R et al. (1996) Mol Gen  
30 Genet. 250(3):329-338).

In this way, it is possible to use ribozymes (e.g. hammerhead ribozymes; Haselhoff and Gerlach (1988) Nature 334:585-591) in order to catalytically cleave the mRNA of a marker protein to be  
35 reduced and thus prevent translation. The ribozyme technique may increase the efficiency of an antisense strategy. Processes for expressing ribozymes in order to reduce particular proteins have been described in (EP 0 291 533, EP 0 321 201, EP 0 360 257). Ribozyme expression has likewise been described in plant cells  
40 (Steinecke P et al. (1992) EMBO J 11(4):1525-1530; de Feyter R et al. (1996) Mol Gen Genet. 250(3):329-338). Suitable target sequences and ribozymes may be determined, for example, as described in "Steinecke P, Ribozymes, Methods in Cell Biology 50, Galbraith et al. eds, Academic Press, Inc. (1995), pp. 449-460",  
45 by calculating the secondary structures of ribozyme RNA and target RNA and by the interaction thereof (Bayley CC et al. (1992) Plant Mol Biol. 18(2):353-361; Lloyd AM and Davis RW et al. (1994) Mol Gen Genet. 242(6):653-657). It is possible, for exam-

## 29

ple, to construct derivatives of the Tetrahymena L-19 IVS RNA which have regions complementary to the mRNA of the marker protein to be suppressed (see also US 4,987,071 and US 5,116,742). Alternatively, such ribozymes may also be identified via a selection process from a library of various ribozymes (Bartel D and Szostak JW (1993) Science 261:1411-1418).

d) Introducing a sense ribonucleic acid sequence of a marker protein (MP-senseRNA) for inducing a cosuppression

10

Expression of a marker protein ribonucleic acid sequence (or a part thereof) in sense orientation may result in a cosuppression of the corresponding marker protein gene. Expression of sense RNA with homology to an endogenous marker protein gene may reduce or switch off expression of the latter, as has been described similarly for antisense approaches (Jorgensen et al. (1996) Plant Mol Biol 31(5):957-973; Goring et al. (1991) Proc Natl Acad Sci USA 88:1770-1774; Smith et al. (1990) Mol Gen Genet 224:447-481; Napoli et al. (1990) Plant Cell 2:279-289; Van der Krol et al. (1990) Plant Cell 2:291-99). In this context, the introduced construct may represent completely or only partially the homologous gene to be reduced. The possibility of translation is not required. The application of this technique to plants has been described (e.g. Napoli et al. (1990) Plant Cell 2:279-289; in US 5,034,323.

The cosuppression is preferably carried out using a sequence which is essentially identical to at least part of the nucleic acid sequence coding for a marker protein, for example the nucleic acid sequence according to SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45 or 47.

The MP-senseRNA is preferably chosen in such a way that a translation of the marker protein or a part thereof cannot occur. For this purpose, for example, the 5'-untranslated or 3'-untranslated region may be chosen or else the ATG start codon may be deleted or mutated.

40

e) Introducing DNA- or protein-binding factors against marker protein genes, marker protein RNAs or proteins

Marker protein expression may also be reduced using specific DNA-binding factors, for example factors of the zinc finger transcription factor type. These factors attach to the genomic sequence of the endogenous target gene, preferably in the

45

## 30

regulatory regions, and cause a reduction in expression. Appropriate processes for preparing corresponding factors have been described (Dreier B et al. (2001) J Biol Chem 276(31):29466-78; Dreier B et al. (2000) J Mol Biol 303(4):489-502; Beerli RR et al. (2000) Proc Natl Acad Sci USA 97 (4):1495-1500; Beerli RR et al. (2000) J Biol Chem 275(42):32617-32627; Segal DJ and Barbas CF 3rd. (2000) Curr Opin Chem Biol 4(1):34-39; Kang JS and Kim JS (2000) J Biol Chem 275(12):8742-8748; Beerli RR et al. (1998) Proc Natl Acad Sci USA 95(25):14628-14633; Kim JS et al. (1997) Proc Natl Acad Sci USA 94(8):3616-3620; Klug A (1999) J Mol Biol 293(2):215-218; Tsai SY et al. (1998) Adv Drug Deliv Rev 30(1-3):23-31; Mapp AK et al. (2000) Proc Natl Acad Sci USA 97(8):3930-3935; Sharrocks AD et al. (1997) Int J Biochem Cell Biol 29(12):1371-1387; Zhang L et al. (2000) J Biol Chem 275(43):33850-33860).

These factors may be selected using any segment of a marker protein gene. This section is preferably in the region of the promoter region. However, for gene suppression, it may also be in the region of the coding exons or introns.

It is also possible to introduce factors which inhibit the marker protein itself into a cell. These protein-binding factors may be, for example, aptamers (Famulok M and Mayer G (1999) Curr Top Microbiol Immunol 243:123-36) or antibodies or antibody fragments or single-chain antibodies. Obtaining these factors has been described (Owen M et al. (1992) Biotechnology (N Y) 10(7):790-794; Franken E et al. (1997) Curr Opin Biotechnol 8(4):411-416; Whitelam (1996) Trend Plant Sci 1:286-272).

30

f) Introducing viral nucleic acid sequences and expression constructs causing the degradation of marker protein RNA

Marker protein expression may also be effectively implemented by inducing the specific degradation of marker protein RNA by the plant with the aid of a viral expression system (Amplikon; Angell SM et al. (1999) Plant J 20(3):357-362). These systems, also referred to as "VIGS" (viral induced gene silencing), introduce nucleic acid sequences with homology to the transcript of a marker protein to be reduced into the plant by means of viral vectors. Transcription is then switched off, presumably mediated by plant defence mechanisms against viruses. Appropriate techniques and processes have been described (Ratcliff F et al. (2001) Plant J 25(2):237-45; Fagard M und Vaucheret H (2000) Plant Mol Biol 43(2-3):285-93; Anandalakshmi R et al. (1998) Proc Natl Acad Sci USA 95(22):13079-84; Ruiz MT (1998) Plant Cell 10(6):937-46).

## 31

VIGS-mediated reduction is preferably implemented using a sequence which is essentially identical to at least part of the nucleic acid sequence coding for a marker protein, for example the nucleic acid sequence according to SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45 or 47.

- g) Introducing constructs for generating a functional loss or a functional reduction of marker protein genes

The skilled worker knows numerous possible processes of how to modify genomic sequences in a targeted manner. These include, in particular, processes such as the generation of knockout mutants by means of targeted homologous recombination, for example by generating stop codons, shifts in the reading frame etc. (Hohn B and Puchta H (1999) Proc Natl Acad Sci USA 96:8321-8323) or the targeted deletion or inversion of sequences by means of, for example, sequence-specific recombinases or nucleases (see below).

20

In a preferred embodiment, the marker protein gene is inactivated by introducing a sequence-specific recombinase. Thus it is possible, for example, for the marker protein gene to include recognition sequences for sequence-specific recombinases or to be flanked by such sequences, and introducing the recombinase then deletes or inverts particular sequences of the marker protein gene, thus leading to inactivation of the marker protein gene. A corresponding procedure is depicted diagrammatically in Fig. 1.

30

Appropriate processes for deletion/inversion of sequences by means of sequence-specific recombinase systems are known to the skilled worker. Examples which may be mentioned are the Cre/lox system of bacteriophage P1 (Dale EC and Ow DW (1991) Proc Natl Acad Sci USA 88:10558-10562; Russell SH et al. (1992) Mol Gen Genet 234:49-59; Osborne BI et al. (1995) Plant J 7:687-701), the yeast FLP/FRT system (Kilby NJ et al. (1995) Plant J 8:637-652; Lyznik LA et al. (1996) Nucl Acids Res 24:3784-3789), the Gin recombinase of the Mu phage, the E. coli Pin recombinase and the R/RS system of the pSR1 plasmids (Onouchi H et al. (1995) Mol Gen Genet 247:653-660; Sugita K et al. (2000) Plant J. 22:461-469). In these systems, the recombinase (for example Cre or FLP) interacts specifically with its particular recombination sequences (34 bp lox-Sequenz and, respectively, 47 bp FRT sequence). Preference is given to the bacteriophage P1 Cre/lox and the yeast FLP/FRT systems. The FLP/FRT and cre/lox recombinase systems have already been applied in plant systems (Odell et al. (1990) Mol Gen Genet 223:369-378). Preference is given to introducing the recombinase

## 32

by means of recombinant expression starting from an expression cassette included on a DNA construct.

5 The activity or amount of the marker protein may also be reduced  
by a targeted deletion in the marker protein gene, for example by  
sequence-specific induction of DNA double-strand breaks at a rec-  
ognition sequence for specific induction of DNA double-strand  
breaks in or close to the nucleic acid sequence coding for a  
10 marker protein. In its simplest embodiment (cf. Fig. 2, A and B)  
an enzyme is to this end introduced with the transformation  
construct, which generates at least one double-strand break in  
such a way that the resulting illegitimate recombination or dele-  
tion causes a reduction in the activity or amount of marker pro-  
tein, for example by inducing a shift in the reading frame or  
15 deletion of essential sequences.

The efficiency of this approach may be increased by the sequence  
coding for the marker protein being flanked by sequences (A and,  
20 respectively, A') which have a sufficient length and homology to  
one another in order to recombine with one another as a conse-  
quence of the induced double-strand break and thus to cause, due  
to an intramolecular homologous recombination, a deletion of the  
sequence coding for the marker protein. Fig. 3 depicts diagram-  
25 matically a corresponding procedure in an exemplary embodiment of  
this variant.

The amount, function and/or activity of the marker protein may  
also be reduced by a targeted insertion of nucleic acid sequences  
30 (for example of the nucleic acid sequence to be inserted within  
the scope of the process of the invention) into the sequence cod-  
ing for a marker protein (e.g. by means of intermolecular homolo-  
gous recombination). This embodiment of the process of the inven-  
35 tion is particularly advantageous and preferred, since, in  
addition to the general advantages of the process of the inven-  
tion, it makes it moreover also possible to insert the nucleic  
acid sequence to be inserted into the plant genome in a reproduc-  
ible, predictable, location-specific manner. This avoids the  
40 positional effects which otherwise occur in the course of a ran-  
dom, location-unspecific insertion (and which may manifest them-  
selves, for example, in the form of different levels of expres-  
sion of the transgene or in unintended inactivation of endogenous  
genes). Preference is given to using as an "anti-marker protein"  
45 compound in the course of this embodiment a DNA construct which  
comprises at least part of the sequence of a marker protein gene  
or neighbouring sequences and which can thus specifically recom-  
bine with said sequences in the target cell so that a deletion,



## 33

addition or substitution of at least one nucleotide alters the marker protein gene in such a way that the functionality of said marker protein gene is reduced or completely removed. The alteration may also affect the regulatory elements (e.g. the promoter) of the marker protein gene so that the coding sequence remains unaltered, but expression (transcription and/or translation) does not occur and is reduced. In conventional homologous recombination, the sequence to be inserted is flanked at its 5' and/or 3' end by further nucleic acid sequences (A' and, respectively, B') which have a sufficient length and homology to corresponding sequences of the marker protein gene (A and, respectively, B) for making homologous recombination possible. The length is usually in a range from several hundred bases to several kilobases (Thomas KR and Capecchi MR (1987) Cell 51:503; Strepp et al. (1998) Proc Natl Acad Sci USA 95(8):4368-4373). The homologous recombination is carried out by transforming the plant cell containing the recombination construct by using the process described below and selecting successfully recombined clones based on the subsequently inactivated marker protein. Although homologous recombination is a relatively rare event in plant organisms, a selection pressure may be avoided by recombination into the marker protein gene, allowing a selection of the recombined cells and sufficient efficiency of the process. Fig. 4 diagrammatically depicts a corresponding procedure in an exemplary embodiment of this variant.

In an advantageous embodiment of the invention, however, insertion into the marker protein gene is facilitated by means of further functional elements. The term is to be understood as being comprehensive and means the use of sequences or of transcripts or polypeptides derived therefrom which are capable of increasing the efficiency of the specific integration into a marker protein gene. Various processes are available to the skilled worker for this purpose. However, preference is given to implementing the insertion by inducing a sequence-specific double-strand break in or close to the marker protein gene.

In a preferred embodiment of the invention, the marker protein is inactivated (i.e. the amount, expression, activity or function is reduced) by integrating a DNA sequence into a marker protein gene, with the process preferably comprising the following steps:

- i) introducing an insertion construct and at least one enzyme suitable for inducing DNA double-strand breaks at a recognition sequence for targeted induction of DNA double-strand

## 34

breaks in or close to the marker protein gene, and

5 ii) inducing DNA double-strand breaks at the recognition sequences for targeted induction of DNA double-strand breaks in or close to the marker protein gene, and

10 iii) inserting the insertion construct into the marker protein gene, with the functionality of the marker protein gene and, preferably, the functionality of the recognition sequence for targeted induction of DNA double-strand breaks is inactivated so that the enzyme suitable for induction of DNA double-strand breaks can no longer cut said recognition sequence, and

15 iv) selecting plants or plant cells in which the insertion construct has been inserted into the marker protein gene.

20 The insertion construct, preferably, comprises the nucleic acid sequence to be inserted into the genome but may also be used separately therefrom.

25 "Enzyme suitable for inducing DNA double-strand breaks at the recognition sequence for targeted induction of DNA double-strand breaks" ("DSBI enzyme" for "double-strand-break inducing enzyme" hereinbelow) means generally all those enzymes which are capable of generating sequence-specifically double-strand breaks in double-stranded DNA. Examples which may be mentioned but which are not limiting are: .

30 1. Restriction endonucleases, preferably type II restriction endonucleases, particularly preferably Homing endonucleases as described in detail hereinbelow.

35 2. Artificial nucleases as described in detail hereinbelow, such as, for example, chimeric nucleases, mutated restriction or Homing endonucleases or RNA protein particles derived from group II mobile introns.

40 Both natural and artificially prepared DSBI enzymes are suitable. Preference is given to all of those DSBI enzymes whose recognition sequence is known and which can either be obtained in the form of their proteins (for example by purification) or be expressed using their nucleic acid sequence.

45

Preference is given to selecting the DSBI enzyme, with the knowledge of its specific recognition sequence, in such a way that it

## 35

possesses, apart from the target recognition sequence, no further functional recognition regions in the genome of the target plant. Very particular preference is therefore given to Homing endonucleases (overview: Belfort M and Roberts RJ (1997) *Nucleic Acids Res* 25:3379-3388; Jasin M (1996) *Trends Genet* 12:224-228; Internet: <http://rebase.neb.com/rebase/rebase.homing.html>; Roberts RJ and Macelis D (2001) *Nucl Acids Res* 29: 268-269). The latter full said requirement, owing to their long recognition sequences. The sequences coding for Homing endonucleases of this kind may be isolated, for example, from the *Chlamydomonas* chromoplast genome (Turmel M et al. (1993) *J Mol Biol* 232:446-467). Suitable Homing endonucleases are listed under the abovementioned internet address. Examples of Homing endonucleases which may be mentioned are those like F-SceI, F-SceII, F-SuvI, F-TevI, F-TevII, I-AmalI, I-AniI, I-CeuI, I-CeuAIIP, I-ChuI, I-CmoI, I-CpaI, I-CpaII, I-CreI, I-CrepsbIP, I-CrepsbIIP, I-CrepsbIIIP, I-CrepsbIVP, I-CsmI, I-CvuI, I-CvuAIP, I-DdiII, I-DirI, I-DmoI, I-HspNIP, I-LlaI, I-MsoI, I-NaaI, I-NanI, I-NclIP, I-NgrIP, I-NitI, I-NjaI, I-Nsp236IP, I-PakI, I-PboIP, I-PcuIP, I-PcuAI, I-PcuVI, I-PgrIP, I-PobIP, I-PorI, I-PorIIP, I-PpbIP, I-PpoI, I-SPBetaIP, I-ScaI, I-SceI, I-SceII, I-SceIII, I-SceIV, I-SceV, I-SceVI, I-SceVII, I-SexIP, I-SneIP, I-SpomCP, I-SpomIP, I-SpomIIP, I-SquIP, I-Ssp6803I, I-SthPhiJP, I-SthPhiST3P, I-SthPhiS3bP, I-TdeIP, I-TevI, I-TevII, I-TevIII, I-UarAP, I-UarHGPA1P, I-UarHGPA13P, I-VinIP, I-ZbiIP, PI-MtuI, PI-MtuHIP, PI-MtuHIIP, PI-PfuI, PI-PfuII, PI-PkoI, PI-PkoII, PI-PspI, PI-Rma43812IP, PI-SPBetaIP, PI-SceI, PI-TfuI, PI-TfuII, PI-ThyI, PI-TliI, PI-TliII. Preference is given here to those Homing endonucleases whose gene sequences are already known, such as, for example, F-SceI, I-CeuI, I-ChuI, I-DmoI, I-CpaI, I-CpaII, I-CreI, I-CsmI, F-TevI, F-TevII, I-TevI, I-TevII, I-AniI, I-CvuI, I-LlaI, I-NanI, I-MsoI, I-NitI, I-NjaI, I-PakI, I-PorI, I-PpoI, I-ScaI, I-Ssp6803I, PI-PkoI, PI-PkoII, PI-PspI, PI-TfuI, PI-TliI.

35 Very particular preference is given to

- I-CeuI (Cote MJ and Turmel M (1995) *Curr Genet* 27:177-183.; Gauthier A et al. (1991) *Curr Genet* 19:43-47; Marshall (1991) *Gene* 104:241-245; GenBank Acc. No.: Z17234 nucleotides 5102 to 5758),
- I-ChuI (Cote V et al. (1993) *Gene* 129:69-76; GenBank Acc. No.: L06107, nucleotides 419 to 1075),

## 36

- I-CmoeI (Drouin M et al. (2000) Nucl Acids Res 28:4566-4572),
- 5 - I-CpaI from *Chlamydomonas pallidostigmatica* (GenBank Acc. No.: L36830, nucleotides 357 to 815; Turmel M et al. (1995) Nucleic Acids Res 23:2519-2525; Turmel, M et al. (1995) Mol Biol Evol 12:533-545)
- 10 - I-CpaII (Turmel M et al. (1995) Mol Biol Evol 12:533-545; GenBank Acc. No.: L39865, nucleotides 719 to 1423),
- I-CreI (Wang J et al. (1997) Nucleic Acids Res 25: 3767-3776; Dürrenberger, F and Rochaix JD (1991) EMBO J 10:3495-3501; GenBank Acc. No.: X01977, nucleotides 571 to 1062),
- 15 - I-CsmI (Ma DP et al. (1992) Plant Mol Biol 18:1001-1004)
- I-NanI (Elde M et al. (1999) Eur J Biochem. 259:281-288; GenBank Acc. No.: X78280, nucleotides 418 to 1155),
- 20 - I-NitI (GenBank Acc. No.: X78277, nucleotides 426 to 1163),
- I-NjaI (GenBank Acc. No.: X78279, nucleotides 416 to 1153),
- 25 - I-PpoI (Muscarella DE and Vogt VM (1989) Cell 56:443-454; Lin J and Vogt VM (1998) Mol Cell Biol 18:5809-5817; GenBank Acc. No.: M38131, nucleotides 86 to 577),
- 30 - I-PspI (GenBank Acc. No.: U00707, nucleotides 1839 to 3449),
- I-ScaI (Monteilhet C et al. (2000) Nucleic Acids Res 28: 1245-1251; GenBank Acc. No.: X95974, nucleotides 55 to 465)
- 35 - I-SceI (WO 96/14408; US 5,962,327, therein Seq ID NO: 1),
- Endo SceI (Kawasaki et al. (1991) J Biol Chem 266:5342-5347, identical to F-SceI; GenBank Acc. No.: M63839, nucleotides 159 to 1589),
- 40 - I-SceII (Sarguiel B et al. (1990) Nucleic Acids Res 18:5659-5665),
- 45 - I-SceIII (Sarguiel B et al. (1991) Mol Gen Genet. 255:340-341),

## 37

- I-Ssp6803I (GenBank Acc. No.: D64003, nucleotides 35372 to 35824),
- I-TevI (Chu et al. (1990) Proc Natl Acad Sci USA 87:3574-3578; Bell-Pedersen et al. (1990) Nucleic Acids Res 18:3763-3770; GenBank Acc. No.: AF158101, nucleotides 144431 to 143694),
- 10 - I-TevII (Bell-Pedersen et al. (1990) Nucleic Acids Res 18:3763-3770; GenBank Acc. No.: AF158101, nucleotides 45612 to 44836),
- I-TevIII (Eddy et al. (1991) Genes Dev. 5:1032-1041).

15

Very particular preference is given to commercially available Homing endonucleases such as I-CeuI, I-SceI, I-PpoI, PI-PspI or PI-SceI. Most preference is given to I-SceI and I-PpoI. While the gene coding for I-PpoI may be utilized in its natural form, the gene coding for I-SceI possesses an editing site. Since, in contrast to yeast mitochondria, the appropriate editing is not carried out in higher plants, an artificial sequence encoding the I-SceI protein must be used for heterologous expression of this enzyme (US 5,866,361).

25

The enzymes may be purified from their source organisms in the manner familiar to the skilled worker and/or the nucleic acid sequence encoding said enzymes may be cloned. The sequences of various enzymes have been deposited with GenBank (see above).

30

Artificial DSB1 enzymes which may be mentioned by way of example are chimeric nucleases which are composed of an unspecific nuclease domain and a sequence-specific DNA-binding domain (e.g. consisting of zinc fingers) (Smith J et al. (2000) Nucl Acids Res 28(17):3361-3369; Bibikova M et al. (2001) Mol Cell Biol. 21:289-297). Thus, for example, the catalytic domain of the restriction endonuclease FokI has been fused to zinc finger-binding domains, thereby defining the specificity of the endonuclease (Chandrasegaran S & Smith J (1999) Biol Chem 380:841-848; Kim YG & Chandrasegaran S (1994) Proc Natl Acad Sci USA 91:883-887; Kim YG et al. (1996) Proc Natl Acad Sci USA 93:1156-1160). The described technique has also been used previously for imparting a predefined specificity to the catalytic domain of the yeast Ho endonuclease by fusing said domain to the zinc finger domain of transcription factors (Nahon E & Raveh D (1998) Nucl Acids Res 26:1233-1239). It is possible, using suitable mutation and selection processes, to adapt existing Homing endonucleases to any de-

sired recognition sequence.

As mentioned, zinc finger proteins are particularly suitable as DNA-binding domains within chimeric nucleases. These DNA-binding zinc finger domains may be adapted to any DNA sequence. Appropriate processes for preparing corresponding zinc finger domains have been described and are known to the skilled worker (Beerli RR et al. (2000) *Proc Natl Acad Sci* 97(4):1495-1500; Beerli RR et al. (2000) *J Biol Chem* 275(42):32617-32627; Segal DJ and Barbas CF 3rd. (2000) *Curr Opin Chem Biol* 4(1):34-39; Kang JS and Kim JS (2000) *J Biol Chem* 275(12):8742-8748; Beerli RR et al. (1998) *Proc Natl Acad Sci USA* 95(25):14628-14633; Kim JS et al. (1997) *Proc Natl Acad Sci USA* 94(8):3616-3620; Klug A (1999) *J Mol Biol* 293(2):215-218; Tsai SY et al. (1998) *Adv Drug Deliv Rev* 30(1-3):23-31; Mapp AK et al. (2000) *Proc Natl Acad Sci USA* 97(8):3930-3935; Sharrocks AD et al. (1997) *Int J Biochem Cell Biol* 29(12):1371-1387; Zhang L et al. (2000) *J Biol Chem* 275(43):33850-33860). Processes for preparing and selecting zinc finger DNA-binding domains with high sequence specificity have been described (WO 96/06166, WO 98/53059, WO 98/53057). Fusing a DNA-binding domain obtained in this way to the catalytic domain of an endonuclease (such as, for example, the FokI or Ho endonuclease) enables chimeric nucleases to be prepared which have any desired specificity and which may be used as DSB1 enzymes advantageously within the scope of the present invention.

Artificial DSB1 enzymes with altered sequence specificity may also be generated by mutating already known restriction endonucleases or Homing endonucleases, using methods familiar to the skilled worker. Besides the mutagenesis of Homing endonucleases, the mutagenesis of maturases is of particular interest for the purpose of obtaining an altered substrate specificity. Maturases frequently share many features with Homing endonucleases and, if appropriate, can be converted into nucleases by carrying out few mutations. This has been shown, for example, for the maturase in the bakers' yeast *bi2* intron. Only two mutations in the maturase-encoding open reading frame (ORF) sufficed to impart to this enzyme a Homing-endonuclease activity (Szczepanek & Lazowska (1996) *EMBO J* 15:3758-3767).

Further artificial nucleases may be generated with the aid of mobile group II introns and the proteins encoded by them, or parts of these proteins. Mobile group II introns, together with the proteins encoded by them, form RNA-protein particles which are capable of recognizing and cutting DNA in a sequence-specific manner. In this context, the sequence specificity can be adapted

## 39

to the requirements by mutating particular regions of the intron (see below) (WO 97/10362).

Preference is given to expressing the DSBI enzyme as a fusion  
 5 protein with a nuclear localization sequence (NLS). This NLS sequence enables facilitated transport into the nucleus and increases the efficiency of the recombination system. Various NLS sequences are known to the skilled worker and described, inter alia, in Jicks GR and Raikhel NV (1995) Annu. Rev. Cell Biol.  
 10 11:155-188. For example, the NLS sequence of the SV40 large antigen is preferred for plant organisms. Very particular preference is given to the following NLS sequences:

NLS1: N-Pro-Lys-Thr-Lys-Arg-Lys-Val-C  
 15  
 NLS2: N-Pro-Lys-Lys-Lys-Arg-Lys-Val-C

Owing to the small size of many DSBI enzymes (such as, for example, the Homing endonucleases), an NLS sequence is not absolutely  
 20 necessary, however. These enzymes are able to pass through the nuclear pores also without this assistance.

"Recognition sequence for targeted induction of DNA double-strand  
 25 breaks" means in general those sequences which allow recognition and cleavage by the DSBI enzyme under the conditions in the eukaryotic cell or organism used in this case. In this context, mention is made, by way of example but not by limitation, in table 1 below of the recognition sequences for the particular DSBI enzymes listed.  
 30

Table 1: Recognition sequences and source organisms of DSBI enzymes ("^" indicates the cleavage site of the DSBI enzyme within a recognition sequence)

35

DSBI enzyme	Source organism	Recognition sequence
40 CRE	Bacteriophage P1	5'-AACTCTCATCGCTTCGGATAACTTCCTGTTATCCGAAACAT ATCACTCACTTTGGTGATTTCACCGTAACTGTCTATGATTAATG -3'
FLP	Saccharomyces cerevisiae	5'-GAAGTTCCTATTCCGAAGTTCCTATTCTCTAGAAAGTA- TAGGAACTTC-3'
45 R	pSR1 plasmids	5'-CGAGATCATATCACTGTGGACGTTGATGAAAGAATACGTTA TTCTTTCATCAAATCGT

## 40

	P-element transposase	Drosophila	5'-CTAGATGAAATAACATAAAGGTGG
5	I-AniI	Aspergillus nidulans	5'-TTGAGGAGGTT^TCTCTGTAAATAANNNNNNNNNNNNNNN 3'-AACTCCTCCAAAGAGACATTTATTNNNNNNNNNNNNNNNN^
	I-DdiI	Dictyostelium discoideumAX3	5'-TTTTTTGGTCATCCAGAAGTATAT 3'-AAAAAACCAG^TAGGTCTTCATATA
10	I-CvuI	Chlorella vulgaris	5'-CTGGGTTCAAAACGTCGTGA^GACAGTTTGG 3'-GACCCAAGTTTTGCAG^CACTCTGTCAAACC
	I-CsmI	Chlamydomonas smithii	5'-GTACTAGCATGGGGTCAAATGTCTTTCTGG
15	I-CmoEI	Chlamydomonas moewusii	5'-TCGTAGCAGCT^CACGGTT 3'-AGCATCG^TCGAGTGCCAA
	I-CreI	Chlamydomonas reinhardtii	5'-CTGGGTTCAAAACGTCGTGA^GACAGTTTGG 3'-GACCCAAGTTTTGCAG^CACTCTGTCAAACC
20	I-ChuI	Chlamydomonas humicola	5'-GAAGGTTTGGCACCTCG^ATGTCGGCTCATC 3'-CTTCCAAACCGTG^GAGCTACAGCCGAGTAG
	I-CpaI	Chlamydomonas pallidostigmatica	5'-CGATCCTAAGGTAGCGAA^ATTCA 3'-GCTAGGATTCCATC^GCTTTAAGT
25	I-CpaII	Chlamydomonas pallidostigmatica	5'-CCCGGCTAACTC^TGTGCCAG 3'-GGGCCGAT^TGAGACACGGTC
30	I-CeuI	Chlamydomonas eugametos	5'-CGTAACTATAACGGTCCTAA^GGTAGCGAA 3'-GCATTGATATTGCCAG^GATTCCATCGCTT
	I-DmoI	Desulfurococcus mobilis	5'-ATGCCTTGCCGGGTAA^GTTCCGGCGCGCAT 3'-TACGGAACGGCC^CATTCAAGGCCGCGCGTA
35	I-SceI	S.cerevisiae	5'-AGTTACGCTAGGGATAA^CAGGGTAATATAG 3'-TCAATGCGATCCC^TATTGTCCCATTATATC 5'-TAGGGATAA^CAGGGTAAT 3'-ATCCC^TATTGTCCCATTAT ("Core" sequence)
40	I-SceII	S.cerevisiae	5'-TTTTGATTCTTTGGTCAACC^TGAAGTATA 3'-AAACTAAGAAACCAG^TGGGACTTCATAT
	I-SceIII	S.cerevisiae	5'-ATTGGAGGTTTTGGTAAC^TATTTATTACC 3'-TAACCTCCAAACC^ATTGATAAATAATGG
45	I-SceIV	S.cerevisiae	5'-TCTTTTCTCTTGATTA^GCCCTAATCTACG 3'-AGAAAAGAGAAC^TAATCGGGATTAGATGC



	I-SceV	S.cerevisiae	5'-ATAATTTCTTCTTAGTAATGCC 3'-TTATTAAAAGAAGATCATTA^CGG
5	I-SceVI	S.cerevisiae	5'-GTTATTTAATG^TTTTAGTAGTTGG 3'-CAATAAATTACAAATCATCA^ACC
	I-SceVII	S.cerevisiae	5'-TGTCACATTGAGGTGC ACTAGTTATTAC
10	PI-SceI	S.cerevisiae	5'-ATCTATGTCGGGTGC^GGAGAAAGAGGTAAT 3'-TAGATACAGCC^CACGCCCTCTTTCTCCATTA
	F-SceI	S.cerevisiae	5'-GATGCTGTAGGC^ATAGGCTTG GTT 3'-CTACGACA^TCCGTATCCGAACCAA
15	F-SceII	S.cerevisiae	5'-CTTTC CGCAACA^GT AAAATT 3'-GAAAGGCG^TTGTCATTTTAA
	I-HmuI	Bacillus subtilis bacteriophage SPO1	5'-AGTAATGAGCCTAACGCTCAGCAA 3'-TCATTACTCGGATTGC^GAGTCGTT
20	I-HmuII	Bacillus subtilis bacteriophage SP82	5'-AGTAATGAGCCTAACGCTCAACAANNNNNNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNNNNNNNNNN
25	I-LlaI	Lactococcus lactis	5'-CACATCCATAAC^CATATCATTTTT 3'-GTGTAGGTATTGGTATAGTAA^AAA
	I-MsoI	Monomastix species	5'-CTGGGTTCAAACGTCGTGA^GACAGTTTGG 3'-GACCCAAGTTTTCAG^CACTCTGTCAAACC
30	I-NanI	Naegleria andersoni	5'-AAGTCTGGTGCCA^GCACCCGC 3'-TTCAGACC^ACGGTCGTGGGCG
	I-NitI	Naegleria italica	5'-AAGTCTGGTGCCA^GCACCCGC 3'-TTCAGACC^ACGGTCGTGGGCG
35	I-NjaI	Naegleria jamiesoni	5'-AAGTCTGGTGCCA^GCACCCGC 3'-TTCAGACC^ACGGTCGTGGGCG
	I-PakI	Pseudendoclonium akinetum	5'-CTGGGTTCAAACGTCGTGA^GACAGTTTGG 3'-GACCCAAGTTTTCAG^CACTCTGTCAAACC
40	I-PosI	Pyrobaculum organotrophum	5'-GCGAGCCCCGTAAGGGT^GTGTACGGG 3'-CGCTCGGGCATT^CCCACACATGCC
	I-PpoI	Physarum polycephalum	5'-TAAGTATGACTCTCTTAA^GGTAGCCAAAT 3'-ATTGATACTGAGAG^AATTCCATCGGTTA
45	I-ScaI	Saccharomyces capensis	5'-TGTCACATTGAGGTGCACT^AGTTATTAC 3'-ACAGTGTAAC TCCAC^GTGATCAATAATG

## 42

	I-Ssp6803I	Synechocystis species	5'-GTCGGGCT^CATAACCCGAA 3'-CAGCCCGAGTA^TTGGGCTT
5	PI-PfuI	Pyrococcus furiosus Vcl	5'-GAAGATGGGAGGAGGG^ACCGGACTCAACTT 3'-CTTCTACCCTCC^TCCCTGGCCTGAGTTGAA
	PI-PfuII	Pyrococcus furiosus Vcl	5'-ACGAATCCATGTGGAGA^AGAGCCTCTATA 3'-TGCTTAGGTACAC^CTCTTCTCGGAGATAT
10	PI-PkoI	Pyrococcus kodakaraensis KOD1	5'-GATTTTAGAT^CCCTGTACC 3'-CTAAAA^TCTAGGGACATGG
	PI-PkoII	Pyrococcus kodakaraensis KOD1	5'-CAGTACTACG^GTTAC 3'-GTCATG^ATGCCAATG
15	PI-PspI	Pyrococcus sp.	5'-AAAATCCTGGCAAACAGCTATTAT^GGGTAT 3'-TTTTAGGACCGTTTGTGAT^AATACCCATA
20	PI-TfuI	Thermococcus fumicolans ST557	5'-TAGATTTTAGGT^CGCTATATCCTTCC 3'-ATCTAAAA^TCCAGCGATATAGGAAGG
	PI-TfuII	Thermococcus fumicolans ST557	5'-TAYGCNGAYACN^GACGGYTTYT 3'-ATRCGNCT^RTGNCTGCCRAARA
25	PI-ThyI	Thermococcus hydrothermalis	5'-TAYGCNGAYACN^GACGGYTTYT 3'-ATRCGNCT^RTGNCTGCCRAARA
30	PI-TliI	Thermococcus litoralis	5'-TAYGCNGAYACNGACG^YTTYT 3'-ATRCGNCTRTGNC^TGCCRAARA
	PI-TliII	Thermococcus litoralis	5'-AAATTGCTTGCAAACAGCTATTACGGCTAT
35	I-TevI	Bacteriophage T4	5'-AGTGGTATCAAC^GCTCAGTAGATG 3'-TCACCATAGT^TGCGAGTCATCTAC
	I-TevII	Bacteriophage T4	5'-GCTTATGAGTATGAAGTGAACACGT^TATTC 3'-CGAATAC^TCATACTTCACTTGTG^CAATAAG
40	F-TevI	Bacteriophage T4	5'-GAAACACAAGA^AATGTTAGTAAANNNNNNNNNNNNN 3'-CTTTGTGTTCTTTACAAATCATTTNNNNNNNNNNNNNN^
	F-TevII	Bacteriophage T4	5'-TTTAATCCTCGCTTC^AGATATGGCAACTG 3'-AAATTAGGAGCGA^AGTCTATACCGTTGAC

45

Relatively small deviations (degenerations) of the recognition sequence which nevertheless make possible recognition and cleavage by the particular DSBII enzyme are also included here. Such

## 43

deviations, also in connection with different basic conditions such as, for example, calcium or magnesium concentration, have been described (Argast GM et al. (1998) J Mol Biol 280:345-353). Core sequences of these recognition sequences are also included.

- 5 It is known that the inner portions of the recognition sequences also suffice for an induced double-strand break and that the outer portions are not necessarily relevant but may contribute to determining the cleavage efficiency. Thus, for example, an 18bp core sequence can be defined for I-SceI.

10

Said DSBI recognition sequences may be localized in various positions in or close to a marker protein gene and, for example when the marker protein used is a transgene, may already be incorporated when constructing the marker protein expression cassette.

- 15 Various possible localizations are illustrated by way of example in Figs. 2-A, 2-B, 3 and 5 and in the descriptions thereof.

- In a further advantageous embodiment, the insertion sequence comprises at least one homology sequence A which has a sufficient length and a sufficient homology to a sequence A' in the marker protein gene in order to ensure homologous recombination between A and A'. The insertion sequence is preferably flanked by two sequences A and B which have a sufficient length and a sufficient homology to a sequence A' and, respectively, B' in the marker protein gene in order to ensure homologous recombination between A and A' and, respectively, B and B'.

- "Sufficient length" means, with respect to the homology sequences A, A' and B, B', preferably sequences with a length of at least 100 base pairs, preferably at least 250 base pairs, particularly preferably at least 500 base pairs, very particularly preferably at least 1000 base pairs, most preferably of at least 2500 base pairs.

35

- "Sufficient homology" means, with respect to the homology sequences, preferably sequences whose homology to one another is at least 70%, preferably 80%, preferentially at least 90%, particularly preferably at least 95%, very particularly preferably at least 99%, most preferably 100%, over a length of at least 20 base pairs, preferably at least 50 base pairs, particularly preferably at least 100 base pairs, very particularly preferably at least 250 base pairs, most preferably at least 500 base pairs.

45

Homology between two nucleic acids means the identity of the nucleic acid sequence over in each case the entire sequence length, which identity is calculated by way of comparison with the aid of

10 a further preferred embodiment, the recombination efficiency is increased by a combination with processes which promote homologous recombination. Such systems have been described and comprise, by way of example, expression of proteins such as RecA or treatment with PARP inhibitors. It has been demonstrated that the

15 intrachromosomal homologous recombination in tobacco plants can be increased by using PARP inhibitors (Puchta H et al. (1995) Plant J 7:203-210). The use of these inhibitors can further increase the rate of homologous recombination in the recombinant constructs, after inducing the sequence-specific DNA double-strand break, and thus the efficiency of the deletion of the

20 transgene sequences. Various PARP inhibitors may be used here. Preference is given to including inhibitors such as 3-amino benzamide, 8-hydroxy-2-methylquinazolin-4-one (NU1025), 1,11b-dihydro-[2H]benzopyrano[4,3,2-de]isoquinolin-3-one (GPI 6150),

25 5-aminoisoquinolinone, 3,4-dihydro-5-[4-(1-piperidinyl)butoxy]-1(2H)-isoquinolinone or the substances described in WO 00/26192, WO 00/29384, WO 00/32579, WO 00/64878, WO 00/68206, WO 00/67734, WO 01/23386 and WO 01/23390.

30 Further suitable methods are the introduction of nonsense mutations into endogenous marker protein genes, for example by means of introducing RNA/DNA oligonucleotides into the plant (Zhu et al. (2000) Nat Biotechnol 18(5):555-558). Point mutations may also be generated by means of DNA-RNA hybrids which are also

35 known as "chimeraplasty" (Cole-Strauss et al. (1999) Nucl Acids Res 27(5):1323-1330; Kmiec (1999) Gene therapy American Scientist 87(3):240-247).

40 The methods of dsRNAi, cosuppression by means of sense RNA and VIGS (virus induced gene silencing) are also referred to as post-transcriptional gene silencing (PTGS). PTGS processes are particularly advantageous because the demands on the homology between the marker protein gene to be reduced and the transgenically expressed sense or dsRNA nucleic acid sequence are lower than, for

45 example, in the case of a traditional antisense approach. Thus it is possible, using the marker protein nucleic acid sequences from one species, to effectively reduce also expression of homologous

## 45

marker protein proteins in other species, without it being absolutely necessary to isolate and to elucidate the structure of the marker protein homologues occurring there. Considerably less labor is therefore required.

5

"Introduction" comprises within the scope of the invention any processes which are suitable for introducing an "anti-marker protein" compound, directly or indirectly, into a plant or a cell, compartment, tissue, organ or seeds of said plant or generating  
10 said compound there. The introduction may result in a transient presence of an "anti-marker protein" compound (for example a dsRNA or a recombinase) or else in a permanent (stable) presence.

15 According to the different nature of the approaches described above, the "anti-marker protein" compound may exert its function directly (for example by way of insertion into an endogenous marker protein gene). However, said function may also be exerted indirectly after transcription into an RNA (for example in anti-sense approaches) or after transcription and translation into a  
20 protein (for example in the case of recombinases or DSBI enzymes). The invention comprises both directly and indirectly acting "anti-marker protein" compounds.

25 Introducing comprises, for example, processes such as transfection, transduction or transformation.

"Anti-marker protein" compounds thus comprises, for example, also expression cassettes capable of implementing expression (i.e.  
30 transcription and, if appropriate, translation) of, for example, an MP-dsRNA, an MP-antisenseRNA, a sequence-specific recombinase or a DSBI enzyme in a plant cell.

"Expression cassette" means within the scope of the present invention generally those constructions in which a nucleic acid sequence to be expressed is functionally linked to at least one genetic control sequence, preferably a promoter sequence.  
Expression cassettes preferably consist of double-stranded DNA and may have a linear or circular structure.  
40

A functional linkage means, for example, the sequential arrangement of a promoter with a nucleic acid sequence to be transcribed (for example coding for an MP-dsRNA or a DSBI enzyme) and, if appropriate, further regulatory elements such as, for example, a  
45 terminator and/or polyadenylation signals in such a way that each of the regulatory elements can fulfill its function during transcription of the nucleic acid sequence, depending on the arrange-

## 46

ment of the nucleic acid sequences. In this context, function can mean, for example, the control of expression, i.e. transcription and/or translation, of the nucleic acid sequence (e.g. coding for an MP-dsRNA or a DSBI enzyme). In this context, control comprises, for example, initiating, increasing, controlling or suppressing the expression, i.e. transcription and, if appropriate, translation. This does not necessarily require a direct linkage in the chemical sense. Genetic control sequences such as, for example, enhancer sequences, may exert their function on the target sequence also from positions further afar or even from different DNA molecules. Preference is given to arrangements in which the nucleic acid sequence to be transcribed is positioned downstream of the sequence acting as promoter so that both sequences are covalently connected to one another. The distance between the promoter sequence and the nucleic acid sequence to be expressed transgenically is here preferably less than 200 base pairs, particularly preferably less than 100 base pairs, very particularly preferably less than 50 base pairs.

The skilled worker knows various ways of obtaining any of the expression cassettes of the invention. An expression cassette of the invention is prepared, for example, preferably by direct fusion of a nucleic acid sequence acting as promoter to a nucleotide sequence to be expressed (e.g. coding for an MP-dsRNA or a DSBI enzyme). A functional linkage may be produced by means of common recombination and cloning techniques, as are described, for example, in Maniatis T, Fritsch EF and Sambrook J (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and in Silhavy TJ et al. (1984) Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and in Ausubel FM et al. (1987) Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley Interscience.

The expression cassettes of the invention preferably comprise a promoter 5' upstream of the particular nucleic acid sequence to be expressed transgenically and a terminator sequence as an additional genetic control sequence 3' downstream and also, if appropriate, further customary regulatory elements, in each case functionally linked to the nucleic acid sequence to be expressed transgenically.

The term "genetic control sequences" is to be understood broadly and means all those sequences which have an influence on the making or function of the expression cassette of the invention. For example, genetic control sequences ensure transcription and, if

## 47

appropriate, translation in prokaryotic or eukaryotic organisms. Genetic control sequences are described, for example, in "Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990)" or "Gruber and Crosby, in: 5 Methods in Plant Molecular Biology and Biotechnology, CRC Press, Boca Raton, Florida, eds.: Glick and Thompson, Chapter 7, 89-108" and in the references quoted there.

10 Genetic control sequences comprise, in particular in plants, functional promoters. Preferred promoters suitable for the expression cassettes are in principle any promoters capable of controlling expression of genes, in particular foreign genes, in plants.

15 Plant-specific promoters or promoters functional in plants or in a plant cell means in principle any promoter capable of controlling expression of genes, in particular foreign genes, in at least one plant or one part, cell, tissue, culture of a plant. In 20 this context, expression may be, for example, constitutive, inducible or development-dependent. Preference is given to:

a) Constitutive promoters

25 "Constitutive" promoters means those promoters which ensure expression in numerous, preferably all, tissues over a relatively large period of plant development, preferably at all points in time of plant development (Benfey et al. (1989) EMBO J 8:2195-2202). Preference is given in particular to using a 30 plant promoter or a promoter which is derived from a plant virus. Particular preference is given to the promoter of the 35S transcript of the CaMV cauliflower mosaic virus (Franck et al. (1980) Cell 21:285-294; Odell et al. (1985) Nature 313:810-812; Shewmaker et al. (1985) Virology 140:281-288; 35 Gardner et al. (1986) Plant Mol Biol 6:221-228) or the 19S CaMV promoter (US 5,352,605; WO 84/02913; Benfey et al. (1989) EMBO J 8:2195-2202) and also to the promoter of the Arabidopsis thaliana nitrilase-1 gene (GenBank Acc. No.: Y07648, nucleotides 2456 (alternatively 2861) to 4308 or 40 alternatively 4340 or 4344. (e.g. bp 2456 to 4340).

Another suitable constitutive promoter is the rubisco small subunit (SSU) promoter (US 4,962,028), the leguminB promoter 45 (GenBank Acc. No.: X03677), the promoter of the Agrobacterium nopaline synthase, the TR dual promoter, the Agrobacterium OCS (octopine synthase) promoter, the ubiquitin promoter (Holtorf S et al. (1995) Plant Mol Biol 29:637-649), the ubi-

## 48

quitin 1 promoter (Christensen et al. (1992) Plant Mol Biol 18:675-689; Bruce et al. (1989) Proc Natl Acad Sci USA 86:9692-9696), the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits or the promoter of a proline-rich protein from wheat (WO 91/13991), and further promoters of genes whose constitutive expression in plants is known to the skilled worker.

10 b) Tissue-specific promoters

Preference is given to promoters with specificities for the anthers, ovaries, flowers, leaves, stems, roots or seeds.

15

Seed-specific promoters comprise, for example, the promoter of phaseolin (US 5,504,200; Bustos MM et al. (1989) Plant Cell 1(9):839-53), of the 2S albumin (Joseffson LG et al. (1987) J Biol Chem 262:12196-12201), of legumin (Shirsat A et al. (1989) Mol Gen Genet 215(2): 326-331), of USP (unknown seed protein; Bäumlein H et al. (1991) Mol Gen Genet 225(3):459-67), of napin (US 5,608,152; Stalberg K et al. (1996) L Planta 199:515-519), of the sucrose-binding protein (WO 00/26388), of legumin B4 (LeB4; Bäumlein H et al. (1991) Mol Gen Genet 225: 121-128; Baeumlein et al. (1992) Plant Journal 2(2):233-9; Fiedler U et al. (1995) Biotechnology (NY) 13(10):1090f), of oleosin (WO 98/45461) or of Bce4 (WO 91/13980). Further suitable seed-specific promoters are those of the genes coding for the high molecular weight glutenin (HMWG), gliadin, branching enzyme, ADP glucose pyrophosphatase (AGPase) or starch synthase. Preference is further given to promoters which allow seed-specific expression in monocotyledones such as corn, barley, wheat, rye, rice, etc. promoters which may be employed advantageously are the promoter of the lpt2 or lpt1 gene (WO 95/15389, WO 95/23230) and the promoters described in WO 99/16890 (hordein, glutelin, oryzin, prolamin, gliadin, zein, kasirin or secalin promoters). Further seed-specific promoters are described in WO 89/03887.

Tuber-, storage-root- or root-specific promoters comprise, for example, the class I patatin promoter (B33) or the promoter of the potato cathepsin D inhibitor.

45

Leaf-specific promoters comprise, for example, the promoter of the potato cytosolic FBPase (WO 97/05900), the



## 49

SSU promoter (small subunit) of rubisco (ribulose-1,5-bisphosphate carboxylase) or the potato ST-LSI promoter (Stockhaus et al. (1989) EMBO J 8:2445-2451).

5 Flower-specific promoters comprise, for example, the phytoene synthase promoter (WO 92/16635) or the promoter of the P-rr gene (WO 98/22593).

10 - Anther-specific promoters comprise, for example, the 5126 promoter (US 5,689,049, US 5,689,051), the glob-1 promoter and the  $\gamma$ -zein promoter.

c) Chemically inducible promoters

15 Chemically inducible promoters allow expression control as a function of an exogenous stimulus (review article: Gatz et al. (1997) Ann Rev Plant Physiol Plant Mol Biol 48:89-108). Examples which may be mentioned are: the PRP1 promoter (Ward  
20 et al. (1993) Plant Mol Biol 22:361-366), a salicylic acid-inducible promoter (WO 95/19443), a benzenesulfonamide-inducible promoter (EP-A 0 388 186), a tetracycline-inducible promoter (Gatz et al. (1992) Plant J 2:397-404), an abscisic acid-inducible promoter (EP 0 335 528) and an ethanol- or  
25 cyclohexanone-inducible promoter (WO 93/21334). Also suitable is the promoter of the glutathione S-transferase isoform II gene (GST-II-27), which may be activated by exogenously applied safeners such as, for example, N,N-diallyl-2,2-dichloroacetamide (WO 93/01294) and which is functional in numerous  
30 tissues of both monocotyledones and dicotyledones.

Particular preference is given to constitutive or inducible promoters.

35 Preference is further given to plastid-specific promoters for targeted expression in the plastids. Suitable promoters are described, for example, in WO 98/55595 or WO 97/06250. promoters which may be mentioned here are the rpo B promoter element, the atoB promoter element, the clpP promoter element (see also WO  
40 99/46394) and the 16SrDNA promoter element. Viral promoters are also suitable (WO 95/16783).

45 Targeted expression in plastids may also be achieved by using, for example, a bacterial or bacteriophage promoter, introducing the resulting expression cassette into the plastid DNA and then expressing expression by means of a fusion protein of a bacterial or bacteriophage polymerase and a plastid transit peptide. US

## 50

5,925,806 describes an appropriate process.

Genetic control sequences further comprise also the 5'-untranslated regions, introns or noncoding 3' region of genes, such as, 5 for example, the actin-1 intron, or the Adh1-S introns 1, 2 and 6 (general overview: The Maize Handbook, Chapter 116, Freeling and Walbot, Eds., Springer, New York (1994)). These sequences have been shown to be able to play a significant functions in the regulation of gene expression. Thus it has been demonstrated that 10 5'-untranslated sequences may increase transient expression of heterologous genes. They may further promote tissue specificity (Rouster J et al.(1998) Plant J. 15:435-440). As an example of translation enhancers, mention may be made of the 5' leader sequence of the tobacco mosaic virus (Gallie et al. (1987) Nucl 15 Acids Res 15:8693-8711).

Polyadenylation signals suitable as control sequences are in particular polyadenylation signals of plant genes and also *Agrobacterium tumefaciens* T-DNA polyadenylation signals. Examples of 20 particularly suitable terminator sequences are the OCS (octopine synthase) terminator and the NOS (nopaline synthase) terminator (Depicker A et al (1982) J Mol Appl Genet 1:561-573) and also the terminators of soybean actin, RUBISCO or alpha-amylase from wheat 25 (Baulcombe DC et al (1987) Mol Gen Genet 209:33-40).

Advantageously, the expression cassette may contain one or more "enhancer sequences" functionally linked to the promoter, which make increased transgenic expression of the nucleic acid sequence 30 possible.

Genetic control sequences further means sequences coding for fusion proteins consisting of a signal peptide sequence. The expression of a target gene is possible in any desired cell 35 compartment, such as, for example, the endomembrane system, the vacuole and the chloroplasts. Desired glycosylation reactions, in particular foldings, and the like are possible by utilizing the secretory pathway. Secretion of the target protein to the cell surface or secretion into the culture medium, for example when 40 using suspension-cultured cells or protoplasts, is also possible. The target sequences required for this may both be taken into account in individual vector variations and be introduced into the vector together with the target gene to be cloned by using a 45 suitable cloning strategy. Target sequences which may be used are both endogenous, if present, and heterologous sequences. Additional heterologous sequences which are preferred for functional linkage but not limited thereto are further targeting sequences

## 51

for ensuring subcellular localization in the apoplast, in the vacuole, in plastids, in the mitochondrion, in the endoplasmic reticulum (ER), in the nucleus, in elaioplasts or other compartments; and also translation enhancers such as the 5' leader sequence from tobacco mosaic virus (Gallie et al. (1987) Nucl Acids Res 15: 8693-8711) and the like. The process of transporting proteins which are per se not located in the plastids specifically into said plastids has been described (Klosgen RB and Weil JH (1991) Mol Gen Genet 225(2):297-304; Van Breusegem F et al. (1998) Plant Mol Biol 38(3):491-496).

Control sequences are furthermore understood to be those which make possible a homologous recombination or insertion into the genome of a host organism or allow the removal from the genome. Methods such as the cre/lox technique allow the expression cassette to be removed tissue-specifically, possibly inducibly from the genome of the host organism (Sauer B. Methods. 1998; 14(4):381-92). Here, particular flanking sequences are attached to the target gene (lox sequences), which make subsequent removal by means of the cre recombinase possible.

Preferably, the expression cassette, consisting of a linkage of the promoter to the nucleic acid sequence to be transcribed, may have been integrated into a vector and may be transferred into the plant cell or organism, for example, by transformation, according to any of the processes described below.

"Transgenic" means preferably, for example with respect to a transgenic expression cassette, a transgenic expression vector, a transgenic organism or to processes for transgenic expression of nucleic acids, all constructions brought about by genetic engineering methods or processes using said constructions, in which either

- a) the nucleic acid sequence to be expressed, or
- b) the promoter functionally linked to the nucleic acid sequence to be expressed according to a), or
- c) (a) and (b)

are not located in their natural, genetic environment (i.e. at their natural chromosomal locus) or have been modified by genetic engineering methods, the modification possibly being, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. Natural genetic environment

## 52

means the natural chromosomal locus in the source organism or the presence in a genomic library.

5 "Transgenic" means, with respect to expression ("transgenic expression"), preferably all expressions achieved using a transgenic expression cassette, transgenic expression vector or transgenic organism, according to the definitions indicated above.

10 The DNA constructs employed within the scope of the process of the invention and the vectors derived therefrom may contain further functional elements. The term functional element is to be understood broadly and means all of those elements which influence the preparation, propagation or function of the DNA  
15 constructs or of vectors or organisms derived therefrom. Examples which may be mentioned without being limited thereto are:

1. Selection markers

20 Selection markers comprise, for example, those nucleic acid or protein sequences whose expression gives to a cell, tissue or organism an advantage (positive selection marker) or disadvantage (negative selection marker) over cells which do not express said nucleic acid or protein. Positive selection markers act, for example,  
25 by detoxifying a substance acting on the cell in an inhibitory manner (e.g. resistance to antibiotics/herbicides) or by forming a substance which enables the plant to regenerate better or grow more under the chosen conditions (for example nutritive markers, hormone-producing markers such as ipt; see below).  
30 Another type of positive selection marker comprises mutated proteins or RNAs which are not sensitive to a selective agent (e.g. 16S rRNA mutants which are insensitive to spectinomycin). Negative selection markers act, for example, by catalyzing the formation of a toxic substance in the transformed cells (e.g. the codA  
35 gene).

1.1 Positive selection markers:

40 In order to further increase the efficiency, the DNA constructs may comprise additional positive selection markers. In a preferred embodiment, the process of the invention may thus be carried out in the form of a dual selection in which a sequence coding for a resistance to at least one toxin, antibiotic or  
45 herbicide is introduced together with the nucleic acid sequence to be inserted and selection is carried out additionally by using the toxin, antibiotic or herbicide.

## 53

Appropriate proteins and sequences of positive selection markers and also selection processes are familiar to the skilled worker. The selection marker imparts to the successfully transformed cells a resistance to a biocide (e.g. a herbicide such as phosphinothricin, glyphosate or bromoxynil), a metabolism inhibitor such as 2-deoxyglucose 6-phosphate (WO 98/45456) or an antibiotic such as, for example, tetracycline, ampicillin, kanamycin, G 418, neomycin, bleomycin or hygromycin. Selection markers which may be mentioned by way of example are:

10

- phosphinothricin acetyltransferases (PAT) which acetylate the free amino group of the glutamine synthase inhibitor phosphinothricin (PPT) and thus detoxify PPT (de Block et al. (1987) EMBO J 6:2513-2518) (also referred to as Bialaphos® resistance gene (bar)). Corresponding sequences are known to the skilled worker (from *Streptomyces hygroscopicus* GenBank Acc. No.: X17220 and X05822, from *Streptomyces viridochromogenes* GenBank Acc. No.: M 22827 and X65195; US 5,489,520). Furthermore, synthetic genes have been described for expression in plastids. A synthetic PAT gene is described in Becker et al. (1994) Plant J 5:299-307. The genes impart a resistance to the herbicide Bialaphos or glufosinate and are frequently used markers in transgenic plants (Vickers JE et al. (1996) Plant Mol Biol Reporter 14:363-368; Thompson CJ et al. (1987) EMBO J 6:2519-2523).

- 5-enolpyruvylshikimate 3-phosphate synthases (EPSPS) which impart a resistance to glyphosate (N-(phosphonomethyl) glycine). The molecular target of the unselective herbicide glyphosate is 5-enolpyruvyl-3-phosphoshikimate synthase (EPSPS). This enzyme has a key function in the biosynthesis of aromatic amino acids in microbes and plants but not in mammals (Steinrücken HC et al. (1980) Biochem Biophys Res Commun 94:1207-1212; Levin JG and Sprinson DB (1964) J Biol Chem 239:1142-1150; Cole DJ (1985) Mode of action of glyphosate a literature analysis, p. 48-74. In: Grossbard E and Atkinson D (eds.). The herbicide glyphosate. Butterworths, Boston.). Preference is given to using glyphosate-tolerant EPSPS variants as selection markers (Padgett SR et al. (1996). New weed control opportunities: development of soybeans with a Roundup Ready™ gene. In: Herbicide Resistant Crops (Duke, S.O., ed.), pp. 53-84. CRC Press, Boca Raton, FL; Saroha MK and Malik VS (1998) J Plant Biochemistry and Biotechnology 7:65-72). The EPSPS gene of *Agrobacterium* sp. strain CP4 has a natural tolerance for glyphosate, which can be transferred to appropriate transgenic plants. The CP4 EPSPS gene was cloned from *Agrobacterium* sp. strain CP4 (Pad-

## 54

- gette SR et al. (1995) Crop Science 35(5):1451-1461). Sequences of EPSPS enzymes which are glyphosate-tolerant have been described (inter alia in US 5,510,471; US 5,776,760; US 5,864,425; US 5,633,435; US 5,627,061; US 5,463,175; EP 0 218 571). Further sequences are described under GenBank Acc. No: X63374 or M10947.
- 5
- Glyphosat<sup>®</sup>-degrading enzymes (gox gene; glyphosate oxidoreductase). GOX (for example *Achromobacter* sp. glyphosate oxidoreductase) catalyzes the cleavage of a C-N bond in glyphosate which is thus converted to aminomethylphosphonic acid (AMPA) and glyoxylate. GOX can thereby impart a resistance to glyphosate (Padgett et al. (1996) J Nutr 126(3):702-16; Shah D et al. (1986) Science 233:478-481).
  - 10
  - 15
  - The deh gene encodes a dehalogenase which inactivates Dalapon<sup>®</sup> (GenBank Acc. No.: AX022822, AX022820 and WO 99/27116)
  - 20
  - The bxn genes encode bromoxynil-degrading nitrilase enzymes (Genbank Acc. No: E01313 and J03196).
  - Neomycin phosphotransferases impart a resistance to antibiotics (aminoglycosides) such as neomycin, G418, hygromycin, paromomycin or kanamycin by reducing the inhibiting action of said antibiotics by means of a phosphorylation reaction. Particular preference is given to the nptII gene. Sequences can be obtained from GenBank (AF080390; AF080389). Moreover, the gene is already part of numerous expression vectors and can be isolated therefrom using processes familiar to the skilled worker (AF234316; AF234315; AF234314). The NPTII gene encodes an aminoglycoside 3'-O-phosphotransferase from *E. coli*, Tn5 (GenBank Acc. No: U00004 position 1401-2300; Beck et al. (1982) Gene 19 327-336).
  - 25
  - 30
  - 35
  - 40
  - 45
  - The DOGR1 gene was isolated from the yeast *Saccharomyces cerevisiae* (EP-A 0 807 836) and encodes a 2-deoxyglucose 6-phosphate phosphatase which imparts a resistance to 2-DOG (Randez-Gil et al. (1995) Yeast 11:1233-1240; Sanz et al. (1994) Yeast 10:1195-1202, GenBank Acc. No.: NC001140; position 194799-194056).
  - Acetolactate synthases which impart a resistance to imidazolinone/sulfonylurea herbicides (GenBank Acc. No.: X51514; Sathasivan K et al. (1990) Nucleic Acids Res. 18(8):2188; AB049823; AF094326; X07645; X07644; A19547; A19546; A19545;

55

I05376; I05373; AL133315)

- Hygromycin phosphotransferases (e.g. GenBank Acc. No.: X74325) which impart a resistance to the antibiotic hygromycin. The gene is part of numerous expression vectors and may be isolated therefrom using processes familiar to the skilled worker (such as, for example, polymerase chain reaction) (GenBank Acc. No.: AF294981; AF234301; AF234300; AF234299; AF234298; AF354046; AF354045).
- Genes of resistance to
  - a) Chloramphenicol (chloramphenicol acetyltransferase),
  - b) tetracycline (inter alia GenBank Acc. No.: X65876; X51366). Moreover, the gene is already part of numerous expression vectors and may be isolated therefrom using processes familiar to the skilled worker (such as, for example, polymerase chain reaction)
  - c) Streptomycin (inter alia GenBank Acc. No.: AJ278607).
  - d) Zeocin, the corresponding resistance gene is part of numerous cloning vectors (e.g. GenBank Acc. No.: L36849) and may be isolated therefrom using processes familiar to the skilled worker (such as, for example, polymerase chain reaction).
  - e) Ampicillin ( $\beta$ -lactamase gene; Datta N, Richmond MH (1966) Biochem J 98(1):204-9; Heffron F et al (1975) J. Bacteriol 122: 250-256; Bolivar F et al. (1977) Gene 2:95-114). The sequence is part of numerous cloning vectors and may be isolated therefrom using processes familiar to the skilled worker (such as, for example, polymerase chain reaction).

Genes such as isopentenyl transferase from *Agrobacterium tumefaciens* (strain:PO22) (Genbank Acc. No.: AB025109) may also be used as selection markers. The *ipt* gene is a key enzyme of cytokinin biosynthesis. Its overexpression facilitates the regeneration of plants (e.g. selection on cytokinin-free medium). The process for utilizing the *ipt* gene has been described (Ebinuma H et al. (2000) Proc Natl Acad Sci USA 94:2117-2121; Ebinuma H et al. (2000) Selection of Marker-free transgenic plants using the onco-genes (*ipt*, *rol* A, B, C) of *Agrobacterium* as selectable markers, In Molecular Biology of Woody Plants. Kluwer Academic Publish-

ers).

Various other positive selection markers which impart to the transformed plants a growth advantage over untransformed plants and also processes for their use are described, inter alia, in EP-A 0 601 092. Examples which may be mentioned are  $\beta$ -glucuronidase (in connection with cytokinin glucuronide, for example), mannose 6-phosphate isomerase (in connection with mannose), UDP-galactose 4-epimerase (in connection with galactose, for example).

For a selection marker functional in plastids, particular preference is given to those which impart a resistance to spectinomycin, streptomycin, kanamycin, lincomycin, gentamycin, hygromycin, methotrexat, bleomycin, phleomycin, blasticidin, sulfonamide, phosphinothricin, chlorsulfuron, bromoxymil, glyphosate, 2,4-datrazine, 4-methyltryptophan, nitrate, S-aminoethyl-L-cysteine, lysine/threonine, aminoethyl-cysteine or betainealdehyde. Particular preference is given to the genes aadA, nptII, BADH, FLARE-S (a fusion of aadA and GFP, described in Khan MS & Maliga P (1999) Nature Biotech 17:910-915). Especially suitable is the aadA gene (Svab Z and Maliga P (1993) Proc Natl Acad Sci USA 90:913-917). Modified 16S rDNA and also betainealdehyde dehydrogenase (BADH) from spinach have also been described (Daniell H et al. (2001) Trends Plant Science 6:237-239; Daniell H et al. (2001) Curr Genet 39:109-116; WO 01/64023; WO 01/64024; WO 01/64850). Lethal agents such as, for example, glyphosate may also be utilized in connection with correspondingly detoxifying or resistance enzymes (WO 01/81605).

The concentrations of the antibiotics, herbicides, biocides or toxins, which are used in each case for selection, must be adapted to the particular test conditions or organisms. Examples which may be mentioned for plants are kanamycin (Km) 50 mg/L, hygromycin B 40 mg/L, phosphinothricin (Ppt) 6 mg/L, spectinomycin (Spec) 500 mg/L.

## 2. Reporter genes

Reporter genes code for readily quantifiable proteins and thus ensure, via intrinsic color or enzyme activity, an evaluation of the transformation efficiency and of the location or time of expression. In this context, very particular preference is given to genes coding for reporter proteins (see also Schenborn E, Groskreutz D (1999) Mol Biotechnol 13(1):29-44) such as



## 57

- green fluorescence protein (GFP) (Chui WL et al. (1996) Curr Biol 6:325-330; Leffell SM et al. (1997) Biotechniques 23(5):912-8; Sheen et al. (1995) Plant J 8(5):777-784; Haseloff et al. (1997) Proc Natl Acad Sci USA 94(6): 2122-2127; Reichel et al. (1996) Proc Natl Acad Sci USA 93(12):5888-5893; Tian et al. (1997) Plant Cell Rep 16:267-271; WO 97/41228)
- chloramphenicol transferase
- luciferase (Millar et al. (1992) Plant Mol Biol Rep 10: 324-414; Ow et al. (1986) Science 234:856-859); allows bioluminescence detection
- $\beta$ -galactosidase (encodes an enzyme for which various chromogenic substrates are available)
- $\beta$ -glucuronidase (GUS) (Jefferson et al. (1987) EMBO J 6: 3901-3907) or the uidA gene (encode enzymes for which various chromogenic substrates are available)
- R-locus gene product which regulates production of anthocyanin pigments (red color) in plant tissue and thus makes possible a direct analysis of the promoter activity without addition of additional auxiliary substances or chromogenic substrates (Dellaporta et al. (1988) In: Chromosome Structure and Function: Impact of New Concepts, 18<sup>th</sup> Stadler Genetics Symposium, 11:263-282)
- tyrosinase (Katz et al. (1983) J Gen Microbiol 129:2703-2714), enzyme which oxidizes tyrosine to give DOPA and dopaquinone which consequently form the readily detectable melanine.
- aequorin (Prasher et al. (1985) Biochem Biophys Res Commun 126(3):1259-1268), may be used in calcium-sensitive bioluminescence detection.
- 3. Origins of replication which ensure propagation of the expression cassettes or vectors of the invention, for example in E. coli. Examples which may be mentioned are ORI (origin of DNA replication), the pBR322 ori or the P15A ori (Sambrook et al.: Molecular Cloning. A Laboratory Manual, 2<sup>nd</sup> ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

## 58

4. Elements, for example border sequences, which enable agrobacteria-mediated transfer into plant cells for transfer and integration into the plant genome, such as, for example, the right or left border of T-DNA or the vir region.
- 5
5. Multiple cloning regions (MCS) allow and facilitate the insertion of one or more nucleic acid sequences.
- 10 Nucleic acid sequences (e.g. expression cassettes) may be introduced into a plant organism or cells, tissues, organs, parts or seeds thereof by advantageously using vectors which contain said sequences. Vectors may be, by way of example, plasmids, cos-
- 15 mids, phages, viruses or else agrobacteria. The sequences may be inserted into the vector (preferably a plasmid vector) via suitable restriction cleavage sites. The resulting vector may first be introduced into E. coli and amplified. Correctly transformed E. coli are selected, grown and the recombinant vector is obtained using methods familiar to the skilled worker. Restriction
- 20 analysis and sequencing may serve to check the cloning step. Preference is given to those vectors which make possible a stable integration into the host genome.

The preparation of a transformed organism (or a transformed cell or tissue) requires that the corresponding DNA (e.g. the transformation vector) or RNA is introduced into the corresponding host cell. For this process which is referred to as transformation (or transduction or transfection), a multiplicity of methods and vectors are available (Keown et al. (1990) Methods in En-

30 zymology 185:527-537; Plant Molecular Biology and Biotechnology (CRC Press, Boca Raton, Florida), Chapter 6/7, pp. 71-119 (1993); White FF (1993) Vectors for Gene Transfer in Higher Plants; in: Transgenic Plants, Vol. 1, Engineering and Utilization, Editors: Kung and Wu R, Academic Press, 15-38; Jenes B et al. (1993) Tech-

35 niques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, editors: Kung and R. Wu, Academic Press, pp.128-143; Potrykus (1991) Annu Rev Plant Physiol Plant Molec Biol 42:205-225; Halford NG, Shewry PR (2000) Br Med Bull 56(1):62-73).

40

For example, the DNA or RNA may be introduced directly by micro-injection (WO 92/09696, WO 94/00583, EP-A 0 331 083, EP-A 0 175 966) or by bombardment with DNA or RNA-coded microparticles (biolistic processes using the gene gun "particle bombardment";

45 US 5,100,792; EP-A 0 444 882; EP-A 0 434 616; Fromm ME et al. (1990) Bio/Technology 8(9):833-9; Gordon-Kamm et al. (1990) Plant Cell 2:603). The cell may also be permeabilized chemically, for

## 59

example with polyethylene glycol, so as to enable the DNA to reach the cell by means of diffusion. The DNA may also take place by means of protoplast fusion to other DNA-containing units such as minicells, cells, lysosomes or liposomes (Freeman et al.

- 5 (1984) Plant Cell Physiol. 29:1353ff; US 4,536,475). Electroporation is another suitable method for introducing DNA, in which the cells are permeabilized reversibly by an electric impulse (EP-A 290 395, WO 87/06614). Further processes comprise the calcium-phosphate-mediated transformation, DEAE-dextran-mediated transformation, the incubation of dry embryos in DNA-containing solution or other methods of direct introduction of DNA (DE 4 005 152, WO 90/12096, US 4,684,611). Appropriate processes have been described (e.g. in Bilang et al. (1991) Gene 100:247-250; Scheid et al. (1991) Mol Gen Genet 228:104-112; Guerche et al. (1987) 10 Plant Science 52:111-116; Neuhaus et al. (1987) Theor Appl Genet 75:30-36; Klein et al. (1987) Nature 327:70-73; Howell et al. (1980) Science 208:1265; Horsch et al. (1985) Science 227:1229-1231; DeBlock et al. (1989) Plant Physiology 91:694-701; Methods for Plant Molecular Biology (Weissbach and Weissbach, eds.) Academic Press Inc. (1988); and Methods in Plant Molecular Biology (Schuler and Zielinski, eds.) Academic Press Inc. (1989)). Physiological methods of introducing DNA into plant cells have been reviewed by Oard (1991) Biotech Adv 9:1-11.

- 25 In the case of these "direct" transformation methods, no particular requirements are made on the plasmid used. It is possible to use simple plasmids such as those of the pUC series, pBR322, M13mp series, pACYC184 etc.

30

- Besides these "direct" transformation techniques, transformation may also be carried out by bacterial infection by means of Agrobacterium (e.g. EP 0 116 718), viral infection by means of viral vectors (EP 0 067 553; US 4,407,956; WO 95/34668; WO 93/03161) or 35 by means of pollen (EP 0 270 356; WO 85/01856; US 4,684,611).

- Transformation is preferably carried out by means of agrobacteria which contain disarmed Ti-plasmid vectors, using the latter's 40 natural ability to transfer genes to plants (EP-A 0 270 355; EP-A 0 116 718). Agrobacterium transformation is widespread for transforming dicotyledones, but is also increasingly applied to monocotyledones (Toriyama et al. (1988) Bio/Technology 6: 1072-1074; Zhang et al. (1988) Plant Cell Rep 7:379-384; Zhang et al. (1988) Theor Appl Genet 76:835-840; Shimamoto et al. (1989) Nature 45 338:274-276; Datta et al. (1990) Bio/Technology 8: 736-740; Christou et al. (1991) Bio/Technology 9:957-962; Peng et al. (1991) International Rice Research Institute, Manila, Philippines

## 60

563-574; Cao et al. (1992) Plant Cell Rep 11:585-591; Li et al. (1993) Plant Cell Rep 12:250-255; Rathore et al. (1993) Plant Mol Biol 21:871-884; Fromm et al. (1990) Bio/Technology 8:833-839; Gordon-Kamm et al. (1990) Plant Cell 2:603-618; D'Halluin et al. 5 (1992) Plant Cell 4:1495-1505; Walters et al. (1992) Plant Mol Biol 18:189-200; Koziel et al. (1993) Biotechnology 11:194-200; Vasil IK (1994) Plant Mol Biol 25:925-937; Weeks et al. (1993) Plant Physiol 102:1077-1084; Somers et al. (1992) Bio/Technology 10:1589-1594; WO 92/14828; Hiei et al. (1994) Plant J 6:271-282).

10

The strains most often used for agrobacterial transformation, *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*, contain a plasmid (Ti and Ri plasmids, respectively), which is transferred to the plant after agrobacterial infection. Part of this plasmid, 15 called T-DNA (transferred DNA), is integrated into the genome of the plant cell. Alternatively, *Agrobacterium* may also transfer binary vectors (mini Ti plasmids) to plants and integrate them into the genome of said plants.

20

The application of *Agrobacterium tumefaciens* to the transformation of plants, using tissue culture explants, has been described (inter alia, Horsch RB et al. (1985) Science 225:1229ff; Fraley et al. (1983) Proc Natl Acad Sci USA 80: 4803-4807; Bevans et al. 25 (1983) Nature 304:184-187). Many *Agrobacterium tumefaciens* strains are capable of transferring genetic material, such as, for example, the strains EHA101[pEHA101], EHA105[pEHA105], LBA4404[pAL4404], C58C1[pMP90] and C58C1[pGV2260] (Hood et al. (1993) Transgenic Res 2:208-218; Hoekema et al. (1983) Nature 303:179-181; Koncz and Schell (1986) Gen Genet 204:383-396; De- 30 blaere et al. (1985) Nucl Acids Res 13: 4777-4788).

When using agrobacteria, the expression cassette must be integrated into special plasmids, either a shuttle or intermediate 35 vector or a binary vector. When using a Ti or Ri plasmid for transformation, then at least the right border, but usually the right and left borders of the Ti or Ri plasmid T-DNA are connected as a flanking region to the expression cassette to be introduced. Preference is given to using binary vectors. Binary 40 vectors may replicate both in *E. coli* and in agrobacteria and contain the components required for transfer into a plant system. They normally contain a selection marker gene for selection of transformed plants (e.g. the nptII gene which imparts a resistance to kanamycin) and a linker or polylinker flanked by the 45 right and left T-DNA border sequences. They contain moreover, outside the T-DNA border sequence, also a selection marker which enables transformed *E. coli* and/or agrobacteria to be selected

## 61

(e.g. the nptIII gene which imparts a resistance to kanamycin). Corresponding vectors may be transformed directly into *Agrobacterium* (Holsters et al. (1978) *Mol Gen Genet* 163:181-187).

- 5 Binary vectors are based, for example, on "broad host range" plasmids such as pRK252 (Bevan et al. (1984) *Nucl Acid Res* 12:8711-8720) and pTJS75 (Watson et al. (1985) *EMBO J* 4(2):277-284). A large group of the binary vectors used is derived from pBIN19 (Bevan et al. (1984) *Nucl Acid Res* 12:8711-8720).
- 10 Hajdukiewicz et al. developed a binary vector (pPZP) which is smaller and more efficient than the previously customary vectors (Hajdukiewicz et al. (1994) *Plant Mol Biol* 25:989-994). Improved and particularly preferred binary vector systems for *Agrobacterium*-mediated transformation are described in WO 02/00900.
- 15

The *agrobacteria* transformed with a vector of this kind may then be used in the known manner for transforming plants, in particular crop plants such as, for example, oilseed rape, for example

20 by bathing wounded leaves or leaf sections in an *agrobacterial* solution and subsequently culturing them in suitable media. The transformation of plants by *agrobacteria* has been described (White FF, *Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization*, edited by S.D. Kung and R. Wu, Academic Press, 1993, pp. 15-38; Jenes B et al. (1993) *Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization*, edited by S.D. Kung and R. Wu, Academic Press, pp.128-143; Potrykus (1991) *Annu Rev Plant Physiol Plant Molec Biol* 42:205-225). Transgenic plants may

25 be regenerated in the known manner from the transformed cells of the wounded leaves or leaf sections.

30

Different explants, cell plants, tissues, organs, embryos, seeds, microspores or other unicellular or multicellular cellular structures derived from a plant organism may be used for transformation. Transformation processes adjusted to the particular explants, cultures or tissues are known to the skilled worker. Examples which may be mentioned are: shoot internodes (Fry J et al. (1987) *Plant Cell Rep.* 6:321-325), hypocotyls (Radke SE et al. (1988) *Theor Appl Genet* 75:685-694; Schröder M et al. (1994) *Physiologia Plant* 92: 37-46.; Stefanov I et al. (1994) *Plant Sci.* 95:175-186; Weier et al. (1997) *Fett/Lipid* 99:160-165), cotyledonous petioles (Meloney MM et al. (1989) *Plant Cell Rep* 8:238-242; Weier D et al. (1998) *Molecular Breeding* 4:39-46), microspores and proembryos (Pechnan (1989) *Plant Cell Rep.* 8:387-390) and flower stalks (Boulter ME et al. (1990) *Plant Sci* 70:91-99; Guerche P et al. (1987) *Mol Gen Genet* 206:382-386). In the case

35

40

45

## 62

of a direct gene transfer, mesophyll protoplasts (Chapel PJ & Glimelius K (1990) Plant Cell Rep 9: 105-108; Golz et al. (1990) Plant Mol Biol 15:475-483) or else hypocotyl protoplasts (Bergmann P & Glimelius K (1993) Physiologia Plant 88:604-611) and microspores (Chen JL et al. (1994) Theor Appl Genet 88:187-192; Jonesvilleneuve E et al. (1995) Plant Cell Tissue and Organ Cult 40:97-100) and shoot sections (Seki M et al. (1991) Plant Mol Biol 17:259-263) may be employed successfully.

- 10 Stably transformed cells, i.e. those which contain the introduced DNA integrated into the DNA of the host cell, may be selected from untransformed cells by using the selection process of the invention. The plants obtained may be grown and crossed in the usual way. Preferably, two or more generations should be cultured  
15 in order to ensure that the genomic integration is stable and can be inherited.

As soon as a transformed plant cell has been prepared, it is possible to obtain a complete plant by using processes known to the skilled worker. This involves, for example, starting from callus cultures, individual cells (e.g. protoplasts) or leaf disks (Vasil et al. (1984) Cell Culture and Somatic Cell Genetics of Plants, Vol I, II and III, Laboratory Procedures and Their Applications, Academic Press; Weissbach and Weissbach (1989) Methods for Plant Molecular Biology, Academic Press). It is possible to induce from these still undifferentiated callus cell masses the formation of shoot and root in the known manner. The seedlings obtained may be planted out and grown. Appropriate processes have been described (Fennell et al. (1992) Plant Cell Rep. 11: 567-570; Stoeger et al. (1995) Plant Cell Rep. 14:273-278; Jahne et al. (1994) Theor Appl Genet 89:525-533).

- The efficacy of expressing the transgenically expressed nucleic acids may be determined, for example, *in vitro* by shoot-meristem propagation using any of the selection methods described above. Moreover, changes in the type and level of expression of a target gene and the effect on the phenotype of the plant may be tested in greenhouse experiments using test plants.

40

The process of the invention is preferably used within the framework of plant biotechnology for generating plants having advantageous properties. The "nucleic acid sequence to be inserted" into the genome of the plant cell or the plant organism preferably  
45 comprises at least one expression cassette, said expression cassette being able to express, under the control of a promoter functional in plant cells or plant organisms, an RNA and/or a

## 63

protein which do not cause reduction of the expression, amount, activity and/or function of a marker protein but, particularly preferably, impart to the plant genetically altered in this way an advantageous phenotype. Numerous genes and proteins which may  
5 be used for achieving an advantageous phenotype, for example for the increase in quality of foodstuff or for producing particular chemicals or pharmaceuticals (Dunwell JM (2000) J Exp Bot 51 Spec No:487-96) are known to the skilled worker.

10 Thus it is possible to improve the suitability of the plants or the seeds thereof as foodstuff or feedstuff, for example by altering the compositions and/or the content of metabolites, in particular proteins, oils, vitamins and/or starch. It is also  
15 possible to increase the growth rate, yield or resistance to biotic or abiotic stress factors. Advantageous effects may be achieved both by transgenic expression of nucleic acids or proteins and by targeted reduction of the expression of endogenous genes, with respect to the phenotype of the transgenic plant. The  
20 advantageous effects which may be achieved in the transgenic plant comprise, for example:

- increased resistance to pathogens (biotic stress)
- 25 - increased resistance to environmental factors such as heat, cold, frost, drought, UV light, oxidative stress, wetness, salt, etc. (abiotic stress)
- 30 - increased yield
- improved quality, for example increased nutritional value, increased storability

35 The invention further relates to the use of the transgenic plants prepared according to the process of the invention and of the cells, cell cultures, plants or propagation material such as seeds or fruits derived from said plants, for preparing foodstuff  
40 or feedstuff, pharmaceuticals or fine chemicals such as, for example, enzymes, vitamins, amino acids, sugars, fatty acids, natural and synthetic flavorings, aroma substances and colorants. Particular preference is given to the production of triacyl glycerides, lipids, oils, fatty acids, starch, tocopherols and  
45 tocotrienols and also carotenoids. Genetically modified plants of the invention, which may be consumed by humans and animals may also be used as foodstuff or feedstuff, for example, directly or after preparation known per se.

## 64

As already mentioned above, the process of the invention comprises in a particularly advantageous embodiment, in a process step downstream of the selection, the deletion of the sequence coding for the marker protein (e.g. mediated by recombinase or as  
5 described in W003/004659) or the elimination by crossing and/or segregation of said sequences. (It is obvious to the skilled worker that, for this purpose, the nucleic acid sequence integrated into the genome and the sequence coding for the marker protein should have a separate chromosomal locus in the trans-  
10 formed cells. This, however, is the case in the majority of the resulting plants, merely for reasons of statistics). This procedure is particularly advantageous if the marker protein is a transgene which otherwise does not occur in the plant to be transformed. Although the resulting plant may still possibly con-  
15 tain the compound for reducing the expression, amount, activity and/or function of the marker protein, said compound would have no longer any "counterpart" in the form of said marker protein, and thus would have no effect. This is particularly the case if the marker protein is derived from a non-plant organism and/or is  
20 synthetic (for example the codA protein). It is, however, also possible to use plant marker proteins from other plant species, which otherwise do not occur in the cell to be transformed (i.e. if not introduced as transgene). Said marker proteins are referred to as "nonendogenous" marker proteins within the scope of  
25 the present invention.

Very particularly advantageously, the compound for reducing the expression, amount, activity and/or function of the marker protein is an RNA. After deletion or elimination by crossing/segregation,  
30 the resulting transgenic plant would have no longer any unnecessary (and, if appropriate, undesired) foreign protein. The sole foreign protein would be possibly the protein resulting from the nucleic acid sequence inserted into the genome. For reasons of product approval, this embodiment is particularly advantageous.  
35 As described above, said RNA may be an antisense RNA or, particularly preferably, a double-stranded RNA. It may be expressed separately from the RNA coding for the target protein but also, possibly, on the same strand as the latter.

40 In summary, the particularly advantageous embodiment comprises the following features:

A process for preparing transformed plant cells or organisms, which comprises the following steps:  
45

- a) transforming a population of plant cells which comprises at least one non-endogenous (preferably non-plant) marker pro-



## 65

- tein capable of converting directly or indirectly a substance X which is nontoxic for said population of plant cells into a substance Y which is toxic for said population, with at least one nucleic acid sequence to be inserted in combination with
- 5 at least one nucleic acid sequence coding for a ribonucleic acid sequence capable of reducing the expression, amount, activity and/or function of said marker protein, and
- 10 b) treating said population of plant cells with the substance X at a concentration which causes a toxic effect for nontransformed cells, due to the conversion by the marker protein, and
- 15 c) selecting transformed plant cells (and/or populations of plant cells, such as plant tissues or plants) whose genome contains said nucleic acid sequence and which have a growth advantage over nontransformed cells, due to the action of said compound, from said population of plant cells, the
- 20 selection being carried out under conditions under which the marker protein can exert its toxic effect on the nontransformed cells, and
- d) regenerating fertile plants, and
- 25 e) eliminating by crossing the nucleic acid sequence coding for the marker protein and isolating fertile plants whose genome contains said nucleic acid sequence but does not contain any
- 30 longer the sequence coding for the marker protein.

## Sequences

- 35 SEQ ID NO: 1 Nucleic acid sequence coding for E. coli cytosine deaminase (codA)
- SEQ ID NO: 2 amino acid sequence coding for E. coli cytosine deaminase (codA)
- 40 SEQ ID NO: 3 Nucleic acid sequence coding for E. coli cytosine deaminase (codA), with modified start codon (GTG/ATG) for expression in eukaryotes
- 45 SEQ ID NO: 4 Amino acid sequence coding for E. coli cytosine deaminase (codA), with modified start codon (GTG/ATG) for expression in eukaryotes

## 66

- SEQ ID NO: 5 Nucleic acid sequence coding for *Streptomyces griseolus* cytochrome P450-SU1 (suaC)
- 5 SEQ ID NO: 6 Amino acid sequence coding for *Streptomyces griseolus* cytochrome P450-SU1 (suaC)
- SEQ ID NO: 7 Nucleic acid sequence coding for *Agrobacterium tumefaciens* indoleacetamide hydrolase (tms2)
- 10 SEQ ID NO: 8 Amino acid sequence coding for *Agrobacterium tumefaciens* indoleacetamide hydrolase (tms2)
- 15 SEQ ID NO: 9 Nucleic acid sequence coding for *Agrobacterium tumefaciens* indoleacetamide hydrolase (tms2)
- SEQ ID NO: 10 Amino acid sequence coding for *Agrobacterium tumefaciens* indoleacetamide hydrolase (tms2)
- 20 SEQ ID NO: 11 Nucleic acid sequence coding for *Xanthobacter autotrophicus* haloalkane dehalogenase (dh1A)
- 25 SEQ ID NO: 12 Amino acid sequence coding for *Xanthobacter autotrophicus* haloalkane dehalogenase (dh1A)
- SEQ ID NO: 13 Nucleic acid sequence coding for Herpes simplex Virus 1 thymidine kinase
- 30 SEQ ID NO: 14 Amino acid sequence coding for Herpes simplex Virus 1 thymidine kinase
- 35 SEQ ID NO: 15 Nucleic acid sequence coding for Herpes simplex Virus 1 thymidine kinase
- SEQ ID NO: 16 Amino acid sequence coding for Herpes simplex Virus 1 thymidine kinase
- 40 SEQ ID NO: 17 Nucleic acid sequence coding for *Toxoplasma gondii* hypoxanthine-xanthine-guanine phosphoribosyl transferase
- 45 SEQ ID NO: 18 Amino acid sequence coding for *Toxoplasma gondii* hypoxanthine-xanthine-guanine phosphoribosyl transferase

## 67

- SEQ ID NO: 19 Nucleic acid sequence coding for E. coli xanthine-guanine phosphoribosyl transferase
- 5 SEQ ID NO: 20 Amino acid sequence coding for E. coli xanthine-guanine phosphoribosyl transferase
- SEQ ID NO: 21 Nucleic acid sequence coding for E. coli xanthine-guanine phosphoribosyl transferase
- 10 SEQ ID NO: 22 Amino acid sequence coding for E. coli xanthine-guanine phosphoribosyl transferase
- SEQ ID NO: 23 Nucleic acid sequence coding for E. coli purine nucleoside phosphorylase (deoD)
- 15 SEQ ID NO: 24 Nucleic acid sequence coding for E. coli purine nucleoside phosphorylase (deoD)
- 20 SEQ ID NO: 25 Nucleic acid sequence coding for Burkholderia caryophylli phosphonate monoester hydrolase (pehA)
- SEQ ID NO: 26 Amino acid sequence coding for Burkholderia caryophylli phosphonate monoester hydrolase (pehA)
- 25 SEQ ID NO: 27 Nucleic acid sequence coding for Agrobacterium rhizogenes tryptophan oxygenase (aux1)
- 30 SEQ ID NO: 28 Amino acid sequence coding for Agrobacterium rhizogenes tryptophan oxygenase (aux1)
- SEQ ID NO: 29 Nucleic acid sequence coding for Agrobacterium rhizogenes indoleacetamide hydrolase (aux2)
- 35 SEQ ID NO: 30 Amino acid sequence coding for Agrobacterium rhizogenes indoleacetamide hydrolase (aux2)
- SEQ ID NO: 31 Nucleic acid sequence coding for Agrobacterium tumefaciens tryptophan oxygenase (aux1)
- 40 SEQ ID NO: 32 Amino acid sequence coding for Agrobacterium tumefaciens tryptophan oxygenase (aux1)
- 45 SEQ ID NO: 33 Nucleic acid sequence coding for Agrobacterium tumefaciens indoleacetamide hydrolase (aux2)

## 68

- SEQ ID NO: 34 Amino acid sequence coding for *Agrobacterium tumefaciens* indoleacetamide hydrolase (aux2)
- 5 SEQ ID NO: 35 Nucleic acid sequence coding for *Agrobacterium vitis* indoleacetamide hydrolase (aux2)
- SEQ ID NO: 36 Amino acid sequence coding for *Agrobacterium vitis* indoleacetamide hydrolase (aux2)
- 10 SEQ ID NO: 37 Nucleic acid sequence coding for *Arabidopsis thaliana* 5-methylthioribose kinase (mtrK)
- SEQ ID NO: 38 Amino acid sequence coding for *Arabidopsis thaliana* 5-methylthioribose kinase (mtrK)
- 15 SEQ ID NO: 39 Nucleic acid sequence coding for *Klebsiella pneumoniae* 5-methylthioribose kinase (mtrK)
- 20 SEQ ID NO: 40 Amino acid sequence coding for *Klebsiella pneumoniae* 5-methylthioribose kinase (mtrK)
- SEQ ID NO: 41 Nucleic acid sequence coding for *Arabidopsis thaliana* alcohol dehydrogenase (adh)
- 25 SEQ ID NO: 42 Amino acid sequence coding for *Arabidopsis thaliana* alcohol dehydrogenase (adh)
- 30 SEQ ID NO: 43 Nucleic acid sequence coding for *Hordeum vulgare* (barley) alcohol dehydrogenase (adh)
- SEQ ID NO: 44 Amino acid sequence coding for *Hordeum vulgare* (barley) alcohol dehydrogenase (adh)
- 35 SEQ ID NO: 45 Nucleic acid sequence coding for *Oryza sativa* (rice) alcohol dehydrogenase (adh)
- 40 SEQ ID NO: 46 Amino acid sequence coding for *Oryza sativa* (rice) alcohol dehydrogenase (adh)
- SEQ ID NO: 47 Nucleic acid sequence coding for *Zea mays* (corn) alcohol dehydrogenase (adh)
- 45 SEQ ID NO: 48 Amino acid sequence coding for *Zea mays* (corn) alcohol dehydrogenase (adh)

## 69

- SEQ ID NO: 49 Nukleic acid sequence coding for a sense RNA fragment of E. coli cytosine deaminase (codARNAi-sense)
- 5 SEQ ID NO: 50 Oligonucleotide primer codA5'HindIII  
5'-AAGCTTGGCTAACAGTGTCGAATAACG-3'
- SEQ ID NO: 51 Oligonucleotide primer codA3'SalI  
5'-GTCGACGACAAAATCCCTTCCTGAGG-3'
- 10
- SEQ ID NO: 52 Nucleic acid sequence coding for an antisense RNA fragment of E. coli cytosine deaminase (codARNAi-anti)
- 15
- SEQ ID NO: 53 Oligonucleotide primer codA5'EcoRI  
5'-GAATTCGGCTAACAGTGTCGAATAACG-3'
- SEQ ID NO: 54 Oligonucleotide primer codA3'BamHI  
5'-GGATCCGACAAAATCCCTTCCTGAGG-3'
- 20
- SEQ ID NO: 55 Vector construct pBluKS-nitP-STLS1-35S-T
- SEQ ID NO: 56 Expression vector pSUN-1
- 25
- SEQ ID NO: 57 Transgenic expression vector pSUN-1-codA-RNAi
- SEQ ID NO: 58 Transgenic expression vector pSUN1-codA-RNAi-  
At.Act.-2-At.Als-R-ocsT
- 30
- SEQ ID NO: 59 Nukleic acid sequence coding for 5-methylthioribose kinase (mtrK) from corn (Zea mays); fragment
- 35
- SEQ ID NO: 60 Amino acid sequence coding for 5-methylthioribose kinase (mtrK) from corn (Zea mays); fragment
- SEQ ID NO: 61 Nucleic acid sequence coding for 5-methylthioribose kinase (mtrK) from oilseed rape (Brassica napus), fragment
- 40
- SEQ ID NO: 62 Amino acid sequence coding for 5-methylthioribose kinase (mtrK) from oilseed rape (Brassica napus), fragment
- 45
- SEQ ID NO: 63 Nucleic acid sequence coding for 5-methylthioribose kinase (mtrK) from oilseed rape (Brassica na-

70

pus), fragment

- 5 SEQ ID NO: 64 Amino acid sequence coding for 5-methylthioribose  
kinase (mtrK) from oilseed rape (*Brassica napus*),  
fragment
- 10 SEQ ID NO: 65 Nucleic acid sequence coding for 5-methylthiori-  
bose kinase (mtrK) from rice (*Oryza sativa*), frag  
ment
- 15 SEQ ID NO: 66 Amino acid sequence coding for 5-methylthioribose  
kinase (mtrK) from rice (*Oryza sativa*), fragment
- 20 SEQ ID NO: 67 Nucleic acid sequence coding for 5-methylthiori-  
bose kinase (mtrK) from soybean (*Glycine max*),  
fragment
- 25 SEQ ID NO: 68 Amino acid sequence coding for 5-methylthioribose  
kinase (mtrK) from soybean (*Glycine max*), fragment
- 30 SEQ ID NO: 69 Oligonucleotide primer codA5'C-term  
5'-CGTGAATACGGCGTGGAGTCG-3'
- 35 SEQ ID NO: 70 Oligonucleotide primer codA3'C-term  
5'-CGGCAGGATAATCAGGTTGG-3'
- 40 SEQ ID NO: 71 Oligonucleotide primer 35sT 5' primer  
5'-GTCAACGTAACCAACCCTGC-3'
- 45

## 71

## Figures

Fig.1: Inactivation of the marker protein gene by means of introducing a recombinase

5

P: promoter  
MP: Sequence coding for a marker protein  
R1/R2: Recombinase recognition sequences  
R: Recombinase or sequence coding for recombinase.

10

In a preferred embodiment, the marker protein gene is inactivated by introducing a sequence-specific recombinase. Preference is given to its expressing the recombinase, as depicted here, starting from an expression cassette.

15

The marker protein gene is flanked by recognition sequences for sequence-specific recombinases, with sequences of said marker protein gene being deleted by introducing said recombinase and thus said marker protein gene being inactivated.

20

Fig.2-A: Inactivation of the marker protein gene by the action of a sequence-specific nuclease

25

P: promoter  
DS: Recognition sequence for targeted induction of DNA double-strand breaks  
MP-DS-MP': Sequence coding for a marker protein, comprising a DS  
nDS: Inactivated DS  
E: Sequence-specific enzyme for targeted induction of DNA double-strand breaks

30

35

The marker protein gene may be established by a targeted mutation or deletion in the marker protein gene, for example by sequence-specific induction of DNA double-strand breaks at a recognition sequence for targeted induction of DNA double-strand breaks in or close to the marker protein gene (P-MP). The double-strand break may occur in the coding region or else the noncoding (such as, for example, the promoter) region, induces an illegitimate recombination (nonhomologous DNA-end joining) and thus, for example, a shift in the reading frame of said marker protein.

40

45

## 72

Fig.2-B: Inactivation of the marker protein gene by the action of a sequence-specific nuclease

5           P:           promoter  
           DS:           Recognition sequence for targeted induction  
                           of DNA double-strand breaks  
           MP:           Sequence coding for a marker protein  
           nDS:          Inactivated DS  
 10          E:           Sequence-specific enzyme for targeted  
                           induction of DNA double-strand breaks

15           The marker protein gene may be established by a targeted  
           deletion by sequence-specific induction of more than one  
           sequence-specific DNA double-strand break in or close to  
           said marker protein gene. The double-strand breaks may  
           occur in the coding region or else the noncoding (such  
           as, for example, the promoter) region and induce a dele-  
           tion in the marker protein gene. The marker protein gene  
 20           is preferably flanked by DS sequences and is completely  
           deleted by the action of enzyme E.

Fig. 3: Inactivation of the marker protein gene by inducing an  
 25           intramolecular homologous recombination, due to the ac-  
           tion of a sequence-specific nuclease

30           A/A':       Sequences with a sufficient length and homolo-  
                           gy to one another, in order to recombine with  
                           one another as a consequence of the induced  
                           double-strand break  
           P:           promoter  
           DS:           Recognition sequence for targeted induction  
                           of DNA double-strand breaks  
           MP:           Sequence coding for a marker protein  
 35          E:           Sequence-specific enzyme for targeted  
                           induction of DNA double-strand breaks

40           The marker protein gene may be inactivated by a deletion  
           by means of intramolecular homologous recombination. Said  
           homologous recombination may be initiated by sequence-  
           specific induction of DNA double-strand breaks at a rec-  
           ognition sequence for targeted induction of DNA double-  
           strand breaks in or close to the marker protein gene. The  
           homologous recombination occurs between the sequences A  
 45           and A' which have a sufficient length and homology to one  
           another in order to recombine with one another as a con-  
           sequence of the induced double-strand break. The recom-



## 73

ination causes a deletion of essential sequences of the marker protein gene.

Fig. 4: Inactivation of the marker protein gene by intermolecular homologous recombination

- 10 A/A': Sequences with a sufficient length and homology to one another in order to recombine with one another
- B/B': Sequences with a sufficient length and homology to one another in order to recombine with one another
- 15 P: promoter
- I: nucleic acid sequence/gene of interest to be inserted
- MP: Sequence coding for a marker protein

20 The marker protein gene (P-MP) may also be inactivated by a targeted insertion into the marker protein gene, for example by means of intermolecular homologous recombination. In this context, the region to be inserted is flanked on its 5' and 3' ends by nucleic acid sequences (A' and B', respectively), which have a sufficient length and homology to corresponding flanking sequences of the marker protein (A and B, respectively) in order to make possible a homologous recombination between A and A' and B and B'. The recombination causes a deletion of essential sequences of the marker protein gene.

30

Fig. 5: Inactivation of the marker protein gene by intermolecular homologous recombination due to the action of a sequence-specific nuclease

- 35 A/A': Sequences with a sufficient length and homology to one another in order to recombine with one another
- B/B': Sequences with a sufficient length and homology to one another in order to recombine with one another
- 40 P: promoter
- I: nucleic acid sequence/gene of interest to be inserted
- 45 MP: Sequence coding for a marker protein
- DS: Recognition sequence for targeted induction of DNA double-strand breaks
- E: Sequence-specific enzyme for targeted

## 74

## induction of DNA double-strand breaks

The marker protein gene may also be inactivated by a targeted insertion into the marker protein gene, for example by means of intermolecular homologous recombination. The homologous recombination may be initiated by sequence-specific induction of DNA double-strand breaks at a recognition sequence for targeted induction of DNA double-strand breaks in or close to the marker protein gene. In this context, the region to be inserted is flanked at its 5' and 3' ends by nucleic acid sequences (A' and B', respectively) which have a sufficient length and homology to corresponding flanking sequences of the marker protein gene (A and B, respectively) in order to make possible a homologous recombination between A and A' and B and B'. The recombination causes a deletion of essential sequences of the marker protein gene.

Fig. 6: Vector map for pBluKS-nitP-STLS1-35S-T (SEQ ID NO: 55)

NitP: promoter of the *A. thaliana* nitrilaseI gene (GenBank Acc. No.: Y07648.2, Hillebrand et al. (1996) Gene 170:197-200)

STLS-1 intron: intron of the potato ST-LS1 gene (Vancanneyt GF et al. (1990) Mol Gen Genet 220(2):245-250).

35S-Term: Terminator of the 35S CaMV gene (cauliflower mosaic virus; Franck et al. (1980) Cell 21:285-294).

Cleavage sites of relevant restriction endonucleases are indicated with their particular cleavage position.

Fig. 7: Vector map for the transgenic expression vector pSUN-1-codA-RNAi (SEQ ID NO: 57)

NitP: promoter of the *A. thaliana* nitrilaseI gene (GenBank Acc. No.: Y07648.2, Hillebrand et al. (1996) Gene 170:197-200)

STLS-1 intron: intron of the potato ST-LS1 gene (Vancanneyt GF et al. (1990) Mol Gen Genet 220(2):245-250).

35S-Term: Terminator of the 35S CaMV gene (cauliflower mosaic virus; Franck et al. (1980) Cell 21:285-294).

## 75

codA-sense: Nucleic acid sequence coding for a sense RNA fragment of E. coli cytosine deaminase (codARNAi-sense; SEQ ID NO: 49)

5 codA-anti: Nucleic acid sequence coding for an antisense RNA fragment of E. coli cytosine deaminase (codARNAi-anti; SEQ ID NO: 52)

10 LB/RB: Left and, respectively, right boundaries of Agrobacterium T-DNA

15 Cleavage sites of relevant restriction endonucleases are indicated with their particular cleavage position. Further elements represent customary elements of a binary Agrobacterium vector (aadA; ColE1; repA)

Fig. 8: Vector map for the transgenic expression vector pSUN1-codA-RNAi-At.Act.-2-At.Als-R-ocsT (SEQ ID NO: 58)

20 NitP: promoter of the A. thaliana nitrilaseI gene (GenBank Acc. No.: Y07648.2, Hillebrand et al. (1996) Gene 170:197-200)

25 STLS-1 intron: intron of the potato ST-LS1 gene (Vancanneyt GF et al. (1990) Mol Gen Genet 220(2):245-250).

30 35S-Term: Terminator of the 35S CaMV gene (cauliflower mosaic virus; Franck et al. (1980) Cell 21:285-294).

35 codA-sense: Nucleic acid sequence coding for a sense RNA fragment of E. coli cytosine deaminase (codARNAi-sense; SEQ ID NO: 49)

codA-anti: Nucleic acid sequence coding for an antisense RNA fragment of E. coli cytosine deaminase (codARNAi-anti; SEQ ID NO: 52)

40 Left border/right border: Left and, respectively, right boundaries of Agrobacterium T-DNA

45 Cleavage sites of relevant restriction endonucleases are indicated with their particular cleavage position. Further elements represent customary elements of a binary Agrobacterium vector (aadA; ColE1; repA)

## 76

Fig.9a-b: Sequence comparison of various 5-methylthioribose (MTR) kinases from various organisms, in particular plant organisms. Sequences from *Klebsiella pneumoniae*, *Clostridium tetani*, *Arabidopsis thaliana* (A.thaliana), oilseed rape (*Brassica napus*), soybean (Soy-1), rice (*Oryza sativa*-1) and also the consensus sequence (Consensus) are shown. Homologous regions can be readily deduced from the consensus sequence.

5  
10

15

20

25

30

35

40

45

## Exemplary embodiments

## General methods

- 5 The chemical synthesis of oligonucleotides may be carried out, for example, in the known manner by using the phosphoamide method (Voet, Voet, 2<sup>nd</sup> Edition, Wiley Press New York, pages 896-897). The cloning steps carried out within the scope of the present invention, such as, for example, restriction cleavages, agarose gel  
10 electrophoresis, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linking of DNA fragments, transformation of *E. coli* cells, cultivation of bacteria, propagation of phages and sequence analysis of recombinant DNA, are carried out as described in Sambrook et al. (1989)  
15 Cold Spring Harbor Laboratory Press; ISBN 0-87969-309-6. The sequencing of recombinant DNA molecules was carried out using a laser fluorescence DNA sequencer from ABI, according to the method of Sanger (Sanger et al. (1977) *Proc Natl Acad Sci* USA 74:5463-5467).

20

Example 1: Preparation of *codA* fragments

- First, a truncated nucleic acid variant of the *codA* gene, modified by the addition of recognition sequences of the restriction  
25 enzymes *Hind*III and *Sal*I, is prepared using the PCR technique. For this purpose, part of the *codA* gene (GeneBank Acc. No.: S56903; SEQ ID NO: 1) is amplified from the *E. coli* source organism by means of the polymerase chain reaction (PCR) using a  
30 sense-specific primer (*codA*5'*Hind*III; SEQ ID NO: 50) and an anti-sense-specific primer (*codA*3'*Sal*I; SEQ ID NO: 51).

*codA*5'*Hind*III: 5'-AAGCTTGGCTAACAGTGTCTGAATAACG-3' (SEQ ID NO: 50)

- 35 *codA*3'*Sal*I: 5'-GTCGACGACAAAATCCCTTCCTGAGG-3' (SEQ ID NO: 51)

The PCR was carried out in 50  $\mu$ l reaction mixture which contained:

- 40 - 2  $\mu$ l (200 ng) of *E. coli* genomic DNA  
- 0.2 mM dATP, dTTP, dGTP, dCTP  
- 1.5 mM Mg(OAc)<sub>2</sub>  
- 5  $\mu$ g of bovine serum albumin  
- 40 pmol of "*codA*5'*Hind*III" primer  
45 - 40 pmol of "*codA*3'*Sal*I" primer  
- 15  $\mu$ l of 3.3 $\times$  rTth DNA Polymerase XLPuffer (PE Applied Biosystems)

## 78

- 5U of rTth DNA Polymerase XL (PE Applied Biosystems)

The PCR is carried out under the following cycle conditions:

- 5      Step 1: 5 minutes 94°C (denaturation)  
         Step 2: 3 seconds 94°C  
         Step 3: 1 minute 60°C (annealing)  
         Step 4: 2 minutes 72°C (elongation)
- 10      30 repeats of steps 2 to 4
- Step 5: 10 minutes 72°C (post elongation)
- 15      Step 6: 4°C (waiting loop)

The amplicon (codARNAi-sense; SEQ ID NO: 49) is cloned using standard methods into the PCR cloning vector pGEM-T (Promega). The identity of the amplicon generated is confirmed by sequencing using the M13F (-40) primer.

Another truncated fragment of the *codA* gene, modified by the addition of recognition sequences of the restriction enzymes EcoRI and BamHI, is amplified using a sense-specific primer (codA5'EcoRI; SEQ ID NO: 53) and an antisense-specific primer (codA3'BamHI; SEQ ID NO: 54).

codA5'EcoRI: 5'-GAATTCGGCTAACAGTGTCTGAATAACG-3' (SEQ ID NO: 53)

30 codA3'BamHI: 5'-GGATCCGACAAAATCCCTTCCTGAGG-3' (SEQ ID NO: 54)

The PCR was carried out in 50 µl reaction mixture which contained:

- 35      - 2 µl (200 ng) of *E. coli* genomic DNA  
         - 0.2 mM dATP, dTTP, dGTP, dCTP  
         - 1.5 mM Mg(OAc)<sub>2</sub>  
         - 5 µg of bovine serum albumin
- 40      - 40 pmol of "codA5'EcoRI" primer  
         - 40 pmol of "codA3'BamHI" primer  
         - 15 µl of 3.3x rTth DNA Polymerase XLPuffer (PE Applied Biosystems)
- 45      - 5U of rTth DNA Polymerase XL (PE Applied Biosystems)

The PCR is carried out under the following cycle conditions:

## 79

- Step 1: 5 minutes 94°C (denaturation)  
Step 2: 3 seconds 94°C  
Step 3: 1 minute 60°C (annealing)  
5 Step 4: 2 minutes 72°C (elongation)  
  
30 repeats of steps 2 to 4  
  
Step 5: 10 minutes 72°C (post elongation)  
10 Step 6: 4°C (waiting loop)

The amplicon (codARNAi-anti; SEQ ID NO: 52) is cloned using standard methods into the PCR cloning vector pGEM-T (Promega). The  
15 identity of the amplicon generated is confirmed by sequencing using the M13F (-40) primer.

Example 2 Preparation of the transgenic expression vector for  
expressing a codA double-stranded RNA

20

The codA fragments generated in example 1 are used for preparing a DNA construct suitable for expressing a double-stranded codA RNA (pSUN-codA-RNAi). The construct is suitable for reducing the steady-state RNA level of the codA gene in transgenic plants and,  
25 as a result therefrom, suppressing codA gene expression by using the double-strand RNA interference (dsRNAi) technique. For this purpose, the codA RNAi cassette is first constructed in the plasmid pBluKS-nitP-STLS1-35S-T and then, in a further cloning step, completely transferred to the pSUN-1 plasmid.

30

The vector pBluKS-nitP-STLS1-35S-T (SEQ ID NO: 55) is a derivative of pBluescript KS (Stratagene) and contains the promoter of the *A. thaliana* nitrilaseI gene (GenBank Acc. No.: Y07648.2, nucleotides 2456 to 4340, Hillebrand et al. (1996) Gene  
35 170:197-200), the STLS-1 intron (Vancanneyt GF et al. (1990) Mol Gen Genet 220(2):245-250), restriction cleavage sites flanking the intron on its 5' and 3' sides and enabling DNA fragments to be inserted in a directed manner, and the terminator of the 35S  
40 CaMV gene (cauliflower mosaic virus; Franck et al. (1980) Cell 21:285-294). Using these restriction cleavage sites (HindIII, SalI, EcoRI, BamHI), the fragments codARNAi-sense (SEQ ID NO: 49) and codARNAi-anti (SEQ ID NO: 52) are inserted into said vector, thereby producing the finished codA RNAi cassette.

45

For this purpose, the codA sense fragment (codARNAi-sense SEQ ID NO: 49) is first excised from the pGEM-T vector, using the enzymes HindIII and SalI, isolated and ligated into the pBluKS-

## 80

nitP-STLS1-35S-T vector under standard conditions. This vector had previously been cleaved using the restriction enzymes HindIII and SalI. Correspondingly positive clones are identified by analytical restriction digest and sequencing.

5

The vector obtained (pBluKS-nitP-codAsense-STLS1-35S-T) is digested using the restriction enzymes BamHI and EcoRI. The codA-anti fragment (codARNAi-anti; SEQ ID NO: 52) is excised from the corresponding pGEM-T vector, using BamHI and EcoRI, isolated and  
10 ligated into the cut vector under standard conditions. Correspondingly positive clones which contain the complete codA-RNAi cassette (pBluKS-nitP-codAsense-STLS1-codAanti-35S-T) are identified by analytical restriction digest and sequencing.

15

The codA-RNAi cassette is transferred into the pSUN-1 vector (SEQ ID NO: 56) by using the SacI and KpnI restriction cleavage sites flanking the cassette. The resulting vector pSUN1-codA-RNAi (see Fig. 7; SEQ ID NO: 57) is used for transforming transgenic  
20 *A.thaliana* plants which express an active codA gene (see below). The plant expression vector pSUN-1 is particularly suitable within the scope of the process of the invention, since it does not contain any other positive selection marker.

25 The resulting vector, pSUN1-codA-RNAi, enables an artificial codA-dsRNA variant consisting of two identical nucleic acid elements which are separated by an intron and inverted to one another to be constitutively expressed. Transcription of this artificial codA-dsRNA variant results in the formation of a  
30 double-stranded RNA molecule, owing to the complementarity of the inverted nucleic acid elements. The presence of this molecule induces the suppression of codA gene expression (accumulation of RNA) by means of double-strand RNA interference.

35 Example 4: Preparation of transgenic *Arabidopsis thaliana* plants

Transgenic *Arabidopsis thaliana* plants which express transgenically the *E. coli* codA gene as a marker protein ("A.  
thaliana-[codA]"), were prepared as described (Kirik et al.

40 (2000) EMBO J 19(20):5562-6).

The *A. thaliana*-[codA] plants are transformed with an *Agrobacterium tumefaciens* strain (GV3101 [pMP90]) on the basis of a modified vacuum infiltration method (Clough S & Bent A (1998) Plant J  
45 16(6):735-43; Bechtold N et al. (1993) CR Acad Sci Paris 1144(2):204-212). The *Agrobacterium tumefaciens* cells used have previously been transformed with the DNA construct described



## 81

- (pSUN1-codA-RNAi). In this way, double transgenic *A. thaliana*-[codA] plants are generated which express an artificial codA double-stranded RNA under the control of the constitutive nitrilase1 promoter. Expression of the codA gene is suppressed as a consequence of the dsRNAi effect induced by the presence of this artificial codA-dsRNA. Said double transgenic plants may be identified owing to their regained ability to grow in the presence of 5-fluorocytosine in the culture medium.
- 10 Seeds of primary transformants are selected on the basis of the regained ability to grow in the presence of 5-fluorocytosine. For this purpose, the T1 seeds of the primary transformants are laid out on selection medium containing 200 µg/ml 5-fluorocytosine.
- 15 These selection plates are incubated under long-day conditions (16 h of light, 21°C/8 h of darkness, 18°C). Seedlings which develop normally in the presence of 5-fluorocytosine are separated after 7 days and transferred to new selection plates. These plates are incubated for another 14 under unchanged conditions.
- 20 The resistant seedlings are then transplanted into soil and cultured under short-day conditions (8 h of light, 21°C/16 h of darkness, 18°C). After 14 days, the young plants are transferred to the greenhouse and cultured under short-day conditions.
- 25 Example 5: Preparation of a plant transformation vector containing an expression cassette for expressing a double-stranded codA RNA and a plant selection marker
- 30 A plant selection marker consisting of a mutated variant of the *A. thaliana* Als gene, coding for the acetolactate synthase under the control of the promoter of the *A. thaliana* actin-2 gene (Meagher RB & Williamson RE (1994) The plant cytoskeleton. In The Plant Cytoskeleton (Meyerowitz, E. & Somerville, C., eds), pp. 1049-1084. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), and the octopine synthase terminator (GIELEN J et al.(1984) EMBO J 3:835-846) is inserted into pSUN1-codA-RNAi (see Fig. 7; SEQ ID NO: 57) (At.Act.-2-At.Als-R-ocST).
- 35 In The Plant Cytoskeleton (Meyerowitz, E. & Somerville, C., eds), pp. 1049-1084. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), and the octopine synthase terminator (GIELEN J et al.(1984) EMBO J 3:835-846) is inserted into pSUN1-codA-RNAi (see Fig. 7; SEQ ID NO: 57) (At.Act.-2-At.Als-R-ocST).
- 40 For this purpose, the pSUN1-codA-RNAi vector is first linearized using the restriction enzyme Pvu II. Subsequently, a linear DNA fragment with blunt ends, coding for a mutated variant of the acetolactate synthase (Als-R gene), is ligated into said linearized vector under standard conditions. Prior to ligation, this
- 45 DNA fragment has been digested with the restriction enzyme KpnI and the protruding ends have been converted into blunt ends by treatment with Pwo DNA polymerase (Roche) according to the

## 82

manufacturer's instructions. This mutated variant of the *A. thaliana* Als gene cannot be inhibited by herbicides of the imidazolinone type. By expressing this mutated A.tAls-R gene, the plants obtain the ability to grow in the presence of the herbicide Pursuit™. Correspondingly positive clones (pSUN1-codA-RNAi-At.Act.-2-At.Als-R-ocsT; SEQ ID NO: 57) are identified by analytical restriction digest and sequencing.

The vector obtained enables an artificial codA RNA variant (consisting of two identical nucleic acid elements which are separated by an intron and inverted to one another) and a mutated variant of the *A. thaliana* Als gene to be expressed constitutively. Transcription of this artificial codA RNA variant results in the formation of a double-stranded RNA molecule, owing to the complementarity of the inverted nucleic acid elements. The presence of this molecule induces the suppression of codA gene expression (accumulation of RNA) by means of double-strand RNA interference. Expression of the Als-R gene imparts to the plants the ability to grow in the presence of herbicides of the imidazolinone type.

Example 6: Preparation of transgenic *Arabidopsis thaliana* plants

Transgenic *Arabidopsis thaliana* plants expressing the *E. coli* codA gene as a marker protein ("A.thaliana-[codA]") were prepared as described (Kirik et al.(2000) EMBO J 19(20):5562-6).

The A.thaliana-[codA] plants are transformed with an *Agrobacterium tumefaciens* strain (GV3101 [pMP90]) on the basis of a modified vacuum infiltration method (Clough S & Bent A (1998) Plant J 16(6):735-43; Bechtold N et al. (1993) CR Acad Sci Paris 1144(2):204-212). The *Agrobacterium tumefaciens* cells used have previously been transformed with the DNA construct described (pSUN1-codA-RNAi-At.Act.-2-At.Als-R-ocsT; SEQ ID NO: 57). In this way, double transgenic *A. thaliana*-[codA] plants are generated which additionally express an artificial codA double-stranded RNA and a herbicide-insensitive variant of the Als gene (Als-R) under the control of the constitutive nitrilase promoter (A.thaliana-[codA]-[codA-RNAi-At.Act.-2-At.Als-R-ocsT]). Expression of the codA gene is suppressed as a consequence of the dsRNAi effect induced by the presence of this artificial codA-dsRNA. These double transgenic plants may be identified owing to their regained ability to grow in the presence of 5-fluorocytosine in the culture medium. In addition, positively transformed plants can be selected owing to their ability to grow in the presence of the her-

## 83

bicide Pursuit in the culture medium.

For the purpose of selection, the T1 seeds of primary transformants are therefore laid out on selection medium containing  
5 100 µg/ml 5-fluorocytosine. These selection plates are incubated under long-day conditions (16 h of light, 21°C/8 h of darkness, 18°C). Seedlings which develop normally in the presence of 5-fluorocytosine are separated after 28 days and transferred to new  
10 selection plates. These plates are incubated for another 14 days under unchanged conditions. The resistant seedlings are then transplanted into soil and cultured under short-day conditions (8 h of light, 21°C/16 h of darkness, 18°C). After a further 14 days, the young plants are transferred to the greenhouse and cul-  
15 tured under short-day conditions.

In addition, seeds of the primary transformants, owing to their ability to grow in the presence of the herbicide Pursuit™, may be selected. It is furthermore possible to carry out dual selection  
20 using the herbicide Pursuit™ and 5-fluorocytosine. For this purpose, the T1 seeds of primary transformants are laid out on selection medium containing the herbicide Pursuit™ at a concentration of 100 nM (in the case of dual selection, 100 µg/ml 5-fluorocytosine is likewise present). These selection plates are  
25 incubated under long-day conditions (16 h of light, 21°C/8 h of darkness, 18°C).

Seedlings which develop normally in the presence of Pursuit™ (Pursuit™ and 5-fluorocytosine) are separated after 28 days and  
30 transferred to new selection plates. These plates are incubated under unchanged conditions for another 14 days. The resistant seedlings are then transplanted into soil and cultured under short-day conditions (8 h of light, 21°C/16 h of darkness, 18°C).  
35 After 14 days, the young plants are transferred to the greenhouse and cultured under short-day conditions.

Example 7: Analysis of the double transgenic *A. thaliana* plants selected using 5-fluorocytosine and/or Pursuit  
40 (*A.thaliana*-[*codA*]-[*codA*-RNAi- *At.Act.-2-At.Als-R-ocST*])

Integration of the T-DNA region of the vector used for transformation, pSUN1-*codA*-RNAi-*A.tAls-R*, into the genomic DNA of the  
45 starting plant (*A.thaliana*-[*codA*]) and the loss of *codA*-specific mRNA in these transgenic plants (*A.thaliana*-[*codA*]-[*codA*-RNAi-*At.Act.-2-At.Als-R-ocST*]) can be detected by applying Southern

## 84

analyses and PCR techniques or Northern analyses.

In order to carry out said analyses, total RNA and DNA are isolated from leaf tissue of the transgenic plants and suitable controls (using the RNeasy Maxi Kit (RNA) and Dneasy Plant Maxi Kit (genomic DNA), respectively, according to the manufacturer's information by Qiagen).

In the PCR analyses, the genomic DNA may be used directly as a basis (template) for the PCR. Total RNA is transcribed to cDNA prior to the PCR. The cDNA synthesis is carried out using the reverse transcriptase Superscript II (Invitrogen) according to the manufacturer's information.

Example 8: Detection of the reduction in the steady-state amount of *codA* RNA in the positively selected double transgenic plants (*A.thaliana* [*codA*]-[*codA*-RNAi-*At.Act.-2-At.Als-R-ocsT*]) in comparison with the starting plants (*A.thaliana* [*codA*]) used for transformation, by means of cDNA synthesis with subsequent PCR amplification.

PCR amplification of the *codA*-specific cDNA:

The cDNA of the *codA* gene (ACCESSION S56903) may be amplified using a sense-specific primer (*codA*5'C-term SEQ ID NO: 69) and an antisense-specific primer (*codA*3'C-term SEQ ID NO: 70). The PCR conditions to be chosen are as follows:

The PCR was carried out in 50 µl reaction mixture which contained:

- 2 µl (200 ng) of cDNA from *A.thaliana* -[*codA*] or *A.thaliana* [*codA*]-[*codA*-RNAi-*At.Act.-2-At.Als-R-ocsT*] plants
- 0.2 mM dATP, dTTP, dGTP, dCTP
- 1.5 mM Mg(OAc)<sub>2</sub>
- 5 µg of bovine serum albumin
- 40 pmol of *codA*5'C-term SEQ ID NO: 69
- 40 pmol of *codA*3'C-term SEQ ID NO: 70
- 15 µl of 3.3x rTth DNA Polymerase XLPuffer (PE Applied Biosystems)
- 5U of rTth DNA Polymerase XL (PE Applied Biosystems)

The PCR was carried out under the following cycle conditions:

Step 1: 5 minutes 94°C (denaturation)

Step 2: 3 seconds 94°C

Step 3: 1 minute 56°C (annealing)

## 85

Step 4: 2 minutes 72°C (elongation)  
30 repeats of steps 2 to 4  
Step 5: 10 minutes 72°C (post elongation)  
Step 6: 4°C (waiting loop)

5 In the positively selected plants, the steady-state amount of the mRNA of the *codA* gene and the amount of CODA protein resulting therefrom is reduced so much that a quantitative conversion of 5-fluorocytosine to 5-fluorouracil can no longer occur. Consequently, these plants (in contrast to the untransformed plants) can grow in the presence of 5-fluorocytosine. Thus it is demonstrated that transgenic plants can be identified owing to the applied principle of preventing expression of a negative selection marker.

15 Example 9: Detection of the DNA coding for *codA*-RNAi by using genomic DNA of the positively selected double transgenic plants (*A.thaliana* [*codA*]-[*codA*-RNAi-*At.Act.-2-At.Als-R-ocST*])

20 The *codA*-RNAi transgene may be amplified using a *codA*-specific primer (e.g. *codA*5'HindIII SEQ ID NO: 50) and a 35S terminator-specific primer (35sT 5' Primer SEQ ID NO: 71). Using this primer combination, it is possible to detect specifically only the DNA coding for the *codA* RNAi construct, since the *codA* gene which was already present in the starting plants (*A.thaliana* [*codA*]) used for transformation is flanked by the nos terminator.

The PCR conditions to be chosen are as follows:

30 The PCR was carried out in a 50 µl reaction mixture which contains:

- 2 µl (200ng) of genomic DNA from the *A.thaliana* [*codA*]-[*codA*-RNAi-*At.Act.-2-At.Als-R-ocST*] plants
- 35 - 0.2 mM dATP, dTTP, dGTP, dCTP
- 1.5 mM Mg(OAc)<sub>2</sub>
- 5 µg of bovine serum albumin
- 40 pmol of *codA*-specific sense primer (SEQ ID NO: 50, 53 or 40 69)
- 40 pmol of 35sT 5' primer SEQ ID NO: 71
- 15 µl of 3.3x rTth DNA Polymerase XLPuffer (PE Applied Biosystems)
- 45 - 5U of rTth DNA Polymerase XL (PE Applied Biosystems)

The PCR was carried out under the following cycle conditions:  
Step 1: 5 minutes 94°C (denaturation)

## 86

- Step 2: 3 seconds 94°C  
Step 3: 1 minute 56°C (annealing)  
Step 4: 2 minutes 72°C (elongation)  
30 repeats of steps 2 to 4  
5 Step 5: 10 minutes 72°C (post elongation)  
Step 6: 4°C (waiting loop)

In this way, it is possible to detect in the positively selected plants integration of the *codA*-RNAi DNA construct into the chromosomal DNA of the starting plants used for transformation. Thus it is demonstrated that transgenic plants can be identified owing to the applied principle of preventing expression of a negative selection marker.

- 15 Example 10: Detection of the reduction in the steady-state amount of *codA* RNA in the positively selected double transgenic plants (*A.thaliana* [*codA*]-[*codA*-RNAi-  
At.Act.-2-At.Als-R-ocsT]) in comparison with the starting plants (*A.thaliana* [*codA*]) used for transformation, by Northern analysis.

## Gel-electrophoretic RNA fractionation:

- 25 For each RNA agarose gel, 3 g of agar are dissolved in 150 ml of H<sub>2</sub>O (f.c. 1.5% (w/v)) in a microwave oven and cooled to 60°C. The addition of 20 ml of 10x MEN (0.2 M MOPS, 50 mM sodium acetate, 10 mM EDTA) and 30 ml of formaldehyde (f.c. 2.2 M) causes further cooling so that the well-mixed solution must be poured speedily.  
30 Formaldehyde prevents the formation of secondary structures in the RNA, and therefore the rate of migration is approximately proportional to the molecular weight (LEHRBACH H et al. (1977) Biochem J 16: 4743-4751). The RNA samples are denatured, prior to application to the gel, in the following mixture: 20 µl of RNA  
35 (1-2 µg/µl), 5 µl of 10x MEN buffer, 6 µl of formaldehyde, 20 µl of formamide.

- The mixture is mixed and incubated at 65°C for 10 minutes. 1/10 volume of sample buffer and 1 µl of ethidium bromide (10 mg/ml) are added and the sample is then applied. Gel electrophoresis is carried out in horizontal gels in 1x MEN at 120 V for two to three hours. After electrophoresis, the gel is photographed under UV light with the aid of a ruler for subsequent determination of the fragment length. This is followed by blotting the RNA to a nylon membrane according to the information in: SAMBROOK J et al. Molecular cloning: A laboratory manual. Cold Spring Harbor, New York, Cold Spring Harbor Laboratory Press, 1989.

## Radioactive labeling of DNA fragments and Northern hybridization

5 The *codA* cDNA fragment (*codARNAi*-sense SEQ ID No: 49) can be labeled using, for example, the High Prime kit sold by Roche Diagnostics. The High Prime kit is based on the "random primed" method for DNA labeling originally described by Feinberg and  
10 Vogelstein. Labeling is carried out by denaturing approx. 25 ng of DNA in 9-11  $\mu$ l of  $H_2O$  at 95°C for 10 min. After a short incubation on ice, 4  $\mu$ l of High Prime solution (contains a random primer mixture, 4 units of Klenow polymerase and 0.125 mM dATP, dTTP and dGTP each in a reaction buffer containing 50% glycerol) and  
15 3-5  $\mu$ l of [ $\alpha^{32}P$ ]dCTP (30-50  $\mu$ Ci) are added. The reaction mixture is incubated at 37°C for at least 10 min and the unincorporated dCTP is then separated from the now radiolabeled DNA by means of gel filtration via a Sephadex G-50 column. The fragment is subsequently denatured at 95°C for 10 min and kept on ice until used. The following hybridization and preincubation buffers are used:

20

Hypo Hybond  
250 mM sodium phosphate buffer pH 7.2  
1 mM EDTA  
7% SDS (g/v)  
25 250 mM NaCl  
10  $\mu$ g/ml ssDNA  
5% polyethylene glycol (PEG) 6000  
40% formamide

30 The hybridization temperature when using Hypo Hybond is 42°C and the duration of hybridization is 16-24 h. The RNA filters are washed using three different solutions: 2 x SSC (300 mM NaCl; 30 mM sodium citrate) + 0.1% SDS, 1 x SSC + 0.1% SDS and 0.1 x SSC + 0.1% SDS. The duration and intensity of washing depend on  
35 the strength of the activity bond. After washing, the filters are sealed in plastic foil and an X-ray film (X-OMat, Kodak) is exposed overnight at -70°C. The signal strength on the X-ray films is a measure of the amount of *codA* mRNA molecules in the total RNA bound on the membranes. Thus it is possible to detect the re-  
40 duction in *codA* mRNA in the positively selected plants compared to the starting plants used for transformation.

In the positively selected plants, the steady-state amount of the mRNA of the *codA* gene and the amount of CODA protein produced resulting therefrom is reduced so much that a quantitative conversion of 5-fluorocytosine to 5-fluorouracil can no longer occur.  
45 Consequently, these plants (in contrast to the untransformed plants) can grow in the presence of 5-fluorocytosine. Thus it is

## 88

demonstrated that transgenic plants can be identified owing to the applied principle of preventing expression of a negative selection marker.

5 Example 11: Summary of the results of "negative-negative"  
selection

Transformation of the codA-transgenic Arabidopsis plants with the  
codA-dsRNA construct (pSUN1-codA-RNAi-At.Act.-2-At.Als-R-ocsT;  
10 SEQ ID NO: 57) results in a significantly increased number of  
double transgenic plants into whose genome the RNAi construct has  
been successfully integrated, in the case of both single selec-  
tion (with 5-fluorocytosine alone) and dual selection (Pursuit™  
and 5-fluorocytosine) (in each case in comparison with untrans-  
15 formed plants). The analysis by means of PCR (see above) confirms  
the double transgenic state for the majority of the plants gener-  
ated in this way. This successfully demonstrates the practicabil-  
ity of the present invention, i.e. the usability of repression of  
a negative marker for positive selection (more or less a "nega-  
20 tive-negative" selection).

25

30

35

40

45



PF 53790

1

## SEQUENCE LISTING

&lt;110&gt; BASF Plant Science GmbH

&lt;120&gt; Novel selection processes

&lt;130&gt; PF53790-AT

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 71

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 1284

&lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1281)

&lt;223&gt; coding for cytosine deaminase (codA)

&lt;400&gt; 1

gtg tcg aat aac gct tta caa aca att att aac gcc cgg tta cca ggc	48
Val Ser Asn Asn Ala Leu Gln Thr Ile Ile Asn Ala Arg Leu Pro Gly	
1 5 10 15	
gaa gag ggg ctg tgg cag att cat ctg cag gac gga aaa atc agc gcc	96
Glu Glu Gly Leu Trp Gln Ile His Leu Gln Asp Gly Lys Ile Ser Ala	
20 25 30	
att gat gcg caa tcc ggc gtg atg ccc ata act gaa aac agc ctg gat	144
Ile Asp Ala Gln Ser Gly Val Met Pro Ile Thr Glu Asn Ser Leu Asp	
35 40 45	
gcc gaa caa ggt tta gtt ata ccg ccg ttt gtg gag cca cat att cac	192
Ala Glu Gln Gly Leu Val Ile Pro Pro Phe Val Glu Pro His Ile His	
50 55 60	
ctg gac acc acg caa acc gcc gga caa ccg aac tgg aat cag tcc ggc	240
Leu Asp Thr Thr Gln Thr Ala Gly Gln Pro Asn Trp Asn Gln Ser Gly	
65 70 75 80	
acg ctg ttt gaa ggc att gaa cgc tgg gcc gag cgc aaa gcg tta tta	288
Thr Leu Phe Glu Gly Ile Glu Arg Trp Ala Glu Arg Lys Ala Leu Leu	
85 90 95	
acc cat gac gat gtg aaa caa cgc gca tgg caa acg ctg aaa tgg cag	336
Thr His Asp Asp Val Lys Gln Arg Ala Trp Gln Thr Leu Lys Trp Gln	
100 105 110	
att gcc aac ggc att cag cat gtg cgt acc cat gtc gat gtt tcg gat	384
Ile Ala Asn Gly Ile Gln His Val Arg Thr His Val Asp Val Ser Asp	
115 120 125	
gca acg cta act gcg ctg aaa gca atg ctg gaa gtg aag cag gaa gtc	432
Ala Thr Leu Thr Ala Leu Lys Ala Met Leu Glu Val Lys Gln Glu Val	
130 135 140	
gcg ccg tgg att gat ctg caa atc gtc gcc ttc cct cag gaa ggg att	480
Ala Pro Trp Ile Asp Leu Gln Ile Val Ala Phe Pro Gln Glu Gly Ile	
145 150 155 160	

PF 53790

2

ttg tgc tat ccc aac ggt gaa gcg ttg ctg gaa gag gcg tta cgc tta	528
Leu Ser Tyr Pro Asn Gly Glu Ala Leu Leu Glu Glu Ala Leu Arg Leu	
165 170 175	
ggg gca gat gta gtg ggg gcg att ccg cat ttt gaa ttt acc cgt gaa	576
Gly Ala Asp Val Val Gly Ala Ile Pro His Phe Glu Phe Thr Arg Glu	
180 185 190	
tac ggc gtg gag tgc ctg cat aaa acc ttc gcc ctg gcg caa aaa tac	624
Tyr Gly Val Glu Ser Leu His Lys Thr Phe Ala Leu Ala Gln Lys Tyr	
195 200 205	
gac cgt ctc atc gac gtt cac tgt gat gag atc gat gac gag cag tgc	672
Asp Arg Leu Ile Asp Val His Cys Asp Glu Ile Asp Asp Glu Gln Ser	
210 215 220	
cgc ttt gtc gaa acc gtt gct gcc ctg gcg cac cat gaa ggc atg ggc	720
Arg Phe Val Glu Thr Val Ala Ala Leu Ala His His Glu Gly Met Gly	
225 230 235 240	
gcg cga gtc acc gcc agc cac acc acg gca atg cac tcc tat aac ggg	768
Ala Arg Val Thr Ala Ser His Thr Thr Ala Met His Ser Tyr Asn Gly	
245 250 255	
gcg tat acc tca cgc ctg ttc cgc ttg ctg aaa atg tcc ggt att aac	816
Ala Tyr Thr Ser Arg Leu Phe Arg Leu Leu Lys Met Ser Gly Ile Asn	
260 265 270	
ttt gtc gcc aac ccg ctg gtc aat att cat ctg caa gga cgt ttc gat	864
Phe Val Ala Asn Pro Leu Val Asn Ile His Leu Gln Gly Arg Phe Asp	
275 280 285	
acg tat cca aaa cgt cgc ggc atc acg cgc gtt aaa gag atg ctg gag	912
Thr Tyr Pro Lys Arg Arg Gly Ile Thr Arg Val Lys Glu Met Leu Glu	
290 295 300	
tcc ggc att aac gtc tgc ttt ggt cac gat gat gtc ttc gat ccg tgg	960
Ser Gly Ile Asn Val Cys Phe Gly His Asp Asp Val Phe Asp Pro Trp	
305 310 315 320	
tat ccg ctg gga acg gcg aat atg ctg caa gtg ctg cat atg ggg ctg	1008
Tyr Pro Leu Gly Thr Ala Asn Met Leu Gln Val Leu His Met Gly Leu	
325 330 335	
cat gtt tgc cag ttg atg ggc tac ggg cag att aac gat ggc ctg aat	1056
His Val Cys Gln Leu Met Gly Tyr Gly Gln Ile Asn Asp Gly Leu Asn	
340 345 350	
tta atc acc cac cac agc gca agg acg ttg aat ttg cag gat tac ggc	1104
Leu Ile Thr His His Ser Ala Arg Thr Leu Asn Leu Gln Asp Tyr Gly	
355 360 365	
att gcc gcc gga aac agc gcc aac ctg att atc ctg ccg gct gaa aat	1152
Ile Ala Ala Gly Asn Ser Ala Asn Leu Ile Ile Leu Pro Ala Glu Asn	
370 375 380	
ggg ttt gat gcg ctg cgc cgt cag gtt ccg gta cgt tat tgc gta cgt	1200
Gly Phe Asp Ala Leu Arg Arg Gln Val Pro Val Arg Tyr Ser Val Arg	
385 390 395 400	
ggc ggc aag gtg att gcc agc aca caa ccg gca caa acc acc gta tat	1248
Gly Gly Lys Val Ile Ala Ser Thr Gln Pro Ala Gln Thr Thr Val Tyr	
405 410 415	
ctg gag cag cca gaa gcc atc gat tac aaa cgt tga	1284
Leu Glu Gln Pro Glu Ala Ile Asp Tyr Lys Arg	
420 425	

PF 53790

3

&lt;210&gt; 2

&lt;211&gt; 427

&lt;212&gt; PRT

&lt;213&gt; Escherichia coli

&lt;400&gt; 2

Val	Ser	Asn	Asn	Ala	Leu	Gln	Thr	Ile	Ile	Asn	Ala	Arg	Leu	Pro	Gly	1	5	10	15
Glu	Glu	Gly	Leu	Trp	Gln	Ile	His	Leu	Gln	Asp	Gly	Lys	Ile	Ser	Ala	20	25	30	
Ile	Asp	Ala	Gln	Ser	Gly	Val	Met	Pro	Ile	Thr	Glu	Asn	Ser	Leu	Asp	35	40	45	
Ala	Glu	Gln	Gly	Leu	Val	Ile	Pro	Pro	Phe	Val	Glu	Pro	His	Ile	His	50	55	60	
Leu	Asp	Thr	Thr	Gln	Thr	Ala	Gly	Gln	Pro	Asn	Trp	Asn	Gln	Ser	Gly	65	70	75	80
Thr	Leu	Phe	Glu	Gly	Ile	Glu	Arg	Trp	Ala	Glu	Arg	Lys	Ala	Leu	Leu	85	90	95	
Thr	His	Asp	Asp	Val	Lys	Gln	Arg	Ala	Trp	Gln	Thr	Leu	Lys	Trp	Gln	100	105	110	
Ile	Ala	Asn	Gly	Ile	Gln	His	Val	Arg	Thr	His	Val	Asp	Val	Ser	Asp	115	120	125	
Ala	Thr	Leu	Thr	Ala	Leu	Lys	Ala	Met	Leu	Glu	Val	Lys	Gln	Glu	Val	130	135	140	
Ala	Pro	Trp	Ile	Asp	Leu	Gln	Ile	Val	Ala	Phe	Pro	Gln	Glu	Gly	Ile	145	150	155	160
Leu	Ser	Tyr	Pro	Asn	Gly	Glu	Ala	Leu	Leu	Glu	Glu	Ala	Leu	Arg	Leu	165	170	175	
Gly	Ala	Asp	Val	Val	Gly	Ala	Ile	Pro	His	Phe	Glu	Phe	Thr	Arg	Glu	180	185	190	
Tyr	Gly	Val	Glu	Ser	Leu	His	Lys	Thr	Phe	Ala	Leu	Ala	Gln	Lys	Tyr	195	200	205	
Asp	Arg	Leu	Ile	Asp	Val	His	Cys	Asp	Glu	Ile	Asp	Asp	Glu	Gln	Ser	210	215	220	
Arg	Phe	Val	Glu	Thr	Val	Ala	Ala	Leu	Ala	His	His	Glu	Gly	Met	Gly	225	230	235	240
Ala	Arg	Val	Thr	Ala	Ser	His	Thr	Thr	Ala	Met	His	Ser	Tyr	Asn	Gly	245	250	255	
Ala	Tyr	Thr	Ser	Arg	Leu	Phe	Arg	Leu	Leu	Lys	Met	Ser	Gly	Ile	Asn	260	265	270	
Phe	Val	Ala	Asn	Pro	Leu	Val	Asn	Ile	His	Leu	Gln	Gly	Arg	Phe	Asp	275	280	285	
Thr	Tyr	Pro	Lys	Arg	Arg	Gly	Ile	Thr	Arg	Val	Lys	Glu	Met	Leu	Glu	290	295	300	
Ser	Gly	Ile	Asn	Val	Cys	Phe	Gly	His	Asp	Asp	Val	Phe	Asp	Pro	Trp	305	310	315	320
Tyr	Pro	Leu	Gly	Thr	Ala	Asn	Met	Leu	Gln	Val	Leu	His	Met	Gly	Leu	325	330	335	

PF 53790

4

His Val Cys Gln Leu Met Gly Tyr Gly Gln Ile Asn Asp Gly Leu Asn  
 340 345 350  
 Leu Ile Thr His His Ser Ala Arg Thr Leu Asn Leu Gln Asp Tyr Gly  
 355 360 365  
 Ile Ala Ala Gly Asn Ser Ala Asn Leu Ile Ile Leu Pro Ala Glu Asn  
 370 375 380  
 Gly Phe Asp Ala Leu Arg Arg Gln Val Pro Val Arg Tyr Ser Val Arg  
 385 390 395 400  
 Gly Gly Lys Val Ile Ala Ser Thr Gln Pro Ala Gln Thr Thr Val Tyr  
 405 410 415  
 Leu Glu Gln Pro Glu Ala Ile Asp Tyr Lys Arg  
 420 425

&lt;210&gt; 3

&lt;211&gt; 1284

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; Description of the artificial sequence: coding for cytosine deaminase (codA)

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(3)

&lt;223&gt; mutation of GTG to ATG start codon for expression in eukaryotic hosts

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1281)

&lt;223&gt; coding for cytosine deaminase (codA)

&lt;400&gt; 3

atg tcg aat aac gct tta caa aca att att aac gcc cgg tta cca ggc 48  
 Met Ser Asn Asn Ala Leu Gln Thr Ile Ile Asn Ala Arg Leu Pro Gly  
 1 5 10 15  
 gaa gag ggg ctg tgg cag att cat ctg cag gac gga aaa atc agc gcc 96  
 Glu Glu Gly Leu Trp Gln Ile His Leu Gln Asp Gly Lys Ile Ser Ala  
 20 25 30  
 att gat gcg caa tcc ggc gtg atg ccc ata act gaa aac agc ctg gat 144  
 Ile Asp Ala Gln Ser Gly Val Met Pro Ile Thr Glu Asn Ser Leu Asp  
 35 40 45  
 gcc gaa caa ggt tta gtt ata ccg ccg ttt gtg gag cca cat att cac 192  
 Ala Glu Gln Gly Leu Val Ile Pro Pro Phe Val Glu Pro His Ile His  
 50 55 60  
 ctg gac acc acg caa acc gcc gga caa ccg aac tgg aat cag tcc ggc 240  
 Leu Asp Thr Thr Gln Thr Ala Gly Gln Pro Asn Trp Asn Gln Ser Gly  
 65 70 75 80  
 acg ctg ttt gaa ggc att gaa cgc tgg gcc gag cgc aaa gcg tta tta 288  
 Thr Leu Phe Glu Ile Glu Arg Trp Ala Glu Arg Lys Ala Leu Leu  
 85 90 95

PF 53790

5

acc cat gac gat gtg aaa caa cgc gca tgg caa acg ctg aaa tgg cag	336
Thr His Asp Asp Val Lys Gln Arg Ala Trp Gln Thr Leu Lys Trp Gln	
100 105 110	
att gcc aac ggc att cag cat gtg cgt acc cat gtc gat gtt tcg gat	384
Ile Ala Asn Gly Ile Gln His Val Arg Thr His Val Asp Val Ser Asp	
115 120 125	
gca acg cta act gcg ctg aaa gca atg ctg gaa gtg aag cag gaa gtc	432
Ala Thr Leu Thr Ala Leu Lys Ala Met Leu Glu Val Lys Gln Glu Val	
130 135 140	
gcg ccg tgg att gat ctg caa atc gtc gcc ttc cct cag gaa ggg att	480
Ala Pro Trp Ile Asp Leu Gln Ile Val Ala Phe Pro Gln Glu Gly Ile	
145 150 155 160	
ttg tcg tat ccc aac ggt gaa gcg ttg ctg gaa gag gcg tta cgc tta	528
Leu Ser Tyr Pro Asn Gly Glu Ala Leu Leu Glu Glu Ala Leu Arg Leu	
165 170 175	
ggg gca gat gta gtg ggg gcg att ccg cat ttt gaa ttt acc cgt gaa	576
Gly Ala Asp Val Val Gly Ala Ile Pro His Phe Glu Phe Thr Arg Glu	
180 185 190	
tac ggc gtg gag tcg ctg cat aaa acc ttc gcc ctg gcg caa aaa tac	624
Tyr Gly Val Glu Ser Leu His Lys Thr Phe Ala Leu Ala Gln Lys Tyr	
195 200 205	
gac cgt ctc atc gac gtt cac tgt gat gag atc gat gac gag cag tcg	672
Asp Arg Leu Ile Asp Val His Cys Asp Glu Ile Asp Asp Glu Gln Ser	
210 215 220	
cgc ttt gtc gaa acc gtt gct gcc ctg gcg cac cat gaa ggc atg ggc	720
Arg Phe Val Glu Thr Val Ala Ala Leu Ala His His Glu Gly Met Gly	
225 230 235 240	
gcg cga gtc acc gcc agc cac acc acg gca atg cac tcc tat aac ggg	768
Ala Arg Val Thr Ala Ser His Thr Thr Ala Met His Ser Tyr Asn Gly	
245 250 255	
gcg tat acc tca cgc ctg ttc cgc ttg ctg aaa atg tcc ggt att aac	816
Ala Tyr Thr Ser Arg Leu Phe Arg Leu Leu Lys Met Ser Gly Ile Asn	
260 265 270	
ttt gtc gcc aac ccg ctg gtc aat att cat ctg caa gga cgt ttc gat	864
Phe Val Ala Asn Pro Leu Val Asn Ile His Leu Gln Gly Arg Phe Asp	
275 280 285	
acg tat cca aaa cgt cgc ggc atc acg cgc gtt aaa gag atg ctg gag	912
Thr Tyr Pro Lys Arg Arg Gly Ile Thr Arg Val Lys Glu Met Leu Glu	
290 295 300	
tcc ggc att aac gtc tgc ttt ggt cac gat gat gtc ttc gat ccg tgg	960
Ser Gly Ile Asn Val Cys Phe Gly His Asp Asp Val Phe Asp Pro Trp	
305 310 315 320	
tat ccg ctg gga acg gcg aat atg ctg caa gtg ctg cat atg ggg ctg	1008
Tyr Pro Leu Gly Thr Ala Asn Met Leu Gln Val Leu His Met Gly Leu	
325 330 335	
cat gtt tgc cag ttg atg ggc tac ggg cag att aac gat ggc ctg aat	1056
His Val Cys Gln Leu Met Gly Tyr Gly Gln Ile Asn Asp Gly Leu Asn	
340 345 350	
tta atc acc cac cac agc gca agg acg ttg aat ttg cag gat tac ggc	1104
Leu Ile Thr His His Ser Ala Arg Thr Leu Asn Leu Gln Asp Tyr Gly	
355 360 365	

PF 53790

6

```

att gcc gcc gga aac agc gcc aac ctg att atc ctg ccg gct gaa aat 1152
ile Ala Ala Gly Asn Ser Ala Asn Leu ile ile Leu Pro Ala Glu Asn
    370                      375                      380

ggg ttt gat gcg ctg cgc cgt cag gtt ccg gta cgt tat tcg gta cgt 1200
Gly Phe Asp Ala Leu Arg Arg Gln Val Pro Val Arg Tyr Ser Val Arg
385                      390                      395                      400

ggc ggc aag gtg att gcc agc aca caa ccg gca caa acc acc gta tat 1248
Gly Gly Lys Val ile Ala Ser Thr Gln Pro Ala Gln Thr Thr Val Tyr
    405                      410                      415

ctg gag cag cca gaa gcc atc gat tac aaa cgt tga 1284
Leu Glu Gln Pro Glu Ala ile Asp Tyr Lys Arg
    420                      425

```

&lt;210&gt; 4

&lt;211&gt; 427

&lt;212&gt; PRT

&lt;213&gt; Artificial sequence

<223> Description of the artificial sequence: coding for  
cytosine deaminase (codA)

&lt;400&gt; 4

```

Met Ser Asn Asn Ala Leu Gln Thr ile ile Asn Ala Arg Leu Pro Gly
 1          5          10          15

Glu Glu Gly Leu Trp Gln ile His Leu Gln Asp Gly Lys ile Ser Ala
    20          25          30

ile Asp Ala Gln Ser Gly Val Met Pro ile Thr Glu Asn Ser Leu Asp
    35          40          45

Ala Glu Gln Gly Leu Val ile Pro Pro Phe Val Glu Pro His ile His
    50          55          60

Leu Asp Thr Thr Gln Thr Ala Gly Gln Pro Asn Trp Asn Gln Ser Gly
    65          70          75          80

Thr Leu Phe Glu Gly ile Glu Arg Trp Ala Glu Arg Lys Ala Leu Leu.
    85          90          95

Thr His Asp Asp Val Lys Gln Arg Ala Trp Gln Thr Leu Lys Trp Gln
    100         105         110

ile Ala Asn Gly ile Gln His Val Arg Thr His Val Asp Val Ser Asp
    115         120         125

Ala Thr Leu Thr Ala Leu Lys Ala Met Leu Glu Val Lys Gln Glu Val
    130         135         140

Ala Pro Trp ile Asp Leu Gln ile Val Ala Phe Pro Gln Glu Gly ile
    145         150         155         160

Leu Ser Tyr Pro Asn Gly Glu Ala Leu Leu Glu Glu Ala Leu Arg Leu
    165         170         175

Gly Ala Asp Val Val Gly Ala ile Pro His Phe Glu Phe Thr Arg Glu
    180         185         190

Tyr Gly Val Glu Ser Leu His Lys Thr Phe Ala Leu Ala Gln Lys Tyr
    195         200         205

Asp Arg Leu ile Asp Val His Cys Asp Glu ile Asp Asp Glu Gln Ser
    210         215         220

```

PF 53790

7

Arg Phe Val Glu Thr Val Ala Ala Leu Ala His His Glu Gly Met Gly  
 225 230 235 240  
 Ala Arg Val Thr Ala Ser His Thr Thr Ala Met His Ser Tyr Asn Gly  
 245 250 255  
 Ala Tyr Thr Ser Arg Leu Phe Arg Leu Leu Lys Met Ser Gly Ile Asn  
 260 265 270  
 Phe Val Ala Asn Pro Leu Val Asn Ile His Leu Gln Gly Arg Phe Asp  
 275 280 285  
 Thr Tyr Pro Lys Arg Arg Gly Ile Thr Arg Val Lys Glu Met Leu Glu  
 290 295 300  
 Ser Gly Ile Asn Val Cys Phe Gly His Asp Asp Val Phe Asp Pro Trp  
 305 310 315 320  
 Tyr Pro Leu Gly Thr Ala Asn Met Leu Gln Val Leu His Met Gly Leu  
 325 330 335  
 His Val Cys Gln Leu Met Gly Tyr Gly Gln Ile Asn Asp Gly Leu Asn  
 340 345 350  
 Leu Ile Thr His His Ser Ala Arg Thr Leu Asn Leu Gln Asp Tyr Gly  
 355 360 365  
 Ile Ala Ala Gly Asn Ser Ala Asn Leu Ile Ile Leu Pro Ala Glu Asn  
 370 375 380  
 Gly Phe Asp Ala Leu Arg Arg Gln Val Pro Val Arg Tyr Ser Val Arg  
 385 390 395 400  
 Gly Gly Lys Val Ile Ala Ser Thr Gln Pro Ala Gln Thr Thr Val Tyr  
 405 410 415  
 Leu Glu Gln Pro Glu Ala Ile Asp Tyr Lys Arg  
 420 425

<210> 5  
 <211> 1221  
 <212> DNA  
 <213> Streptomyces griseolus  
 <220>  
 <221> CDS  
 <222> (1)..(1218)  
 <223> coding for cytochrome P450-Su1 (suaC)

<400> 5  
 atg acc gat acc gcc acg acg ccc cag acc acg gac gca ccc gcc ttc 48  
 Met Thr Asp Thr Ala Thr Thr Pro Gln Thr Thr Asp Ala Pro Ala Phe  
 1 5 10 15  
 ccg agc aac cgg agc tgt ccc tac cag tta ccg gac ggc tac gcc cag 96  
 Pro Ser Asn Arg Ser Cys Pro Tyr Gln Leu Pro Asp Gly Tyr Ala Gln  
 20 25 30  
 ctc cgg gac acc ccc ggc ccc ctg cac cgg gtg acg ctc tac gac ggc 144  
 Leu Arg Asp Thr Pro Gly Pro Leu His Arg Val Thr Leu Tyr Asp Gly  
 35 40 45  
 cgt cag gcg tgg gtg gtg acc aag cac gag gcc gcg cgc aaa ctg ctc 192  
 Arg Gln Ala Trp Val Val Thr Lys His Glu Ala Ala Arg Lys Leu Leu  
 50 55 60

PF 53790

8

ggc gac ccc cgg ctg tcc tcc aac cgg acg gac gac aac ttc ccc gcc	240
Gly Asp Pro Arg Leu Ser Ser Asn Arg Thr Asp Asp Asn Phe Pro Ala	
65 70 75 80	
acg tca ccg cgc ttc gag gcc gtc cgg gag agc ccg cag gcg ttc atc	288
Thr Ser Pro Arg Phe Glu Ala Val Arg Glu Ser Pro Gln Ala Phe Ile	
85 90 95	
ggc ctg gac ccg ccc gag cac ggc acc cgg cgg cgg atg acg atc agc	336
Gly Leu Asp Pro Pro Glu His Gly Thr Arg Arg Arg Met Thr Ile Ser	
100 105 110	
gag ttc acc gtc aag cgg atc aag ggc atg cgc ccc gag gtc gag gag	384
Glu Phe Thr Val Lys Arg Ile Lys Gly Met Arg Pro Glu Val Glu Glu	
115 120 125	
gtg gtg cac ggc ttc ctc gac gag atg ctg gcc gcc ggc ccg acc gcc	432
Val Val His Gly Phe Leu Asp Glu Met Leu Ala Ala Gly Pro Thr Ala	
130 135 140	
gac ctg gtc agt cag ttc gcg ctg ccg gtg ccc tcc atg gtg atc tgc	480
Asp Leu Val Ser Gln Phe Ala Leu Pro Val Pro Ser Met Val Ile Cys	
145 150 155 160	
cga ctc ctc ggc gtg ccc tac gcc gac cac gag ttc ttc cag gac gcg	528
Arg Leu Leu Gly Val Pro Tyr Ala Asp His Glu Phe Phe Gln Asp Ala	
165 170 175	
agc aag cgg ctg gtg cag tcc acg gac gcg cag agc gcg ctc acc gcg	576
Ser Lys Arg Leu Val Gln Ser Thr Asp Ala Gln Ser Ala Leu Thr Ala	
180 185 190	
cgg aac gac ctc gcg ggt tac ctg gac ggc ctc atc acc cag ttc cag	624
Arg Asn Asp Leu Ala Gly Tyr Leu Asp Gly Leu Ile Thr Gln Phe Gln	
195 200 205	
acc gaa ccg ggc gcg ggc ctg gtg ggc gct ctg gtc gcc gac cag ctg	672
Thr Glu Pro Gly Ala Gly Leu Val Gly Ala Leu Val Ala Asp Gln Leu	
210 215 220	
gcc aac ggc gag atc gac cgt gag gaa ctg atc tcc acc gcg atg ctg	720
Ala Asn Gly Glu Ile Asp Arg Glu Glu Leu Ile Ser Thr Ala Met Leu	
225 230 235 240	
ctc ctc atc gcc ggc cac gag acc acg gcc tcg atg acc tcc ctc agc	768
Leu Leu Ile Ala Gly His Glu Thr Thr Ala Ser Met Thr Ser Leu Ser	
245 250 255	
gtg atc acc ctg ctg gac cac ccc gag cag tac gcc gcc ctg cgc gcc	816
Val Ile Thr Leu Leu Asp His Pro Glu Gln Tyr Ala Ala Leu Arg Ala	
260 265 270	
gac cgc agc ctc gtg ccc ggc gcg gtg gag gaa ctg ctc cgc tac ctc	864
Asp Arg Ser Leu Val Pro Gly Ala Val Glu Glu Leu Leu Arg Tyr Leu	
275 280 285	
gcc atc gcc gac atc gcg ggc ggc cgc gtc gcc acg gcg gac atc gag	912
Ala Ile Ala Asp Ile Ala Gly Gly Arg Val Ala Thr Ala Asp Ile Glu	
290 295 300	
gtc gag ggg cac ctc atc cgg gcc ggc gag ggc gtg atc gtc gtc aac	960
Val Glu Gly His Leu Ile Arg Ala Gly Glu Gly Val Ile Val Val Asn	
305 310 315 320	
tcg ata gcc aac cgg gac ggc acg gtg tac gag gac ccg gac gcc ctc	1008
Ser Ile Ala Asn Arg Asp Gly Thr Val Tyr Glu Asp Pro Asp Ala Leu	
325 330 335	



PF 53790

9

```

gac atc cac cgc tcc gcg cgc cac cac ctc gcc ttc ggc ttc ggc gtg 1056
Asp Ile His Arg Ser Ala Arg His His Leu Ala Phe Gly Phe Gly Val
      340                      345                      350

cac cag tgc ctg ggc cag aac ctc gcc cgg ctg gag ctg gag gtc atc 1104
His Gln Cys Leu Gly Gln Asn Leu Ala Arg Leu Glu Leu Glu Val Ile
      355                      360                      365

ctc aac gcc ctc atg gac cgc gtc ccg acg ctg cga ctg gcc gtc ccc 1152
Leu Asn Ala Leu Met Asp Arg Val Pro Thr Leu Arg Leu Ala Val Pro
      370                      375                      380

gtc gag cag ttg gtg ctg cgg ccg ggt acg acg atc cag ggc gtc aac 1200
Val Glu Gln Leu Val Leu Arg Pro Gly Thr Thr Ile Gln Gly Val Asn
      385                      390                      395                      400

gaa ctc ccg gtc acc tgg tga 1221
Glu Leu Pro Val Thr Trp
      405

```

&lt;210&gt; 6

&lt;211&gt; 406

&lt;212&gt; PRT

&lt;213&gt; Streptomyces griseolus

&lt;400&gt; 6

```

Met Thr Asp Thr Ala Thr Thr Pro Gln Thr Thr Asp Ala Pro Ala Phe
  1              5              10              15

Pro Ser Asn Arg Ser Cys Pro Tyr Gln Leu Pro Asp Gly Tyr Ala Gln
      20              25              30

Leu Arg Asp Thr Pro Gly Pro Leu His Arg Val Thr Leu Tyr Asp Gly
      35              40              45

Arg Gln Ala Trp Val Val Thr Lys His Glu Ala Ala Arg Lys Leu Leu
      50              55              60

Gly Asp Pro Arg Leu Ser Asn Arg Thr Asp Asp Asn Phe Pro Ala
      65              70              75              80

Thr Ser Pro Arg Phe Glu Ala Val Arg Glu Ser Pro Gln Ala Phe Ile
      85              90              95

Gly Leu Asp Pro Pro Glu His Gly Thr Arg Arg Arg Met Thr Ile Ser
      100              105              110

Glu Phe Thr Val Lys Arg Ile Lys Gly Met Arg Pro Glu Val Glu Glu
      115              120              125

Val Val His Gly Phe Leu Asp Glu Met Leu Ala Ala Gly Pro Thr Ala
      130              135              140

Asp Leu Val Ser Gln Phe Ala Leu Pro Val Pro Ser Met Val Ile Cys
      145              150              155              160

Arg Leu Leu Gly Val Pro Tyr Ala Asp His Glu Phe Phe Gln Asp Ala
      165              170              175

Ser Lys Arg Leu Val Gln Ser Thr Asp Ala Gln Ser Ala Leu Thr Ala
      180              185              190

Arg Asn Asp Leu Ala Gly Tyr Leu Asp Gly Leu Ile Thr Gln Phe Gln
      195              200              205

Thr Glu Pro Gly Ala Gly Leu Val Gly Ala Leu Val Ala Asp Gln Leu
      210              215              220

```

PF 53790

10

Ala Asn Gly Glu Ile Asp Arg Glu Glu Leu Ile Ser Thr Ala Met Leu  
 225 230 235 240  
 Leu Leu Ile Ala Gly His Glu Thr Thr Ala Ser Met Thr Ser Leu Ser  
 245 250 255  
 Val Ile Thr Leu Leu Asp His Pro Glu Gln Tyr Ala Ala Leu Arg Ala  
 260 265 270  
 Asp Arg Ser Leu Val Pro Gly Ala Val Glu Glu Leu Leu Arg Tyr Leu  
 275 280 285  
 Ala Ile Ala Asp Ile Ala Gly Gly Arg Val Ala Thr Ala Asp Ile Glu  
 290 295 300  
 Val Glu Gly His Leu Ile Arg Ala Gly Glu Gly Val Ile Val Val Asn  
 305 310 315 320  
 Ser Ile Ala Asn Arg Asp Gly Thr Val Tyr Glu Asp Pro Asp Ala Leu  
 325 330 335  
 Asp Ile His Arg Ser Ala Arg His His Leu Ala Phe Gly Phe Gly Val  
 340 345 350  
 His Gln Cys Leu Gly Gln Asn Leu Ala Arg Leu Glu Leu Glu Val Ile  
 355 360 365  
 Leu Asn Ala Leu Met Asp Arg Val Pro Thr Leu Arg Leu Ala Val Pro  
 370 375 380  
 Val Glu Gln Leu Val Leu Arg Pro Gly Thr Thr Ile Gln Gly Val Asn  
 385 390 395 400  
 Glu Leu Pro Val Thr Trp  
 405

&lt;210&gt; 7

&lt;211&gt; 1404

&lt;212&gt; DNA

&lt;213&gt; Agrobacterium tumefaciens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1401)

&lt;223&gt; coding for indoleacetamide hydrolase (tms2)

&lt;400&gt; 7

atg gtg ccc att acc tcg tta gca caa acc cta gaa cgc ctg aga cgg 48  
 Met Val Pro Ile Thr Ser Leu Ala Gln Thr Leu Glu Arg Leu Arg Arg  
 1 5 10 15  
 aaa gac tac tcc tgc tta gaa cta gta gaa act ctg ata gcg cgt tgc 96  
 Lys Asp Tyr Ser Cys Leu Glu Leu Val Glu Thr Leu Ile Ala Arg Cys  
 20 25 30  
 caa gct gca aaa cca tta aat gcc ctt ctg gct aca gac tgg gat ggc 144  
 Gln Ala Ala Lys Pro Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Gly  
 35 40 45  
 ttg cgg cga agc gcc aaa aaa att gat cgt cat gga aac gcc gga tta 192  
 Leu Arg Arg Ser Ala Lys Lys Ile Asp Arg His Gly Asn Ala Gly Leu  
 50 55 60  
 ggt ctt tgc ggc att cca ctc tgt ttt aag gcg aac atc gcg acc ggc 240  
 Gly Leu Cys Gly Ile Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly  
 65 70 75 80

PF 53790

11

ata ttt cct aca agc gct gct act ccg gcg ctg ata aac cac ttg cca	288
Ile Phe Pro Thr Ser Ala Ala Thr Pro Ala Leu Ile Asn His Leu Pro	
85 90 95	
aag ata cca tcc cgc gtc gca gaa aga ctt ttt tca gct gga gca ctg	336
Lys Ile Pro Ser Arg Val Ala Glu Arg Leu Phe Ser Ala Gly Ala Leu	
100 105 110	
ccg ggt gcc tcg gga aac atg cat gag tta tcg ttt gga att acg agc	384
Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser	
115 120 125	
aac aac tat gcc acc ggt gcg gtg cgg aac ccg tgg aat cca agt ctg	432
Asn Asn Tyr Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu	
130 135 140	
ata cca gga ggc tca agc ggt ggt gtg gct gct gcg gtg gca agc cga	480
Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Val Ala Ser Arg	
145 150 155 160	
ttg atg tta ggc ggc ata ggc acc gat acc ggt gca tct gtt cgc cta	528
Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu	
165 170 175	
ccc gca gcc ctg tgt ggc gta gta gga ttt cga ccg acg ctt gct cga	576
Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Leu Ala Arg	
180 185 190	
tat cca aga gat cgg ata ata ccg gtc agc ccc acc cgg gac acc gcc	624
Tyr Pro Arg Asp Arg Ile Ile Pro Val Ser Pro Thr Arg Asp Thr Ala	
195 200 205	
gga atc ata gcg cag tgc gta gcc gat gtt ata atc ctc gac cag gtg	672
Gly Ile Ile Ala Gln Cys Val Ala Asp Val Ile Ile Leu Asp Gln Val	
210 215 220	
att tcc gga cgg tcg gcg aaa att tca ccc atg ccg ctg aag ggg ctt	720
Ile Ser Gly Arg Ser Ala Lys Ile Ser Pro Met Pro Leu Lys Gly Leu	
225 230 235 240	
cgg atc ggc ctc ccc act acc tac ttt tac gat gac ctt gat gct gat	768
Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp	
245 250 255	
gtg gcc ttc gca gct gaa acg acg att cgc ttg cta gcc aac aga ggc	816
Val Ala Phe Ala Ala Glu Thr Thr Ile Arg Leu Leu Ala Asn Arg Gly	
260 265 270	
gta acc ttt gtt gaa gcc gac atc ccc cac cta gag gaa ctg aat agt	864
Val Thr Phe Val Glu Ala Asp Ile Pro His Leu Glu Glu Leu Asn Ser	
275 280 285	
ggg gca agt ttg cca att gcg ctt tac gaa ttt cca cac gct cta aaa	912
Gly Ala Ser Leu Pro Ile Ala Leu Tyr Glu Phe Pro His Ala Leu Lys	
290 295 300	
aag tat ctc gac gat ttt gtg gga aca gtt tct ttt tct gac gtt atc	960
Lys Tyr Leu Asp Asp Phe Val Gly Thr Val Ser Phe Ser Asp Val Ile	
305 310 315 320	
aaa gga att cgt agc ccc gat gta gcg aac att gtc agt gcg caa att	1008
Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Val Ser Ala Gln Ile	
325 330 335	
gat ggg cat caa att tcc aac gat gaa tat gaa ctg gcg cgt caa tcc	1056
Asp Gly His Gln Ile Ser Asn Asp Glu Tyr Glu Leu Ala Arg Gln Ser	
340 345 350	

PF 53790

12

```

ttc agg cca agg ctc cag gcc act tat cgg aat tac ttc aga ctc tat 1104
Phe Arg Pro Arg Leu Gln Ala Thr Tyr Arg Asn Tyr Phe Arg Leu Tyr
355 360 365

cag tta gat gca atc ctt ttc cca act gca ccc tta gcg gcc aaa gcc 1152
Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Ala Ala Lys Ala
370 375 380

ata ggt cag gag tcg tca gtc atc cac aat ggc tca atg atg aac act 1200
Ile Gly Gln Glu Ser Ser Val Ile His Asn Gly Ser Met Met Asn Thr
385 390 395 400

ttc aag atc tac gtg cga aat gtg gac cca agc agc aac gca ggc cta 1248
Phe Lys Ile Tyr Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu
405 410 415

cct ggg ttg agc ctt cct gcc tgc ctt aca cct gat cgc ttg cct gtt 1296
Pro Gly Leu Ser Leu Pro Ala Cys Leu Thr Pro Asp Arg Leu Pro Val
420 425 430

gga atg gaa att gat gga tta gcg ggg tca gac cac cgt ctg tta gca 1344
Gly Met Glu Ile Asp Gly Leu Ala Gly Ser Asp His Arg Leu Leu Ala
435 440 445

atc ggg gca gca tta gaa aaa gcc ata aat ttt cct tcc ttt ccc gat 1392
Ile Gly Ala Ala Leu Glu Lys Ala Ile Asn Phe Pro Ser Phe Pro Asp
450 455 460

gct ttt aat tag 1404
Ala Phe Asn
465

<210> 8
<211> 467
<212> PRT
<213> Agrobacterium tumefaciens

<400> 8
Met Val Pro Ile Thr Ser Leu Ala Gln Thr Leu Glu Arg Leu Arg Arg
1 5 10 15
Lys Asp Tyr Ser Cys Leu Glu Leu Val Glu Thr Leu Ile Ala Arg Cys
20 25 30
Gln Ala Ala Lys Pro Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Gly
35 40 45
Leu Arg Arg Ser Ala Lys Lys Ile Asp Arg His Gly Asn Ala Gly Leu
50 55 60
Gly Leu Cys Gly Ile Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly
65 70 75 80
Ile Phe Pro Thr Ser Ala Ala Thr Pro Ala Leu Ile Asn His Leu Pro
85 90 95
Lys Ile Pro Ser Arg Val Ala Glu Arg Leu Phe Ser Ala Gly Ala Leu
100 105 110
Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser
115 120 125
Asn Asn Tyr Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu
130 135 140
Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Val Ala Ser Arg
145 150 155 160

```

PF 53790

13

Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu  
 165 170 175  
 Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Leu Ala Arg  
 180 185 190  
 Tyr Pro Arg Asp Arg Ile Ile Pro Val Ser Pro Thr Arg Asp Thr Ala  
 195 200 205  
 Gly Ile Ile Ala Gln Cys Val Ala Asp Val Ile Ile Leu Asp Gln Val  
 210 215 220  
 Ile Ser Gly Arg Ser Ala Lys Ile Ser Pro Met Pro Leu Lys Gly Leu  
 225 230 235 240  
 Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp  
 245 250 255  
 Val Ala Phe Ala Ala Glu Thr Thr Ile Arg Leu Leu Ala Asn Arg Gly  
 260 265 270  
 Val Thr Phe Val Glu Ala Asp Ile Pro His Leu Glu Glu Leu Asn Ser  
 275 280 285  
 Gly Ala Ser Leu Pro Ile Ala Leu Tyr Glu Phe Pro His Ala Leu Lys  
 290 295 300  
 Lys Tyr Leu Asp Asp Phe Val Gly Thr Val Ser Phe Ser Asp Val Ile  
 305 310 315 320  
 Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Val Ser Ala Gln Ile  
 325 330 335  
 Asp Gly His Gln Ile Ser Asn Asp Glu Tyr Glu Leu Ala Arg Gln Ser  
 340 345 350  
 Phe Arg Pro Arg Leu Gln Ala Thr Tyr Arg Asn Tyr Phe Arg Leu Tyr  
 355 360 365  
 Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Ala Ala Lys Ala  
 370 375 380  
 Ile Gly Gln Glu Ser Ser Val Ile His Asn Gly Ser Met Met Asn Thr  
 385 390 395 400  
 Phe Lys Ile Tyr Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu  
 405 410 415  
 Pro Gly Leu Ser Leu Pro Ala Cys Leu Thr Pro Asp Arg Leu Pro Val  
 420 425 430  
 Gly Met Glu Ile Asp Gly Leu Ala Gly Ser Asp His Arg Leu Leu Ala  
 435 440 445  
 Ile Gly Ala Ala Leu Glu Lys Ala Ile Asn Phe Pro Ser Phe Pro Asp  
 450 455 460  
 Ala Phe Asn  
 465

&lt;210&gt; 9

&lt;211&gt; 1404

&lt;212&gt; DNA

&lt;213&gt; Agrobacterium tumefaciens

&lt;220&gt;

&lt;221&gt; CDS

PF 53790

14

&lt;222&gt; (1)..(1401)

&lt;223&gt; coding for indoleacetamide hydrolase (tms2)

&lt;400&gt; 9

```

atg gtg ccc att acc tcg tta gca caa acc cta gaa cgc ctg aga cgg      48
Met Val Pro Ile Thr Ser Leu Ala Gln Thr Leu Glu Arg Leu Arg Arg
      1              5              10              15

aaa gac tac tcc tgc tta gaa cta gta gaa act ctg ata gcg cgt tgc      96
Lys Asp Tyr Ser Cys Leu Glu Leu Val Glu Thr Leu Ile Ala Arg Cys
      20              25              30

caa gct gca aaa cca tta aat gcc ctt ctg gct aca gac tgg gat ggc      144
Gln Ala Ala Lys Pro Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Gly
      35              40              45

ttg cgg cga agc gcc aaa aaa att gat cgt cat gga aac gcc gga tta      192
Leu Arg Arg Ser Ala Lys Lys Ile Asp Arg His Gly Asn Ala Gly Leu
      50              55              60

ggg ctt tgc ggc att cca ctc tgt ttt aag gcg aac atc gcg acc ggc      240
Gly Leu Cys Gly Ile Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly
      65              70              75              80

ata ttt cct aca agc gct gct act ccg gcg ctg ata aac cac ttg cca      288
Ile Phe Pro Thr Ser Ala Ala Thr Pro Ala Leu Ile Asn His Leu Pro
      85              90              95

aag ata cca tcc cgc gtc gca gaa aga ctt ttt tca gct gga gca ctg      336
Lys Ile Pro Ser Arg Val Ala Glu Arg Leu Phe Ser Ala Gly Ala Leu
      100             105             110

ccg ggt gcc tcg gga aac atg cat gag tta tcg ttt gga att acg agc      384
Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser
      115             120             125

aac aac tat gcc acc ggt gcg gtg cgg aac ccg tgg aat cca agt ctg      432
Asn Asn Tyr Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu
      130             135             140

ata cca gga ggc tca agc ggt ggt gtg gct gct gcg gtg gca agc cga      480
Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Ala Val Ala Ser Arg
      145             150             155             160

ttg atg tta ggc ggc ata ggc acc gat acc ggt gca tct gtt cgc cta      528
Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu
      165             170             175

ccc gca gcc ctg tgt ggc gta gta gga ttt cga ccg acg ctt gct cga      576
Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Leu Ala Arg
      180             185             190

tat cca aga gat cgg ata ata ccg gtc agc ccc acc cgg gac acc gcc      624
Tyr Pro Pro Arg Asp Arg Ile Ile Pro Val Ser Pro Thr Arg Asp Thr Ala
      195             200             205

gga atc ata gcg cag tgc gta gcc gat gtt ata atc ctc gat cag gtg      672
Gly Ile Ile Ala Gln Cys Val Ala Asp Val Ile Ile Leu Asp Gln Val
      210             215             220

att tcc gga cgg tcg gcg aaa att tca ccc atg ccg ctg aag ggg ctt      720
Ile Ser Gly Arg Ser Ala Lys Ile Ser Pro Met Pro Leu Lys Gly Leu
      225             230             235             240

cgg atc ggc ctc ccc act acc tac ttt tac gat gac ctt gat gct gat      768
Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp
      245             250             255

```

PF 53790

15

```

gtg gcc ttc gca gct gaa acg acg att cgc ttg cta gcc aac aga ggc 816
Val Ala Phe Ala Ala Glu Thr Thr Ile Arg Leu Leu Ala Asn Arg Gly
                260                265                270

gta acc ttt gtt gaa gcc gac atc ccc cac cta gag gaa ctg aat agt 864
Val Thr Phe Val Glu Ala Asp Ile Pro His Leu Glu Glu Leu Asn Ser
                275                280                285

ggg gca agt ttg cca att gcg ctt tac gaa ttt cca cac gct cta aaa 912
Gly Ala Ser Leu Pro Ile Ala Leu Tyr Glu Phe Pro His Ala Leu Lys
                290                295                300

aag tat ctc gac gat ttt gtg gga aca gtt tct ttt tct gac gtt atc 960
Lys Tyr Leu Asp Asp Phe Val Gly Thr Val Ser Phe Ser Asp Val Ile
305                310                315                320

aaa gga att cgt agc ccc gat gta gcg aac att gtc agt gcg caa att 1008
Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Val Ser Ala Gln Ile
                325                330                335

gat ggg cat caa att tcc aac gat gaa tat gaa ctg gcg cgt caa tcc 1056
Asp Gly His Gln Ile Ser Asn Asp Glu Tyr Glu Leu Ala Arg Gln Ser
                340                345                350

ttc agg cca agg ctc cag gcc act tat cgg aat tac ttc aga ctc tat 1104
Phe Arg Pro Arg Leu Gln Ala Thr Tyr Arg Asn Tyr Phe Arg Leu Tyr
                355                360                365

cag tta gat gca atc ctt ttc cca act gca ccc tta gcg gcc aaa gcc 1152
Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Ala Ala Lys Ala
370                375                380

ata ggt cag gag tcg tca gtc atc cac aat ggc tca atg ata aac act 1200
Ile Gly Gln Glu Ser Ser Val Ile His Asn Gly Ser Met Ile Asn Thr
385                390                395                400

ttc aag atc tac gtg cga aat gtg gac cca agc agc aac gca ggc cta 1248
Phe Lys Ile Tyr Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu
                405                410                415

cct ggg ttg agc ctt cct gcc tgc ctt aca cct gat cgc ttg cct gtt 1296
Pro Gly Leu Ser Leu Pro Ala Cys Leu Thr Pro Asp Arg Leu Pro Val
                420                425                430

gga atg gaa att gac gga tta gcg ggg tca gac cac cgt ctg tta gca 1344
Gly Met Glu Ile Asp Gly Leu Ala Gly Ser Asp His Arg Leu Leu Ala
                435                440                445

atc ggg gca gca tta gaa aaa gcc ata aat ttt cct tcc ttt ccc gat 1392
Ile Gly Ala Ala Leu Glu Lys Ala Ile Asn Phe Pro Ser Phe Pro Asp
                450                455                460

gct ttt aat tag 1404
Ala Phe Asn
465

<210> 10
<211> 467
<212> PRT
<213> Agrobacterium tumefaciens

<400> 10
Met Val Pro Ile Thr Ser Leu Ala Gln Thr Leu Glu Arg Leu Arg Arg
1 5 10 15

```

PF 53790

16

Lys Asp Tyr Ser Cys Leu Glu Leu Val Glu Thr Leu Ile Ala Arg Cys  
                   20                                  25                                  30  
 Gln Ala Ala Lys Pro Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Gly  
                   35                                  40                                  45  
 Leu Arg Arg Ser Ala Lys Lys Ile Asp Arg His Gly Asn Ala Gly Leu  
                   50                                  55                                  60  
 Gly Leu Cys Gly Ile Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly  
                   65                                  70                                  75                                  80  
 Ile Phe Pro Thr Ser Ala Ala Thr Pro Ala Leu Ile Asn His Leu Pro  
                                   85                                  90                                  95  
 Lys Ile Pro Ser Arg Val Ala Glu Arg Leu Phe Ser Ala Gly Ala Leu  
                                   100                                  105                                  110  
 Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser  
                                   115                                  120                                  125  
 Asn Asn Tyr Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu  
                                   130                                  135                                  140  
 Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Ala Val Ala Ser Arg  
                                   145                                  150                                  155                                  160  
 Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu  
                                   165                                  170                                  175  
 Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Leu Ala Arg  
                                   180                                  185                                  190  
 Tyr Pro Arg Asp Arg Ile Ile Pro Val Ser Pro Thr Arg Asp Thr Ala  
                                   195                                  200                                  205  
 Gly Ile Ile Ala Gln Cys Val Ala Asp Val Ile Ile Leu Asp Gln Val  
                                   210                                  215                                  220  
 Ile Ser Gly Arg Ser Ala Lys Ile Ser Pro Met Pro Leu Lys Gly Leu  
                                   225                                  230                                  235                                  240  
 Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp  
                                   245                                  250                                  255  
 Val Ala Phe Ala Ala Glu Thr Thr Ile Arg Leu Leu Ala Asn Arg Gly  
                                   260                                  265                                  270  
 Val Thr Phe Val Glu Ala Asp Ile Pro His Leu Glu Glu Leu Asn Ser  
                                   275                                  280                                  285  
 Gly Ala Ser Leu Pro Ile Ala Leu Tyr Glu Phe Pro His Ala Leu Lys  
                                   290                                  295                                  300  
 Lys Tyr Leu Asp Asp Phe Val Gly Thr Val Ser Phe Ser Asp Val Ile  
                                   305                                  310                                  315                                  320  
 Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Val Ser Ala Gln Ile  
                                   325                                  330                                  335  
 Asp Gly His Gln Ile Ser Asn Asp Glu Tyr Glu Leu Ala Arg Gln Ser  
                                   340                                  345                                  350  
 Phe Arg Pro Arg Leu Gln Ala Thr Tyr Arg Asn Tyr Phe Arg Leu Tyr  
                                   355                                  360                                  365  
 Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Ala Ala Lys Ala  
                                   370                                  375                                  380



PF 53790

17

Ile Gly Gln Glu Ser Ser Val Ile His Asn Gly Ser Met Ile Asn Thr  
 385 390 395 400  
 Phe Lys Ile Tyr Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu  
 405 410 415  
 Pro Gly Leu Ser Leu Pro Ala Cys Leu Thr Pro Asp Arg Leu Pro Val  
 420 425 430  
 Gly Met Glu Ile Asp Gly Leu Ala Gly Ser Asp His Arg Leu Leu Ala  
 435 440 445  
 Ile Gly Ala Ala Leu Glu Lys Ala Ile Asn Phe Pro Ser Phe Pro Asp  
 450 455 460  
 Ala Phe Asn  
 465

&lt;210&gt; 11

&lt;211&gt; 609

&lt;212&gt; DNA

&lt;213&gt; Xanthobacter autotrophicus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(603)

&lt;223&gt; coding for haloalkane dehalogenase

&lt;400&gt; 11

atg tca acg ttt ttt gaa ccg gag aac gga atg aaa caa aac gcc aaa 48  
 Met Ser Thr Phe Phe Glu Pro Glu Asn Gly Met Lys Gln Asn Ala Lys  
 1 5 10 15  
 acc gaa cga atc ctg gat gtc gcg ctc gaa ttg ctt gag aca gag ggt 96  
 Thr Glu Arg Ile Leu Asp Val Ala Leu Glu Leu Leu Glu Thr Glu Gly  
 20 25 30  
 gag ttt ggt ttg acg atg agg cag gtg gca acg caa gcg gac atg tcc 144  
 Glu Phe Gly Leu Thr Met Arg Gln Val Ala Thr Gln Ala Asp Met Ser  
 35 40 45  
 ctg agc aac gtt cag tac tat ttc aag tcc gag gac ctg ctc ctc gtg 192  
 Leu Ser Asn Val Gln Tyr Tyr Phe Lys Ser Glu Asp Leu Leu Leu Val  
 50 55 60  
 gcc atg gca gac cgt tac ttt caa cgg tgc ctg aca acc atg gct gag 240  
 Ala Met Ala Asp Arg Tyr Phe Gln Arg Cys Leu Thr Thr Met Ala Glu  
 65 70 75 80  
 cat ccg ccc tta tcg gca ggg cgt gat caa cac gcc cag tta aga gcg 288  
 His Pro Pro Leu Ser Ala Gly Arg Asp Gln His Ala Gln Leu Arg Ala  
 85 90 95  
 ttg tta cga gaa ctg ctc ggt cat ggt ctt gag att tcc gag atg tgt 336  
 Leu Leu Arg Glu Leu Leu Gly His Gly Leu Glu Ile Ser Glu Met Cys  
 100 105 110  
 cga ata ttc agg gag tac tgg gca atc gcc acc cgt aat gaa act gtt 384  
 Arg Ile Phe Arg Glu Tyr Trp Ala Ile Ala Thr Arg Asn Glu Thr Val  
 115 120 125  
 cac ggc tat ctc aag tcg tac tat cgg gat ctc gcc gaa gtg atg gct 432  
 His Gly Tyr Leu Lys Ser Tyr Tyr Arg Asp Leu Ala Glu Val Met Ala  
 130 135 140

PF 53790

18

```

gag aag ctt gcg cca ctg gcc agc agc gaa aag gcg ctg gcc gtg gcc 480
Glu Lys Leu Ala Pro Leu Ala Ser Ser Glu Lys Ala Leu Ala Val Ala
145          150          155          160

gta tct ttg gtt att cct tat gtt gag ggg tat tcg gta acg gcc att 528
Val Ser Leu Val Ile Pro Tyr Val Glu Gly Tyr Ser Val Thr Ala Ile
          165          170          175

gca atg ccc gaa tcc att gat acg att tcc gag acg ctg acc aat gtg 576
Ala Met Pro Glu Ser Ile Asp Thr Ile Ser Glu Thr Leu Thr Asn Val
          180          185          190

gtg ttg gag cag ctt cgc atc agc aat tcatga 609
Val Leu Glu Gln Leu Arg Ile Ser Asn
          195          200

```

&lt;210&gt; 12

&lt;211&gt; 201

&lt;212&gt; PRT

&lt;213&gt; Xanthobacter autotrophicus

&lt;400&gt; 12

```

Met Ser Thr Phe Phe Glu Pro Glu Asn Gly Met Lys Gln Asn Ala Lys
 1          5          10          15

Thr Glu Arg Ile Leu Asp Val Ala Leu Glu Leu Leu Glu Thr Glu Gly
          20          25          30

Glu Phe Gly Leu Thr Met Arg Gln Val Ala Thr Gln Ala Asp Met Ser
          35          40          45

Leu Ser Asn Val Gln Tyr Tyr Phe Lys Ser Glu Asp Leu Leu Leu Val
          50          55          60

Ala Met Ala Asp Arg Tyr Phe Gln Arg Cys Leu Thr Thr Met Ala Glu
          65          70          75          80

His Pro Pro Leu Ser Ala Gly Arg Asp Gln His Ala Gln Leu Arg Ala
          85          90          95

Leu Leu Arg Glu Leu Leu Gly His Gly Leu Glu Ile Ser Glu Met Cys
          100          105          110

Arg Ile Phe Arg Glu Tyr Trp Ala Ile Ala Thr Arg Asn Glu Thr Val
          115          120          125

His Gly Tyr Leu Lys Ser Tyr Tyr Arg Asp Leu Ala Glu Val Met Ala
          130          135          140

Glu Lys Leu Ala Pro Leu Ala Ser Ser Glu Lys Ala Leu Ala Val Ala
          145          150          155          160

Val Ser Leu Val Ile Pro Tyr Val Glu Gly Tyr Ser Val Thr Ala Ile
          165          170          175

Ala Met Pro Glu Ser Ile Asp Thr Ile Ser Glu Thr Leu Thr Asn Val
          180          185          190

Val Leu Glu Gln Leu Arg Ile Ser Asn
          195          200

```

&lt;210&gt; 13

&lt;211&gt; 1131

&lt;212&gt; DNA

&lt;213&gt; Herpes simplex virus 1

PF 53790

19

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1128)

&lt;223&gt; coding for thymidine kinase (TK)

&lt;400&gt; 13

atg gct tcg tac ccc tgc cat caa cac gcg tct gcg ttc gac cag gct	48
Met Ala Ser Tyr Pro Cys His Gln His Ala Ser Ala Phe Asp Gln Ala	
1 5 10 15	
gcg cgt tct cgc ggc cat agc aac cga cgt acg gcg ttg cgc cct cgc	96
Ala Arg Ser Arg Gly His Ser Asn Arg Arg Thr Ala Leu Arg Pro Arg	
20 25 30	
cgg cag caa gaa gcc acg gaa gtc cgc ctg gag cag aaa atg ccc acg	144
Arg Gln Gln Glu Ala Thr Glu Val Arg Leu Glu Gln Lys Met Pro Thr	
35 40 45	
cta ctg cgg gtt tat ata gac ggt cct cac ggg atg ggg aaa acc acc	192
Leu Leu Arg Val Tyr Ile Asp Gly Pro His Gly Met Gly Lys Thr Thr	
50 55 60	
acc acg caa ctg ctg gtg gcc ctg ggt tcg cgc gac gat atc gtc tac	240
Thr Thr Gln Leu Leu Val Ala Leu Gly Ser Arg Asp Asp Ile Val Tyr	
65 70 75 80	
gta ccc gag ccg atg act tac tgg cag gtg ctg ggg gct tcc gag aca	288
Val Pro Glu Pro Met Thr Tyr Trp Gln Val Leu Gly Ala Ser Glu Thr	
85 90 95	
atc gcg aac atc tac acc aca caa cac cgc ctc gac cag ggt gag ata	336
Ile Ala Asn Ile Tyr Thr Thr Gln His Arg Leu Asp Gln Gly Glu Ile	
100 105 110	
tcg gcc ggg gac gcg gcg gtg gta atg aca agc gcc cag ata aca atg	384
Ser Ala Gly Asp Ala Ala Val Val Met Thr Ser Ala Gln Ile Thr Met	
115 120 125	
ggc atg cct tat gcc gtg acc gac gcc gtt ctg gct cct cat gtc ggg	432
Gly Met Pro Tyr Ala Val Thr Asp Ala Val Leu Ala Pro His Val Gly	
130 135 140	
ggg gag gct ggg agt tca cat gcc ccg ccc ccg gcc ctc acc ctc atc	480
Gly Glu Ala Gly Ser Ser His Ala Pro Pro Pro Ala Leu Thr Leu Ile	
145 150 155 160	
ttc gac cgc cat ccc atc gcc gcc ctc ctg tgc tac ccg gcc gcg cga	528
Phe Asp Arg His Pro Ile Ala Ala Leu Leu Cys Tyr Pro Ala Ala Arg	
165 170 175	
tac ctt atg ggc agc atg acc ccc cag gcc gtg ctg gcg ttc gtg gcc	576
Tyr Leu Met Gly Ser Met Thr Pro Gln Ala Val Leu Ala Phe Val Ala	
180 185 190	
ctc atc ccg ccg acc ttg ccc ggc aca aac atc gtg ttg ggg gcc ctt	624
Leu Ile Pro Pro Thr Leu Pro Gly Thr Asn Ile Val Leu Gly Ala Leu	
195 200 205	
ccg gag gac aga cac atc gac cgc ctg gcc aaa cgc cag cgc ccc ggc	672
Pro Glu Asp Arg His Ile Asp Arg Leu Ala Lys Arg Gln Arg Pro Gly	
210 215 220	
gag cgg ctt gac ctg gct atg ctg gcc gcg att cgc cgc gtt tac ggg	720
Glu Arg Leu Asp Leu Ala Met Leu Ala Ala Ile Arg Arg Val Tyr Gly	
225 230 235 240	

PF 53790

20

```

ctg ctt gcc aat acg gtg cgg tat ctg cag ggc ggc ggg tcg tgg tgg 768
Leu Leu Ala Asn Thr Val Arg Tyr Leu Gln Gly Gly Gly Ser Trp Trp
                245                250                255

gag gat tgg gga cag ctt tcg ggg acg gcc gtg ccg ccc cag ggt gcc 816
Glu Asp Trp Gly Gln Leu Ser Gly Thr Ala Val Pro Pro Gln Gly Ala
                260                265                270

gag ccc cag agc aac gcg ggc cca cga ccc cat atc ggg gac acg tta 864
Glu Pro Gln Ser Asn Ala Gly Pro Arg Pro His Ile Gly Asp Thr Leu
                275                280                285

ttt acc ctg ttt cgg gcc ccc gag ttg ctg gcc ccc aac ggc gac ctg 912
Phe Thr Leu Phe Arg Ala Pro Glu Leu Leu Ala Pro Asn Gly Asp Leu
                290                295                300

tat aac gtg ttt gcc tgg gcc ttg gac gtc ttg gcc aaa cgc ctc cgt 960
Tyr Asn Val Phe Ala Trp Ala Leu Asp Val Leu Ala Lys Arg Leu Arg
305                310                315                320

ccc atg cac gtc ttt atc ctg gat tac gac caa tcg ccc gcc ggc tgc 1008
Pro Met His Val Phe Ile Leu Asp Tyr Asp Gln Ser Pro Ala Gly Cys
                325                330                335

cgg gac gcc ctg ctg caa ctt acc tcc ggg atg gtc cag acc cac gtc 1056
Arg Asp Ala Leu Leu Gln Leu Thr Ser Gly Met Val Gln Thr His Val
                340                345                350

acc acc cca ggc tcc ata ccg acg atc tgc gac ctg gcg cgc acg ttt 1104
Thr Thr Pro Gly Ser Ile Pro Thr Ile Cys Asp Leu Ala Arg Thr Phe
                355                360                365

gcc cgg gag atg ggg gag gct aac tga 1131
Ala Arg Glu Met Gly Glu Ala Asn
370                375

<210> 14
<211> 376
<212> PRT
<213> Herpes simplex virus 1

<400> 14
Met Ala Ser Tyr Pro Cys His Gln His Ala Ser Ala Phe Asp Gln Ala
 1          5          10          15
Ala Arg Ser Arg Gly His Ser Asn Arg Arg Thr Ala Leu Arg Pro Arg
 20          25          30
Arg Gln Gln Glu Ala Thr Glu Val Arg Leu Glu Gln Lys Met Pro Thr
 35          40          45
Leu Leu Arg Val Tyr Ile Asp Gly Pro His Gly Met Gly Lys Thr Thr
 50          55          60
Thr Thr Gln Leu Leu Val Ala Leu Gly Ser Arg Asp Asp Ile Val Tyr
 65          70          75          80
Val Pro Glu Pro Met Thr Tyr Trp Gln Val Leu Gly Ala Ser Glu Thr
 85          90          95
Ile Ala Asn Ile Tyr Thr Thr Gln His Arg Leu Asp Gln Gly Glu Ile
100          105          110
Ser Ala Gly Asp Ala Ala Val Val Met Thr Ser Ala Gln Ile Thr Met
115          120          125

```

PF 53790

21

Gly Met Pro Tyr Ala Val Thr Asp Ala Val Leu Ala Pro His Val Gly  
 130 135 140  
 Gly Glu Ala Gly Ser Ser His Ala Pro Pro Pro Ala Leu Thr Leu Ile  
 145 150 155 160  
 Phe Asp Arg His Pro Ile Ala Ala Leu Leu Cys Tyr Pro Ala Ala Arg  
 165 170 175  
 Tyr Leu Met Gly Ser Met Thr Pro Gln Ala Val Leu Ala Phe Val Ala  
 180 185 190  
 Leu Ile Pro Pro Thr Leu Pro Gly Thr Asn Ile Val Leu Gly Ala Leu  
 195 200 205  
 Pro Glu Asp Arg His Ile Asp Arg Leu Ala Lys Arg Gln Arg Pro Gly  
 210 215 220  
 Glu Arg Leu Asp Leu Ala Met Leu Ala Ala Ile Arg Arg Val Tyr Gly  
 225 230 235 240  
 Leu Leu Ala Asn Thr Val Arg Tyr Leu Gln Gly Gly Gly Ser Trp Trp  
 245 250 255  
 Glu Asp Trp Gly Gln Leu Ser Gly Thr Ala Val Pro Pro Gln Gly Ala  
 260 265 270  
 Glu Pro Gln Ser Asn Ala Gly Pro Arg Pro His Ile Gly Asp Thr Leu  
 275 280 285  
 Phe Thr Leu Phe Arg Ala Pro Glu Leu Leu Ala Pro Asn Gly Asp Leu  
 290 295 300  
 Tyr Asn Val Phe Ala Trp Ala Leu Asp Val Leu Ala Lys Arg Leu Arg  
 305 310 315 320  
 Pro Met His Val Phe Ile Leu Asp Tyr Asp Gln Ser Pro Ala Gly Cys  
 325 330 335  
 Arg Asp Ala Leu Leu Gln Leu Thr Ser Gly Met Val Gln Thr His Val  
 340 345 350  
 Thr Thr Pro Gly Ser Ile Pro Thr Ile Cys Asp Leu Ala Arg Thr Phe  
 355 360 365  
 Ala Arg Glu Met Gly Glu Ala Asn  
 370 375

&lt;210&gt; 15

&lt;211&gt; 1131

&lt;212&gt; DNA

&lt;213&gt; Herpes simplex virus 1

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1128)

&lt;223&gt; coding for thymidine kinase (TK)

&lt;400&gt; 15

atg gct tcg tac ccc tgc cat caa cac gcg tct gcg ttc gac cag gct 48  
 Met Ala Ser Tyr Pro Cys His Gln His Ala Ser Ala Phe Asp Gln Ala  
 1 5 10 15

gcg cgt tct cgc ggc cat agc aac cga cgt acg gcg ttg cgc cct cgc 96  
 Ala Arg Ser Arg Gly His Ser Asn Arg Arg Thr Ala Leu Arg Pro Arg  
 20 25 30

PF 53790

22

cgg cag caa gaa gcc acg gaa gtc cgc ctg gag cag aaa atg ccc acg	144
Arg Gln Gln Glu Ala Thr Glu Val Arg Leu Glu Gln Lys Met Pro Thr	
35 40 45	
cta ctg cgg gtt tat ata gac ggt cct cac ggg atg ggg aaa acc acc	192
Leu Leu Arg Val Tyr Ile Asp Gly Pro His Gly Met Gly Lys Thr Thr	
50 55 60	
acc acg caa ctg ctg gtg gcc ctg ggt tcg cgc gac gat atc gtc tac	240
Thr Thr Gln Leu Leu Val Ala Leu Gly Ser Arg Asp Asp Ile Val Tyr	
65 70 75 80	
gta ccc gag ccg atg act tac tgg cag gtg ctg ggg gct tcc gag aca	288
Val Pro Glu Pro Met Thr Tyr Trp Gln Val Leu Gly Ala Ser Glu Thr	
85 90 95	
atc gcg aac atc tac acc aca caa cac cgc ctc gac cag ggt gag ata	336
Ile Ala Asn Ile Tyr Thr Thr Gln His Arg Leu Asp Gln Gly Glu Ile	
100 105 110	
tcg gcc ggg gac gcg gcg gtg gta atg aca agc gcc cag ata aca atg	384
Ser Ala Gly Asp Ala Ala Val Val Met Thr Ser Ala Gln Ile Thr Met	
115 120 125	
ggc atg cct tat gcc gtg acc gac gcc gtt ctg gct cct cat gtc ggg	432
Gly Met Pro Tyr Ala Val Thr Asp Ala Val Leu Ala Pro His Val Gly	
130 135 140	
ggg gag gct ggg agt tca cat gcc ccg ccc ccg gcc ctc acc ctc atc	480
Gly Glu Ala Gly Ser His Ala Pro Pro Pro Ala Leu Thr Leu Ile	
145 150 155 160	
ttc gac cgc cat ccc atc gcc gcc ctc ctg tgc tac ccg gcc gcg cga	528
Phe Asp Arg His Pro Ile Ala Ala Leu Leu Cys Tyr Pro Ala Ala Arg	
165 170 175	
tac ctt atg ggc agc atg acc ccc cag gcc gtg ctg gcg ttc gtg gcc	576
Tyr Leu Met Gly Ser Met Thr Pro Gln Ala Val Leu Ala Phe Val Ala	
180 185 190	
ctc atc ccg ccg acc ttg ccc ggc aca aac atc gtg ttg ggg gcc ctt	624
Leu Ile Pro Pro Thr Leu Pro Gly Thr Asn Ile Val Leu Gly Ala Leu	
195 200 205	
ccg gag gac aga cac atc gac cgc ctg gcc aaa cgc cag cgc ccc ggc	672
Pro Glu Asp Arg His Ile Asp Arg Leu Ala Lys Arg Gln Arg Pro Gly	
210 215 220	
gag cgg ctt gac ctg gct atg ctg gcc gcg att cgc cgc gtt tac ggg	720
Glu Arg Leu Asp Leu Ala Met Leu Ala Ala Ile Arg Arg Val Tyr Gly	
225 230 235 240	
ctg ctt gcc aat acg gtg cgg tat ctg cag ggc ggc ggg tcg tgg tgg	768
Leu Leu Ala Asn Thr Val Arg Tyr Leu Gln Gly Gly Gly Ser Trp Trp	
245 250 255	
gag gat tgg gga cag ctt tcg ggg acg gcc gtg ccg ccc cag ggt gcc	816
Glu Asp Trp Gly Gln Leu Ser Gly Thr Ala Val Pro Pro Gln Gly Ala	
260 265 270	
gag ccc cag agc aac gcg ggc cca cga ccc cat atc ggg gac acg tta	864
Glu Pro Gln Ser Asn Ala Gly Pro Arg Pro His Ile Gly Asp Thr Leu	
275 280 285	
ttt acc ctg ttt cgg gcc ccc gag ttg ctg gcc ccc aac ggc gac ctg	912
Phe Thr Leu Phe Arg Ala Pro Glu Leu Leu Ala Pro Asn Gly Asp Leu	
290 295 300	

PF 53790

23

tat aac gtg ttt gcc tgg gcc ttg gac gtc ttg gcc aaa cgc ctc cgt 960  
 Tyr Asn Val Phe Ala Trp Ala Leu Asp Val Leu Ala Lys Arg Leu Arg  
 305 310 315 320  
 ccc atg cac gtc ttt atc ctg gat tac gac caa tcg ccc gcc ggc tgc 1008  
 Pro Met His Val Phe Ile Leu Asp Tyr Asp Gln Ser Pro Ala Gly Cys  
 325 330 335  
 cgg gac gcc ctg ctg caa ctt acc tcc ggg atg gtc cag acc cac gtc 1056  
 Arg Asp Ala Leu Leu Gln Leu Thr Ser Gly Met Val Gln Thr His Val  
 340 345 350  
 acc acc cca ggc tcc ata ccg acg atc tgc gac ctg gcg cgc acg ttt 1104  
 Thr Thr Pro Gly Ser Ile Pro Thr Ile Cys Asp Leu Ala Arg Thr Phe  
 355 360 365  
 gcc cgg gag atg ggg gag gct aac tga 1131  
 Ala Arg Glu Met Gly Glu Ala Asn  
 370 375  
 <210> 16  
 <211> 376  
 <212> PRT  
 <213> Herpes simplex virus 1  
 <400> 16  
 Met Ala Ser Tyr Pro Cys His Gln His Ala Ser Ala Phe Asp Gln Ala  
 1 5 10 15  
 Ala Arg Ser Arg Gly His Ser Asn Arg Arg Thr Ala Leu Arg Pro Arg  
 20 25 30  
 Arg Gln Gln Glu Ala Thr Glu Val Arg Leu Glu Gln Lys Met Pro Thr  
 35 40 45  
 Leu Leu Arg Val Tyr Ile Asp Gly Pro His Gly Met Gly Lys Thr Thr  
 50 55 60  
 Thr Thr Gln Leu Leu Val Ala Leu Gly Ser Arg Asp Asp Ile Val Tyr  
 65 70 75 80  
 Val Pro Glu Pro Met Thr Tyr Trp Gln Val Leu Gly Ala Ser Glu Thr  
 85 90 95  
 Ile Ala Asn Ile Tyr Thr Thr Gln His Arg Leu Asp Gln Gly Glu Ile  
 100 105 110  
 Ser Ala Gly Asp Ala Ala Val Val Met Thr Ser Ala Gln Ile Thr Met  
 115 120 125  
 Gly Met Pro Tyr Ala Val Thr Asp Ala Val Leu Ala Pro His Val Gly  
 130 135 140  
 Gly Glu Ala Gly Ser Ser His Ala Pro Pro Pro Ala Leu Thr Leu Ile  
 145 150 155 160  
 Phe Asp Arg His Pro Ile Ala Ala Leu Leu Cys Tyr Pro Ala Ala Arg  
 165 170 175  
 Tyr Leu Met Gly Ser Met Thr Pro Gln Ala Val Leu Ala Phe Val Ala  
 180 185 190  
 Leu Ile Pro Pro Thr Leu Pro Gly Thr Asn Ile Val Leu Gly Ala Leu  
 195 200 205  
 Pro Glu Asp Arg His Ile Asp Arg Leu Ala Lys Arg Gln Arg Pro Gly  
 210 215 220

PF 53790

24

Glu Arg Leu Asp Leu Ala Met Leu Ala Ala Ile Arg Arg Val Tyr Gly  
 225 230 235 240  
 Leu Leu Ala Asn Thr Val Arg Tyr Leu Gln Gly Gly Gly Ser Trp Trp  
 245 250 255  
 Glu Asp Trp Gly Gln Leu Ser Gly Thr Ala Val Pro Pro Gln Gly Ala  
 260 265 270  
 Glu Pro Gln Ser Asn Ala Gly Pro Arg Pro His Ile Gly Asp Thr Leu  
 275 280 285  
 Phe Thr Leu Phe Arg Ala Pro Glu Leu Leu Ala Pro Asn Gly Asp Leu  
 290 295 300  
 Tyr Asn Val Phe Ala Trp Ala Leu Asp Val Leu Ala Lys Arg Leu Arg  
 305 310 315 320  
 Pro Met His Val Phe Ile Leu Asp Tyr Asp Gln Ser Pro Ala Gly Cys  
 325 330 335  
 Arg Asp Ala Leu Leu Gln Leu Thr Ser Gly Met Val Gln Thr His Val  
 340 345 350  
 Thr Thr Pro Gly Ser Ile Pro Thr Ile Cys Asp Leu Ala Arg Thr Phe  
 355 360 365  
 Ala Arg Glu Met Gly Glu Ala Asn  
 370 375

&lt;210&gt; 17

&lt;211&gt; 840

&lt;212&gt; DNA

<213> *Toxoplasma gondii*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(837)

 <223> coding for hypoxanthine-xanthine-guanine  
 phosphoribosyl transferase (HXGPRTase)

&lt;400&gt; 17

atg gcg tcc aaa ccc att gaa gaa tcc cgg tcg caa aaa cgg agt gcc 48  
 Met Ala Ser Lys Pro Ile Glu Glu Ser Arg Ser Gln Lys Arg Ser Ala  
 1 5 10 15  
 ttc tca gac atc ttc tgt tgt tgc act cct aat gaa ggg gct atc gtg 96  
 Phe Ser Asp Ile Phe Cys Cys Cys Thr Pro Asn Glu Gly Ala Ile Val  
 20 25 30  
 ccc agt gac cca atg gtc tcc acc agt gct cca gca cgc acc agt gct 144  
 Pro Ser Asp Pro Met Val Ser Thr Ser Ala Pro Ala Arg Thr Ser Ala  
 35 40 45  
 cca gcg cgc tcc agt gca ctt caa gac tac ggc aag ggc aag ggc cgt 192  
 Pro Ala Arg Ser Ser Ala Leu Gln Asp Tyr Gly Lys Gly Lys Gly Arg  
 50 55 60  
 att gag ccc atg tat atc ccc gac aac acc ttc tac aac gct gat gac 240  
 Ile Glu Pro Met Tyr Ile Pro Asp Asn Thr Phe Tyr Asn Ala Asp Asp  
 65 70 75 80  
 ttt ctt gtg ccc ccc cac tgc aag ccc tac att gac aaa atc ctc ctc 288  
 Phe Leu Val Pro Pro His Cys Lys Pro Tyr Ile Asp Lys Ile Leu Leu  
 85 90 95



PF 53790

25

```

cct ggt gga ttg gtc aag gac aga gtt gag aag ttg gcg tat gac atc 336
Pro Gly Gly Leu Val Lys Asp Arg Val Glu Lys Leu Ala Tyr Asp Ile
      100      105      110

cac aga act tac ttc ggc gag gag ttg cac atc att tgc atc ctg aaa 384
His Arg Thr Tyr Phe Gly Glu Glu Leu His Ile Ile Cys Ile Leu Lys
      115      120      125

ggc tct cgc ggc ttc ttc aac ctt ctg atc gac tac ctt gcc acc ata 432
Gly Ser Arg Gly Phe Phe Asn Leu Leu Ile Asp Tyr Leu Ala Thr Ile
      130      135      140

cag aag tac agt ggt cgt gag tcc agc gtg ccc ccc ttc ttc gag cac 480
Gln Lys Tyr Ser Gly Arg Glu Ser Ser Val Pro Pro Phe Phe Glu His
      145      150      155      160

tat gtc cgc ctg aag tcc tac cag aac gac aac agc aca ggc cag ctc 528
Tyr Val Arg Leu Lys Ser Tyr Gln Asn Asp Asn Ser Thr Gly Gln Leu
      165      170      175

acc gtc ttg agc gac gac ttg tca atc ttt cgc gac aag cac gtt ctg 576
Thr Val Leu Ser Asp Asp Leu Ser Ile Phe Arg Asp Lys His Val Leu
      180      185      190

att gtt gag gac atc gtc gac acc ggt ttc acc ctc acc gag ttc ggt 624
Ile Val Glu Asp Ile Val Asp Thr Gly Phe Thr Leu Thr Glu Phe Gly
      195      200      205

gag cgc ctg aaa gcc gtc ggt ccc aag tcg atg aga atc gcc acc ctc 672
Glu Arg Leu Lys Ala Val Gly Pro Lys Ser Met Arg Ile Ala Thr Leu
      210      215      220

gtc gag aag cgc aca gat cgc tcc aac agc ttg aag ggc gac ttc gtc 720
Val Glu Lys Arg Thr Asp Arg Ser Asn Ser Leu Lys Gly Asp Phe Val
      225      230      235      240

ggc ttc agc att gaa gac gtc tgg atc gtt ggt tgc tgc tac gac ttc 768
Gly Phe Ser Ile Glu Asp Val Trp Ile Val Gly Cys Cys Tyr Asp Phe
      245      250      255

aac gag atg ttc cgc gac ttc gac cac gtc gcc gtc ctg agc gac gcc 816
Asn Glu Met Phe Arg Asp Phe Asp His Val Ala Val Leu Ser Asp Ala
      260      265      270

gct cgc aaa aag ttc gag aag taa 840
Ala Arg Lys Lys Phe Glu Lys
      275

```

&lt;210&gt; 18

&lt;211&gt; 279

&lt;212&gt; PRT

&lt;213&gt; Toxoplasma gondii

&lt;400&gt; 18

```

Met Ala Ser Lys Pro Ile Glu Glu Ser Arg Ser Gln Lys Arg Ser Ala
 1          5          10          15

Phe Ser Asp Ile Phe Cys Cys Cys Thr Pro Asn Glu Gly Ala Ile Val
      20      25      30

Pro Ser Asp Pro Met Val Ser Thr Ser Ala Pro Ala Arg Thr Ser Ala
      35      40      45

Pro Ala Arg Ser Ser Ala Leu Gln Asp Tyr Gly Lys Gly Lys Gly Arg
      50      55      60

```

PF 53790

26

Ile Glu Pro Met Tyr Ile Pro Asp Asn Thr Phe Tyr Asn Ala Asp Asp  
 65 70 75 80  
 Phe Leu Val Pro Pro His Cys Lys Pro Tyr Ile Asp Lys Ile Leu Leu  
 85 90 95  
 Pro Gly Gly Leu Val Lys Asp Arg Val Glu Lys Leu Ala Tyr Asp Ile  
 100 105 110  
 His Arg Thr Tyr Phe Gly Glu Glu Leu His Ile Ile Cys Ile Leu Lys  
 115 120 125  
 Gly Ser Arg Gly Phe Phe Asn Leu Leu Ile Asp Tyr Leu Ala Thr Ile  
 130 135 140  
 Gln Lys Tyr Ser Gly Arg Glu Ser Ser Val Pro Pro Phe Phe Glu His  
 145 150 155 160  
 Tyr Val Arg Leu Lys Ser Tyr Gln Asn Asp Asn Ser Thr Gly Gln Leu  
 165 170 175  
 Thr Val Leu Ser Asp Asp Leu Ser Ile Phe Arg Asp Lys His Val Leu  
 180 185 190  
 Ile Val Glu Asp Ile Val Asp Thr Gly Phe Thr Leu Thr Glu Phe Gly  
 195 200 205  
 Glu Arg Leu Lys Ala Val Gly Pro Lys Ser Met Arg Ile Ala Thr Leu  
 210 215 220  
 Val Glu Lys Arg Thr Asp Arg Ser Asn Ser Leu Lys Gly Asp Phe Val  
 225 230 235 240  
 Gly Phe Ser Ile Glu Asp Val Trp Ile Val Gly Cys Cys Tyr Asp Phe  
 245 250 255  
 Asn Glu Met Phe Arg Asp Phe Asp His Val Ala Val Leu Ser Asp Ala  
 260 265 270  
 Ala Arg Lys Lys Phe Glu Lys  
 275

&lt;210&gt; 19

&lt;211&gt; 459

&lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(456)

 <223> coding for xanthine-guanine phosphoribosyl  
 transferase (gpt)

&lt;400&gt; 19

atg agc gaa aaa tac atc gtc acc tgg gac atg ttg cag atc cat gca 48  
 Met Ser Glu Lys Tyr Ile Val Thr Trp Asp Met Leu Gln Ile His Ala  
 1 5 10 15  
 cgt aaa ctc gca agc cga ctg atg cct tct gaa caa tgg aaa ggc att 96  
 Arg Lys Leu Ala Ser Arg Leu Met Pro Ser Glu Gln Trp Lys Gly Ile  
 20 25 30  
 att gcc gta agc cgt ggc ggt ctg gta ccg ggt gcg tta ctg gcg cgt 144  
 Ile Ala Val Ser Arg Gly Gly Leu Val Pro Gly Ala Leu Leu Ala Arg  
 35 40 45

PF 53790

27

gaa ctg ggt att cgt cat gtc gat acc gtt tgt att tcc agc tac gat	192
Glu Leu Gly Ile Arg His Val Asp Thr Val Cys Ile Ser Ser Tyr Asp	
50 55 60	
cac gac aac cag cgc gag ctt aaa gtg ctg aaa cgc gca gaa ggc gat	240
His Asp Asn Gln Arg Glu Leu Lys Val Leu Lys Arg Ala Glu Gly Asp	
65 70 75 80	
ggc gaa ggc ttc atc gtt att gat gac ctg gtg gat acc ggt ggt act	288
Gly Glu Gly Phe Ile Val Ile Asp Asp Leu Val Asp Thr Gly Gly Thr	
85 90 95	
gcg gtt gcg att cgt gaa atg tat cca aaa gcg cac ttt gtc acc atc	336
Ala Val Ala Ile Arg Glu Met Tyr Pro Lys Ala His Phe Val Thr Ile	
100 105 110	
ttc gca aaa ccg gct ggt cgt ccg ctg gtt gat gac tat gtt gtt gat	384
Phe Ala Lys Pro Ala Gly Arg Pro Leu Val Asp Asp Tyr Val Val Asp	
115 120 125	
atc ccg caa gat acc tgg att gaa cag ccg tgg gat atg ggc gtc gta	432
Ile Pro Gln Asp Thr Trp Ile Glu Gln Pro Trp Asp Met Gly Val Val	
130 135 140	
ttc gtc ccg cca atc tcc ggt cgc taa	459
Phe Val Pro Pro Ile Ser Gly Arg	
145 150	
<210> 20	
<211> 152	
<212> PRT	
<213> Escherichia coli	
<400> 20	
Met Ser Glu Lys Tyr Ile Val Thr Trp Asp Met Leu Gln Ile His Ala	
1 5 10 15	
Arg Lys Leu Ala Ser Arg Leu Met Pro Ser Glu Gln Trp Lys Gly Ile	
20 25 30	
Ile Ala Val Ser Arg Gly Gly Leu Val Pro Gly Ala Leu Leu Ala Arg	
35 40 45	
Glu Leu Gly Ile Arg His Val Asp Thr Val Cys Ile Ser Ser Tyr Asp	
50 55 60	
His Asp Asn Gln Arg Glu Leu Lys Val Leu Lys Arg Ala Glu Gly Asp	
65 70 75 80	
Gly Glu Gly Phe Ile Val Ile Asp Asp Leu Val Asp Thr Gly Gly Thr	
85 90 95	
Ala Val Ala Ile Arg Glu Met Tyr Pro Lys Ala His Phe Val Thr Ile	
100 105 110	
Phe Ala Lys Pro Ala Gly Arg Pro Leu Val Asp Asp Tyr Val Val Asp	
115 120 125	
Ile Pro Gln Asp Thr Trp Ile Glu Gln Pro Trp Asp Met Gly Val Val	
130 135 140	
Phe Val Pro Pro Ile Ser Gly Arg	
145 150	

PF 53790

28

<210> 21  
 <211> 459  
 <212> DNA  
 <213> Escherichia coli

<220>  
 <221> CDS  
 <222> (1)..(456)  
 <223> coding for xanthine-guanine phosphoribosyl  
 transferase (gpt)

<400> 21  
 atg agc gaa aaa tac atc gtc acc tgg gac atg ttg cag atc cat gca 48  
 Met Ser Glu Lys Tyr Ile Val Thr Trp Asp Met Leu Gln Ile His Ala  
 1 5 10 15  
 cgt aaa ctc gca agc cga ctg atg cct tct gaa caa tgg aaa ggc att 96  
 Arg Lys Leu Ala Ser Arg Leu Met Pro Ser Glu Gln Trp Lys Gly Ile  
 20 25 30  
 att gcc gta agc cgt ggc ggt ctg gta ccg ggt gcg tta ctg gcg cgt 144  
 Ile Ala Val Ser Arg Gly Gly Leu Val Pro Gly Ala Leu Leu Ala Arg  
 35 40 45  
 gaa ctg ggt att cgt cat gtc gat acc gtt tgt att tcc agc tac gat 192  
 Glu Leu Gly Ile Arg His Val Asp Thr Val Cys Ile Ser Ser Tyr Asp  
 50 55 60  
 cac gac aac cag cgc gag ctt aaa gtg ctg aaa cgc gca gaa ggc gat 240  
 His Asp Asn Gln Arg Glu Leu Lys Val Leu Lys Arg Ala Glu Gly Asp  
 65 70 75 80  
 ggc gaa ggc ttc atc gtt att gat gac ctg gtg gat acc ggt ggt act 288  
 Gly Glu Gly Phe Ile Val Ile Asp Asp Leu Val Asp Thr Gly Gly Thr  
 85 90 95  
 gcg gtt gcg att cgt gaa atg tat cca aaa gcg cac ttt gtc acc atc 336  
 Ala Val Ala Ile Arg Glu Met Tyr Pro Lys Ala His Phe Val Thr Ile  
 100 105 110  
 ttc gca aaa ccg gct ggt cgt ccg ctg gtt gat gac tat gtt gtt gat 384  
 Phe Ala Lys Pro Ala Gly Arg Pro Leu Val Asp Asp Tyr Val Val Asp  
 115 120 125  
 atc ccg caa gat acc tgg att gaa cag ccg tgg gat atg ggc gtc gta 432  
 Ile Pro Gln Asp Thr Trp Ile Glu Gln Pro Trp Asp Met Gly Val Val  
 130 135 140  
 ttc gtc ccg cca atc tcc ggt cgc taa 459  
 Phe Val Pro Pro Ile Ser Gly Arg  
 145 150

<210> 22  
 <211> 152  
 <212> PRT  
 <213> Escherichia coli

<400> 22  
 Met Ser Glu Lys Tyr Ile Val Thr Trp Asp Met Leu Gln Ile His Ala  
 1 5 10 15  
 Arg Lys Leu Ala Ser Arg Leu Met Pro Ser Glu Gln Trp Lys Gly Ile  
 20 25 30

29

Ile	Ala	Val	Ser	Arg	Gly	Gly	Leu	Val	Pro	Gly	Ala	Leu	Leu	Ala	Arg
		35					40					45			
Glu	Leu	Gly	Ile	Arg	His	Val	Asp	Thr	Val	Cys	Ile	Ser	Ser	Tyr	Asp
		50					55				60				
His	Asp	Asn	Gln	Arg	Glu	Leu	Lys	Val	Leu	Lys	Arg	Ala	Glu	Gly	Asp
						70					75				80
Gly	Glu	Gly	Phe	Ile	Val	Ile	Asp	Asp	Leu	Val	Asp	Thr	Gly	Gly	Thr
						85				90				95	
Ala	Val	Ala	Ile	Arg	Glu	Met	Tyr	Pro	Lys	Ala	His	Phe	Val	Thr	Ile
								105					110		
Phe	Ala	Lys	Pro	Ala	Gly	Arg	Pro	Leu	Val	Asp	Asp	Tyr	Val	Val	Asp
							120					125			
Ile	Pro	Gln	Asp	Thr	Trp	Ile	Glu	Gln	Pro	Trp	Asp	Met	Gly	Val	Val
							135					140			
Phe	Val	Pro	Pro	Ile	Ser	Gly	Arg								
							150								

<210> 23

**<211> 720**

**<212> DNA**

<213> Escherichia coli

**<220>**

**<221> CDS**

<222> (1)..(717)

<223> coding for purine nucleoside phosphorylase (deoD)

<400> 23

atg gct acc cca cac att aat gca gaa atg ggc gat ttc gct gac gta 48  
Met Ala Thr Pro His Ile Asn Ala Glu Met Gly Asp Phe Ala Asp Val  
1 5 10 15

ggt ttg atg cca ggc gac ccg ctg cgt gcg aag tat att gct gaa act 96  
Val Leu Met Pro Gly Asp Pro Leu Arg Ala Lys Tyr Ile Ala Glu Thr  
20 25 30

ttc ctt gaa gat gcc cgt gaa gtg aac aac gtt cgc ggt atg ctg ggc 144  
Phe Leu Glu Asp Ala Arg Glu Val Asn Asn Val Arg Gly Met Leu Gly  
35 40 45

ttc acc ggt act tac aaa ggc cgc aaa att tcc gta atg ggt cac ggt 192  
Phe Thr Gly Thr Tyr Lys Gly Arg Lys Ile Ser Val Met Gly His Gly  
50 55 60

atg ggt atc ccg tcc tgc tcc atc tac acc aaa gaa ctg atc acc gat 240  
Met Gly Ile Pro Ser Cys Ser Ile Tyr Thr Lys Glu Leu Ile Thr Asp  
65 70 75 80

ttc ggc gtg aag aaa att atc cgc gtg ggt tcc tgt ggc gca gtt ctg 288  
Phe Gly Val Lys Lys Ile Ile Arg Val Gly Ser Cys Gly Ala Val Leu  
85 90 95

ccg cac gta aaa ctg cgc gac gtc gtt atc ggt atg ggt gcc tgc acc 336  
Pro His Val Lys Leu Arg Asp Val Val Ile Gly Met Gly Ala Cys Thr  
100 105 110

gat tcc aaa gtt aac cgc atc cgt ttt aaa gac cat gac ttt gcc gct 384  
Asp Ser Lys Val Asn Arg Ile Arg Phe Lys Asp His Asp Phe Ala Ala  
115 120 125

PF 53790

30

atc gct gac ttc gac atg gtg cgt aac gca gta gat gca gct aaa gca 432  
 Ile Ala Asp Phe Asp Met Val Arg Asn Ala Val Asp Ala Ala Lys Ala  
 130 135 140

ctg ggt att gat gct cgc gtg ggt aac ctg ttc tcc gct gac ctg ttc 480  
 Leu Gly Ile Asp Ala Arg Val Gly Asn Leu Phe Ser Ala Asp Leu Phe  
 145 150 155 160

tac tct ccg gac ggc gaa atg ttc gac gtg atg gaa aaa tac ggc att 528  
 Tyr Ser Pro Asp Gly Glu Met Phe Asp Val Met Glu Lys Tyr Gly Ile  
 165 170 175

ctc ggc gtg gaa atg gaa gcg gct ggt atc tac ggc gtc gct gca gaa 576  
 Leu Gly Val Glu Met Glu Ala Ala Gly Ile Tyr Gly Val Ala Ala Glu  
 180 185 190

ttt ggc gcg aaa gcc ctg acc atc tgc acc gta tct gac cac atc cgc 624  
 Phe Gly Ala Lys Ala Leu Thr Ile Cys Thr Val Ser Asp His Ile Arg  
 195 200 205

act cac gag cag acc act gcc gct gag cgt cag act acc ttc aac gac 672  
 Thr His Glu Gln Thr Thr Ala Ala Glu Arg Gln Thr Thr Phe Asn Asp  
 210 215 220

atg atc aaa atc gca ctg gaa tcc gtt ctg ctg ggc gat aaa gag taa 720  
 Met Ile Lys Ile Ala Leu Glu Ser Val Leu Leu Gly Asp Lys Glu  
 225 230 235

<210> 24  
 <211> 239  
 <212> PRT  
 <213> Escherichia coli

<400> 24  
 Met Ala Thr Pro His Ile Asn Ala Glu Met Gly Asp Phe Ala Asp Val  
 1 5 10 15  
 Val Leu Met Pro Gly Asp Pro Leu Arg Ala Lys Tyr Ile Ala Glu Thr  
 20 25 30  
 Phe Leu Glu Asp Ala Arg Glu Val Asn Asn Val Arg Gly Met Leu Gly  
 35 40 45  
 Phe Thr Gly Thr Tyr Lys Gly Arg Lys Ile Ser Val Met Gly His Gly  
 50 55 60  
 Met Gly Ile Pro Ser Cys Ser Ile Tyr Thr Lys Glu Leu Ile Thr Asp  
 65 70 75 80  
 Phe Gly Val Lys Lys Ile Ile Arg Val Gly Ser Cys Gly Ala Val Leu  
 85 90 95  
 Pro His Val Lys Leu Arg Asp Val Val Ile Gly Met Gly Ala Cys Thr  
 100 105 110  
 Asp Ser Lys Val Asn Arg Ile Arg Phe Lys Asp His Asp Phe Ala Ala  
 115 120 125  
 Ile Ala Asp Phe Asp Met Val Arg Asn Ala Val Asp Ala Ala Lys Ala  
 130 135 140  
 Leu Gly Ile Asp Ala Arg Val Gly Asn Leu Phe Ser Ala Asp Leu Phe  
 145 150 155 160  
 Tyr Ser Pro Asp Gly Glu Met Phe Asp Val Met Glu Lys Tyr Gly Ile  
 165 170 175

<400> 25																
atg	acc	aga	aaa	aat	gtc	ctg	ctt	atc	gtc	gtt	gat	caa	tgg	cga	gca	48
Met	Thr	Arg	Lys	Asn	Val	Leu	Leu	Ile	Val	Val	Asp	Gln	Trp	Arg	Ala	
1				5				10				15				
gat	ttt	atc	cct	cac	ctg	atg	cgg	gcg	gag	ggg	cgc	gaa	cct	ttc	ctt	96
Asp	Phe	Ile	Pro	His	Leu	Met	Arg	Ala	Glu	Gly	Arg	Glu	Pro	Phe	Leu	
20				25				30								
aaa	act	ccc	aat	ctt	gat	cgt	ctt	tgc	cgg	gaa	ggc	ttg	acc	ttc	cgc	144
Lys	Thr	Pro	Asn	Leu	Asp	Arg	Leu	Cys	Arg	Glu	Gly	Leu	Thr	Phe	Arg	
35				40				45								
aat	cat	gtc	acg	acg	tgc	gtg	ccg	tgt	ggt	ccg	gca	agg	gca	agc	ctg	192
Asn	His	Val	Thr	Thr	Cys	Val	Pro	Cys	Gly	Pro	Ala	Arg	Ala	Ser	Leu	
50				55				60								
ctg	acg	ggc	ctc	tac	ctg	atg	aac	cac	cgg	gcg	gtg	cag	aac	act	gtt	240
Leu	Thr	Gly	Leu	Tyr	Leu	Met	Asn	His	Arg	Ala	Val	Gln	Asn	Thr	Val	
65				70				75				80				
ccg	ctt	gac	cag	cgc	cat	cta	aac	ctt	ggc	aag	gcc	ctg	cgc	gcc	att	288
Pro	Leu	Asp	Gln	Arg	His	Leu	Asn	Leu	Gly	Lys	Ala	Leu	Arg	Ala	Ile	
85				90				95								
ggc	tac	gat	ccc	gcg	ctc	att	ggt	tac	acc	acc	acg	aca	cct	gat	ccg	336
Gly	Tyr	Asp	Pro	Ala	Leu	Ile	Gly	Tyr	Thr	Thr	Thr	Thr	Pro	Asp	Pro	
100				105				110								
cgc	aca	acc	tct	gca	agg	gat	ccg	cgt	ttc	acg	gtc	ctg	ggc	gac	atc	384
Arg	Thr	Thr	Ser	Ala	Arg	Asp	Pro	Arg	Phe	Thr	Val	Leu	Gly	Asp	Ile	
115				120				125								
atg	gac	ggc	ttt	cgt	tcg	gtc	ggc	gca	ttc	gag	ccc	aat	atg	gag	ggg	432
Met	Asp	Gly	Phe	Arg	Ser	Val	Gly	Ala	Phe	Glu	Pro	Asn	Met	Glu	Gly	
130				135				140								
tat	ttt	ggc	tgg	gtg	gcg	cag	aac	ggc	ttc	gaa	ctg	cca	gag	aac	cgc	480
Tyr	Phe	Gly	Trp	Val	Ala	Gln	Asn	Gly	Phe	Glu	Leu	Pro	Glu	Asn	Arg	
145				150				155				160				

PF 53790

32

gaa gat atc tgg ctg ccg gaa ggt gaa cat tcc gtt ccc ggt gct acc	528
Glu Asp Ile Trp Leu Pro Glu Gly Glu His Ser Val Pro Gly Ala Thr	
165 170 175	
gac aaa ccg tgc cgc att ccg aag gaa ttt tgc gat tgc aca ttc ttc	576
Asp Lys Pro Ser Arg Ile Pro Lys Glu Phe Ser Asp Ser Thr Phe Phe	
180 185 190	
acg gag cgc gcc ctg aca tat ctg aag ggc agg gac ggc aag cct ttc	624
Thr Glu Arg Ala Leu Thr Tyr Leu Lys Gly Arg Asp Gly Lys Pro Phe	
195 200 205	
ttc ctg cat ctt ggc tat tat cgc ccg cat ccg cct ttc gta gcc tcc	672
Phe Leu His Leu Gly Tyr Tyr Arg Pro His Pro Pro Phe Val Ala Ser	
210 215 220	
gcg ccc tac cat gcg atg tac aaa gcc gaa gat atg cct gcg cct ata	720
Ala Pro Tyr His Ala Met Tyr Lys Ala Glu Asp Met Pro Ala Pro Ile	
225 230 235 240	
cgt gcg gag aat ccg gat gcc gaa gcg gca cag cat ccg ctc atg aag	768
Arg Ala Glu Asn Pro Asp Ala Glu Ala Ala Gln His Pro Leu Met Lys	
245 250 255	
cac tat atc gac cac atc aga cgc ggc tgc ttc ttc cat ggc gcg gaa	816
His Tyr Ile Asp His Ile Arg Arg Gly Ser Phe Phe His Gly Ala Glu	
260 265 270	
ggc tgc gga gca acg ctt gat gaa ggc gaa att cgc cag atg cgc gct	864
Gly Ser Gly Ala Thr Leu Asp Glu Gly Glu Ile Arg Gln Met Arg Ala	
275 280 285	
aca tat tgc gga ctg atc acc gag atc gac gat tgt ctg ggg agg gtc	912
Thr Tyr Cys Gly Leu Ile Thr Glu Ile Asp Asp Cys Leu Gly Arg Val	
290 295 300	
ttt gcc tat ctc gat gaa acc ggt cag tgg gac gac acg ctg att atc	960
Phe Ala Tyr Leu Asp Glu Thr Gly Gln Trp Asp Asp Thr Leu Ile Ile	
305 310 315 320	
ttc acg agc gat cat ggc gaa caa ctg ggc gat cat cac ctg ctc ggc	1008
Phe Thr Ser Asp His Gly Glu Gln Leu Gly Asp His His Leu Leu Gly	
325 330 335	
aag atc ggt tac aat gcc gaa agc ttc cgt att ccc ttg gtc ata aag	1056
Lys Ile Gly Tyr Asn Ala Glu Ser Phe Arg Ile Pro Leu Val Ile Lys	
340 345 350	
gat gcg gga cag aac ccg cac gcc ggc cag atc gaa gaa gcc ttc tcc	1104
Asp Ala Gly Gln Asn Arg His Ala Gly Gln Ile Glu Glu Gly Phe Ser	
355 360 365	
gaa agc atc gac gtc atg ccg acc atc ctc gaa tgg ctg ggc ggg gaa	1152
Glu Ser Ile Asp Val Met Pro Thr Ile Leu Glu Trp Leu Gly Gly Glu	
370 375 380	
acg cct cgc gcc tgc gac ggc cgt tgc ctg ttg ccg ttt ctg gct gag	1200
Thr Pro Arg Ala Cys Asp Gly Arg Ser Leu Leu Pro Phe Leu Ala Glu	
385 390 395 400	
gga aag ccc tcc gac tgg cgc acg gaa cta cat tac gag ttc gat ttt	1248
Gly Lys Pro Ser Asp Trp Arg Thr Glu Leu His Tyr Glu Phe Asp Phe	
405 410 415	
cgc gat gtc ttc tac gat cag ccg cag aac tgc gtc cag ctt tcc cag	1296
Arg Asp Val Phe Tyr Asp Gln Pro Gln Asn Ser Val Gln Leu Ser Gln	
420 425 430	



PF 53790

33

```

gat gat tgc agc ctc tgt gtg atc gag gac gaa aac tac aag tac gtg 1344
Asp Asp Cys Ser Leu Cys Val Ile Glu Asp Glu Asn Tyr Lys Tyr Val
      435                      440                      445

cat ttt gcc gcc ctg ccg ccg ctg ttc ttc gat ctg aag gca gac ccg 1392
His Phe Ala Ala Leu Pro Pro Leu Phe Phe Asp Leu Lys Ala Asp Pro
      450                      455                      460

cat gaa ttc agc aat ctg gct ggc gat cct gct tat gcg gcc ctc gtt 1440
His Glu Phe Ser Asn Leu Ala Gly Asp Pro Ala Tyr Ala Ala Leu Val
      465                      470                      475                      480

cgt gac tat gcc cag aag gca ttg tgc tgg cga ctg tct cat gcc gac 1488
Arg Asp Tyr Ala Gln Lys Ala Leu Ser Trp Arg Leu Ser His Ala Asp
      485                      490                      495

cgg aca ctc acc cat tac aga tcc agc ccg caa ggg ctg aca acg cgc 1536
Arg Thr Leu Thr His Tyr Arg Ser Ser Pro Gln Gly Leu Thr Thr Arg
      500                      505                      510

aac cat tga 1545
Asn His

<210> 26
<211> 514
<212> PRT
<213> Burkholderia caryophylli

<400> 26
Met Thr Arg Lys Asn Val Leu Leu Ile Val Val Asp Gln Trp Arg Ala
  1           5           10           15
Asp Phe Ile Pro His Leu Met Arg Ala Glu Gly Arg Glu Pro Phe Leu
      20           25           30
Lys Thr Pro Asn Leu Asp Arg Leu Cys Arg Glu Gly Leu Thr Phe Arg
      35           40           45
Asn His Val Thr Thr Cys Val Pro Cys Gly Pro Ala Arg Ala Ser Leu
      50           55           60
Leu Thr Gly Leu Tyr Leu Met Asn His Arg Ala Val Gln Asn Thr Val
      65           70           75           80
Pro Leu Asp Gln Arg His Leu Asn Leu Gly Lys Ala Leu Arg Ala Ile
      85           90           95
Gly Tyr Asp Pro Ala Leu Ile Gly Tyr Thr Thr Thr Thr Pro Asp Pro
      100          105          110
Arg Thr Thr Ser Ala Arg Asp Pro Arg Phe Thr Val Leu Gly Asp Ile
      115          120          125
Met Asp Gly Phe Arg Ser Val Gly Ala Phe Glu Pro Asn Met Glu Gly
      130          135          140
Tyr Phe Gly Trp Val Ala Gln Asn Gly Phe Glu Leu Pro Glu Asn Arg
      145          150          155          160
Glu Asp Ile Trp Leu Pro Glu Gly Glu His Ser Val Pro Gly Ala Thr
      165          170          175
Asp Lys Pro Ser Arg Ile Pro Lys Glu Phe Ser Asp Ser Thr Phe Phe
      180          185          190
Thr Glu Arg Ala Leu Thr Tyr Leu Lys Gly Arg Asp Gly Lys Pro Phe
      195          200          205

```

PF 53790

34

Phe Leu His Leu Gly Tyr Tyr Arg Pro His Pro Pro Phe Val Ala Ser  
 210 215 220  
 Ala Pro Tyr His Ala Met Tyr Lys Ala Glu Asp Met Pro Ala Pro Ile  
 225 230 235 240  
 Arg Ala Glu Asn Pro Asp Ala Glu Ala Ala Gln His Pro Leu Met Lys  
 245 250 255  
 His Tyr Ile Asp His Ile Arg Arg Gly Ser Phe Phe His Gly Ala Glu  
 260 265 270  
 Gly Ser Gly Ala Thr Leu Asp Glu Gly Glu Ile Arg Gln Met Arg Ala  
 275 280 285  
 Thr Tyr Cys Gly Leu Ile Thr Glu Ile Asp Asp Cys Leu Gly Arg Val  
 290 295 300  
 Phe Ala Tyr Leu Asp Glu Thr Gly Gln Trp Asp Asp Thr Leu Ile Ile  
 305 310 315 320  
 Phe Thr Ser Asp His Gly Glu Gln Leu Gly Asp His His Leu Leu Gly  
 325 330 335  
 Lys Ile Gly Tyr Asn Ala Glu Ser Phe Arg Ile Pro Leu Val Ile Lys  
 340 345 350  
 Asp Ala Gly Gln Asn Arg His Ala Gly Gln Ile Glu Glu Gly Phe Ser  
 355 360 365  
 Glu Ser Ile Asp Val Met Pro Thr Ile Leu Glu Trp Leu Gly Gly Glu  
 370 375 380  
 Thr Pro Arg Ala Cys Asp Gly Arg Ser Leu Leu Pro Phe Leu Ala Glu  
 385 390 395 400  
 Gly Lys Pro Ser Asp Trp Arg Thr Glu Leu His Tyr Glu Phe Asp Phe  
 405 410 415  
 Arg Asp Val Phe Tyr Asp Gln Pro Gln Asn Ser Val Gln Leu Ser Gln  
 420 425 430  
 Asp Asp Cys Ser Leu Cys Val Ile Glu Asp Glu Asn Tyr Lys Tyr Val  
 435 440 445  
 His Phe Ala Ala Leu Pro Pro Leu Phe Phe Asp Leu Lys Ala Asp Pro  
 450 455 460  
 His Glu Phe Ser Asn Leu Ala Gly Asp Pro Ala Tyr Ala Ala Leu Val  
 465 470 475 480  
 Arg Asp Tyr Ala Gln Lys Ala Leu Ser Trp Arg Leu Ser His Ala Asp  
 485 490 495  
 Arg Thr Leu Thr His Tyr Arg Ser Ser Pro Gln Gly Leu Thr Thr Arg  
 500 505 510  
 Asn His

&lt;210&gt; 27

&lt;211&gt; 2250

&lt;212&gt; DNA

&lt;213&gt; Agrobacterium rhizogenes

&lt;220&gt;

&lt;221&gt; CDS

PF 53790

35

&lt;222&gt; (1)..(2247)

&lt;223&gt; coding for tryptophan oxygenase (aux1)

&lt;400&gt; 27

atg gct gga tcc tcc ttc aca ttg cca tca act ggc tca gcg ccc ctt	48
Met Ala Gly Ser Ser Phe Thr Leu Pro Ser Thr Gly Ser Ala Pro Leu	
1 5 10 15	
gat atg atg ctt atc gat gat tca gat ctg ctg caa ttg ggt ctc cag	96
Asp Met Met Leu Ile Asp Asp Ser Asp Leu Leu Gln Leu Gly Leu Gln	
20 25 30	
cag gta ttc tcg aag cgg tac aca gag aca ccg cag tca cgc tac aaa	144
Gln Val Phe Ser Lys Arg Tyr Thr Glu Thr Pro Gln Ser Arg Tyr Lys	
35 40 45	
ctg acc agg agg gct tct cca gac gtc tca tct ggc gaa ggc aat gtg	192
Leu Thr Arg Arg Ala Ser Pro Asp Val Ser Ser Gly Glu Gly Asn Val	
50 55 60	
cat gcc ctt gcg ttc ata tat gtc aac gct gag acg ttg cag atg atc	240
His Ala Leu Ala Phe Ile Tyr Val Asn Ala Glu Thr Leu Gln Met Ile	
65 70 75 80	
aaa aac gct cga tcg cta acc gaa gcg aac ggc gtc aaa gat ctt gtc	288
Lys Asn Ala Arg Ser Leu Thr Glu Ala Asn Gly Val Lys Asp Leu Val	
85 90 95	
gcc atc gac gtt ccg cca ttt cga aac gac ttc tca aga gcg cta ctc	336
Ala Ile Asp Val Pro Pro Phe Arg Asn Asp Phe Ser Arg Ala Leu Leu	
100 105 110	
ctt caa gtg atc aac ttg ttg gga aac aac cga aat gcc gat gac gat	384
Leu Gln Val Ile Asn Leu Leu Gly Asn Asn Arg Asn Ala Asp Asp Asp	
115 120 125	
ctt agt cac ttc ata gca gtt gct ctc cca aac agc gcc cgc tct aag	432
Leu Ser His Phe Ile Ala Val Ala Leu Pro Asn Ser Ala Arg Ser Lys	
130 135 140	
atc cta acc acg gca ccg ttc gaa gga agc ttg tca gaa aac ttc agg	480
Ile Leu Thr Thr Ala Pro Phe Glu Gly Ser Leu Ser Glu Asn Phe Arg	
145 150 155 160	
ggg ttc ccg atc act cgt gaa gga aat gtg gca tgt gaa gtg cta gcc	528
Gly Phe Pro Ile Thr Arg Glu Gly Asn Val Ala Cys Glu Val Leu Ala	
165 170 175	
tat ggg aat aac ttg atg ccc aag gcc tgc tcc gat tcc ttt cca acc	576
Tyr Gly Asn Asn Leu Met Pro Lys Ala Cys Ser Asp Ser Phe Pro Thr	
180 185 190	
gtg gat ctt ctt tat gac tat ggc aag ttc ttc gag agt tgc gcg gcc	624
Val Asp Leu Leu Tyr Asp Tyr Gly Lys Phe Phe Glu Ser Cys Ala Ala	
195 200 205	
gat gga cgt atc ggt tat ttt cct gaa ggc gtt acg aaa cct aaa gtg	672
Asp Gly Arg Ile Gly Tyr Phe Pro Glu Gly Val Thr Lys Pro Lys Val	
210 215 220	
gct ata att ggc gca ggc ttt tcc ggg ctc gtt gca gcg agc gaa cta	720
Ala Ile Ile Gly Ala Gly Phe Ser Gly Leu Val Ala Ala Ser Glu Leu	
225 230 235 240	
ctt cat gca ggg gta gac gat gtt acg gtg tat gag gcg agt gat cgg	768
Leu His Ala Gly Val Asp Asp Val Thr Val Tyr Glu Ala Ser Asp Arg	
245 250 255	

PF 53790

36

ctt gga gga aag cta tgg tca cac gga ttt aag agt gct cca aat gtg	816
Leu Gly Gly Lys Leu Trp Ser His Gly Phe Lys Ser Ala Pro Asn Val	
260 265 270	
ata gcc gag atg ggg gcc atg cgt ttt ccg cga agt gaa tca tgc ttg	864
Ile Ala Glu Met Gly Ala Met Arg Phe Pro Arg Ser Glu Ser Cys Leu	
275 280 285	
ttc ttc tat ctc aaa aag cac gga ctg gac tcc gtt ggt ctg ttc ccg	912
Phe Phe Tyr Leu Lys Lys His Gly Leu Asp Ser Val Gly Leu Phe Pro	
290 295 300	
aat ccg gga agt gtc gat acc gca ttg ttc tac agg ggc cgt caa tat	960
Asn Pro Gly Ser Val Thr Ala Leu Phe Tyr Arg Gly Arg Gln Tyr	
305 310 315 320	
atc tgg aaa gcg gga gag gag cca ccg gag ctg ttt cgt cgt gtg cac	1008
Ile Trp Lys Ala Gly Glu Glu Pro Pro Glu Leu Phe Arg Arg Val His	
325 330 335	
cat gga tgg cgc gca ttt ttg caa gat ggc tat ctc cat gat gga gtc	1056
His Gly Trp Arg Ala Phe Leu Gln Asp Gly Tyr Leu His Asp Gly Val	
340 345 350	
atg ttg gcg tca ccg tta gca att gtt gac gcc ttg aat tta ggg cat	1104
Met Leu Ala Ser Pro Leu Ala Ile Val Asp Ala Leu Asn Leu Gly His	
355 360 365	
cta cag cag gcg cat ggc ttc tgg caa tct tgg ctc aca tat ttt gag	1152
Leu Gln Gln Ala His Gly Phe Trp Gln Ser Trp Leu Thr Tyr Phe Glu	
370 375 380	
cga gag tct ttc tct tct ggc atc gaa aaa atg ttc ttg ggc aat cat	1200
Arg Glu Ser Phe Ser Ser Gly Ile Glu Lys Met Phe Leu Gly Asn His	
385 390 395 400	
cct ccg ggg ggt gaa caa tgg aat tcc cta gat gac ttg gat ctt ttc	1248
Pro Pro Gly Gly Glu Gln Trp Asn Ser Leu Asp Asp Leu Asp Leu Phe	
405 410 415	
aaa gcg ctg ggt att gga tcc ggc gga ttc ggc cct gta ttt gaa agt	1296
Lys Ala Leu Gly Ile Gly Ser Gly Gly Phe Gly Pro Val Phe Glu Ser	
420 425 430	
ggg ttt atc gag atc ctt cgc tta gtc gtc aac ggg tat gag gat aac	1344
Gly Phe Ile Glu Ile Leu Arg Leu Val Val Asn Gly Tyr Glu Asp Asn	
435 440 445	
gtg cgg ctg agt tac gaa gga att tct gag ctg cct cat agg atc gcc	1392
Val Arg Leu Ser Tyr Glu Gly Ile Ser Glu Leu Pro His Arg Ile Ala	
450 455 460	
tca cag gta att aac ggc aga tct att cgc gag cgt aca att cac gtt	1440
Ser Gln Val Ile Asn Gly Arg Ser Ile Arg Glu Arg Thr Ile His Val	
465 470 475 480	
caa gtc gag cag att gat aga gag gag gat aaa ata aat atc aag atc	1488
Gln Val Glu Gln Ile Asp Arg Glu Glu Asp Lys Ile Asn Ile Lys Ile	
485 490 495	
aaa gga gga aag gtt gag gtc tat gat cga gta ctg gtt aca tcc ggg	1536
Lys Gly Gly Lys Val Glu Val Tyr Asp Arg Val Leu Val Thr Ser Gly	
500 505 510	
ttt gcg aac atc gaa atg cgc cat ctc ctg aca tca agc aac gca ttc	1584
Phe Ala Asn Ile Glu Met Arg His Leu Leu Thr Ser Ser Asn Ala Phe	
515 520 525	

PF 53790

37

ttc cat gca gat gta agc cat gca ata ggg aac agt cat atg act ggt	1632
Phe His Ala Asp Val Ser His Ala Ile Gly Asn Ser His Met Thr Gly	
530 535 540	
gcg tca aaa ctg ttc ttg ctg act aac gaa aaa ttc tgg cta caa cat	1680
Ala Ser Lys Leu Phe Leu Leu Thr Asn Glu Lys Phe Trp Leu Gln His	
545 550 555 560	
cat ttg cca tcg tgc ata ctc acc acc ggc gtt gca aag gca gtt tat	1728
His Leu Pro Ser Cys Ile Leu Thr Thr Gly Val Ala Lys Ala Val Tyr	
565 570 575	
tgc tta gac tat gat ccg cga gat cca agc ggc aaa gga ctg gtg ttg	1776
Cys Leu Asp Tyr Asp Pro Arg Asp Pro Ser Gly Lys Gly Leu Val Leu	
580 585 590	
ata agc tat act tgg gag gat gac tca cat aag ctc cta gcc gtc ccc	1824
Ile Ser Tyr Thr Trp Glu Asp Ser His Lys Leu Leu Ala Val Pro	
595 600 605	
gac aaa aga gaa agg ttc gca tcg ctg cag cgc gat att ggg agg gca	1872
Asp Lys Arg Glu Arg Phe Ala Ser Leu Gln Arg Asp Ile Gly Arg Ala	
610 615 620	
ttc cca gat ttt gcc aag cac cta act cct gca gac ggg aac tat gat	1920
Phe Pro Asp Phe Ala Lys His Leu Thr Pro Ala Asp Gly Asn Tyr Asp	
625 630 635 640	
gat aat atc gtt caa cat gat tgg ctg act gat ccc cac gct ggc gga	1968
Asp Asn Ile Val Gln His Asp Trp Leu Thr Asp Pro His Ala Gly Gly	
645 650 655	
gcg ttt aaa ctg aac cgc aga ggc aac gac gta tat tca gaa agg ctt	2016
Ala Phe Lys Leu Asn Arg Arg Gly Asn Asp Val Tyr Ser Glu Arg Leu	
660 665 670	
ttc ttt cag ccc ttt gac gta atg cat ccc gcg gac gat aag gga ctt	2064
Phe Phe Gln Pro Phe Asp Val Met His Pro Ala Asp Asp Lys Gly Leu	
675 680 685	
tac ttg gcc ggt tgt agc tgt tcc ttc acc gga ggg tgg gtt cat ggt	2112
Tyr Leu Ala Gly Cys Ser Cys Ser Phe Thr Gly Trp Val His Gly	
690 695 700	
gcc att cag acc gca tgc aac gct acg tgt gcg atc att tat ggt tcc	2160
Ala Ile Gln Thr Ala Cys Asn Ala Thr Cys Ala Ile Ile Tyr Gly Ser	
705 710 715 720	
gga cac ctg caa gag cta atc cac tgg cga cac ctc aaa gaa ggt aat	2208
Gly His Leu Gln Glu Leu Ile His Trp Arg His Leu Lys Glu Gly Asn	
725 730 735	
cca ctg gcg cac gct tgg aag cgg tat agg tat caa gcg tga	2250
Pro Leu Ala His Ala Trp Lys Arg Tyr Arg Tyr Gln Ala	
740 745	

&lt;210&gt; 28

&lt;211&gt; 749

&lt;212&gt; PRT

&lt;213&gt; Agrobacterium rhizogenes

&lt;400&gt; 28

Met Ala Gly Ser Ser Phe Thr Leu Pro Ser Thr Gly Ser Ala Pro Leu
1 5 10 15

Asp	Met	Met	Leu 20	Ile	Asp	Asp	Ser	Asp 25	Leu	Leu	Gln	Leu	Gly 30	Leu	Gln
Gln	Val	Phe 35	Ser	Lys	Arg	Tyr	Thr 40	Glu	Thr	Pro	Gln	Ser 45	Arg	Tyr	Lys
Leu	Thr 50	Arg	Arg	Ala	Ser	Pro 55	Asp	Val	Ser	Ser	Gly 60	Glu	Gly	Asn	Val
His 65	Ala	Leu	Ala	Phe	Ile 70	Tyr	Val	Asn	Ala	Glu 75	Thr	Leu	Gln	Met	Ile 80
Lys	Asn	Ala	Arg	Ser 85	Leu	Thr	Glu	Ala	Asn 90	Gly	Val	Lys	Asp	Leu 95	Val
Ala	Ile	Asp	Val 100	Pro	Pro	Phe	Arg	Asn 105	Asp	Phe	Ser	Arg	Ala 110	Leu	Leu
Leu	Gln	Val 115	Ile	Asn	Leu	Leu	Gly 120	Asn	Asn	Arg	Asn	Ala 125	Asp	Asp	Asp
Leu	Ser 130	His	Phe	Ile	Ala	Val 135	Ala	Leu	Pro	Asn	Ser 140	Ala	Arg	Ser	Lys
Ile 145	Leu	Thr	Thr	Ala	Pro 150	Phe	Glu	Gly	Ser	Leu 155	Ser	Glu	Asn	Phe	Arg 160
Gly	Phe	Pro	Ile	Thr 165	Arg	Glu	Gly	Asn 170	Val	Ala	Cys	Glu	Val	Leu 175	Ala
Tyr	Gly	Asn	Asn 180	Leu	Met	Pro	Lys	Ala 185	Cys	Ser	Asp	Ser	Phe 190	Pro	Thr
Val	Asp	Leu 195	Leu	Tyr	Asp	Tyr	Gly 200	Lys	Phe	Phe	Glu	Ser 205	Cys	Ala	Ala
Asp	Gly 210	Arg	Ile	Gly	Tyr	Phe 215	Pro	Glu	Gly	Val	Thr 220	Lys	Pro	Lys	Val
Ala 225	Ile	Ile	Gly	Ala	Gly 230	Phe	Ser	Gly	Leu	Val 235	Ala	Ala	Ser	Glu	Leu 240
Leu	His	Ala	Gly	Val 245	Asp	Asp	Val	Thr	Val 250	Tyr	Glu	Ala	Ser	Asp 255	Arg
Leu	Gly	Gly	Lys 260	Leu	Trp	Ser	His	Gly 265	Phe	Lys	Ser	Ala	Pro 270	Asn	Val
Ile	Ala	Glu 275	Met	Gly	Ala	Met	Arg 280	Phe	Pro	Arg	Ser	Glu 285	Ser	Cys	Leu
Phe	Phe 290	Tyr	Leu	Lys	Lys	His 295	Gly	Leu	Asp	Ser	Val 300	Gly	Leu	Phe	Pro
Asn 305	Pro	Gly	Ser	Val	Asp 310	Thr	Ala	Leu	Phe	Tyr 315	Arg	Gly	Arg	Gln	Tyr 320
Ile	Trp	Lys	Ala	Gly 325	Glu	Glu	Pro	Pro	Glu 330	Leu	Phe	Arg	Arg	Val 335	His
His	Gly	Trp	Arg 340	Ala	Phe	Leu	Gln	Asp 345	Gly	Tyr	Leu	His	Asp 350	Gly	Val
Met	Leu	Ala 355	Ser	Pro	Leu	Ala	Ile 360	Val	Asp	Ala	Leu	Asn 365	Leu	Gly	His
Leu	Gln 370	Gln	Ala	His	Gly	Phe 375	Trp	Gln	Ser	Trp	Leu 380	Thr	Tyr	Phe	Glu

PF 53790

39

Arg	Glu	Ser	Phe	Ser	Ser	Gly	Ile	Glu	Lys	Met	Phe	Leu	Gly	Asn	His
385					390					395					400
Pro	Pro	Gly	Gly	Glu	Gln	Trp	Asn	Ser	Leu	Asp	Asp	Leu	Asp	Leu	Phe
				405					410					415	
Lys	Ala	Leu	Gly	Ile	Gly	Ser	Gly	Gly	Phe	Gly	Pro	Val	Phe	Glu	Ser
			420					425					430		
Gly	Phe	Ile	Glu	Ile	Leu	Arg	Leu	Val	Val	Asn	Gly	Tyr	Glu	Asp	Asn
			435				440					445			
Val	Arg	Leu	Ser	Tyr	Glu	Gly	Ile	Ser	Glu	Leu	Pro	His	Arg	Ile	Ala
			450				455				460				
Ser	Gln	Val	Ile	Asn	Gly	Arg	Ser	Ile	Arg	Glu	Arg	Thr	Ile	His	Val
465					470					475					480
Gln	Val	Glu	Gln	Ile	Asp	Arg	Glu	Glu	Asp	Lys	Ile	Asn	Ile	Lys	Ile
				485					490					495	
Lys	Gly	Gly	Lys	Val	Glu	Val	Tyr	Asp	Arg	Val	Leu	Val	Thr	Ser	Gly
			500					505					510		
Phe	Ala	Asn	Ile	Glu	Met	Arg	His	Leu	Leu	Thr	Ser	Ser	Asn	Ala	Phe
			515				520					525			
Phe	His	Ala	Asp	Val	Ser	His	Ala	Ile	Gly	Asn	Ser	His	Met	Thr	Gly
			530				535				540				
Ala	Ser	Lys	Leu	Phe	Leu	Leu	Thr	Asn	Glu	Lys	Phe	Trp	Leu	Gln	His
545					550					555					560
His	Leu	Pro	Ser	Cys	Ile	Leu	Thr	Thr	Gly	Val	Ala	Lys	Ala	Val	Tyr
				565					570					575	
Cys	Leu	Asp	Tyr	Asp	Pro	Arg	Asp	Pro	Ser	Gly	Lys	Gly	Leu	Val	Leu
			580					585					590		
Ile	Ser	Tyr	Thr	Trp	Glu	Asp	Asp	Ser	His	Lys	Leu	Leu	Ala	Val	Pro
			595				600					605			
Asp	Lys	Arg	Glu	Arg	Phe	Ala	Ser	Leu	Gln	Arg	Asp	Ile	Gly	Arg	Ala
			610			615					620				
Phe	Pro	Asp	Phe	Ala	Lys	His	Leu	Thr	Pro	Ala	Asp	Gly	Asn	Tyr	Asp
625					630					635					640
Asp	Asn	Ile	Val	Gln	His	Asp	Trp	Leu	Thr	Asp	Pro	His	Ala	Gly	Gly
				645					650					655	
Ala	Phe	Lys	Leu	Asn	Arg	Arg	Gly	Asn	Asp	Val	Tyr	Ser	Glu	Arg	Leu
			660					665					670		
Phe	Phe	Gln	Pro	Phe	Asp	Val	Met	His	Pro	Ala	Asp	Asp	Lys	Gly	Leu
			675				680					685			
Tyr	Leu	Ala	Gly	Cys	Ser	Cys	Ser	Phe	Thr	Gly	Gly	Trp	Val	His	Gly
	690					695					700				
Ala	Ile	Gln	Thr	Ala	Cys	Asn	Ala	Thr	Cys	Ala	Ile	Ile	Tyr	Gly	Ser
705					710					715					720
Gly	His	Leu	Gln	Glu	Leu	Ile	His	Trp	Arg	His	Leu	Lys	Glu	Gly	Asn
				725					730					735	
Pro	Leu	Ala	His	Ala	Trp	Lys	Arg	Tyr	Arg	Tyr	Gln	Ala			
			740					745							

PF 53790

40

```

<210> 29
<211> 1401
<212> DNA
<213> Agrobacterium rhizogenes
<220>
<221> CDS
<222> (1)..(1398)
<223> coding for indoleacetamide hydrolase
<400> 29
atg gtg acc ctc tcc tcg atc acc gag acg ctt aaa tgt ctc agg gaa 48
Met Val Thr Leu Ser Ser Ile Thr Glu Thr Leu Lys Cys Leu Arg Glu
1 5 10 15
aga aaa tac tcg tgc ttt gag tta atc gaa acg ata ata gcc cgc tgt 96
Arg Lys Tyr Ser Cys Phe Glu Leu Ile Glu Thr Ile Ile Ala Arg Cys
20 25 30
gaa gca gca aga tcc tta aac gcc ttt ctg gaa acc gac tgg gcg cac 144
Glu Ala Ala Arg Ser Leu Asn Ala Phe Leu Glu Thr Asp Trp Ala His
35 40 45
cta cgg tgg act gcc agc aaa atc gat caa cac gga ggt gcc ggt gtt 192
Leu Arg Trp Thr Ala Ser Lys Ile Asp Gln His Gly Gly Ala Gly Val
50 55 60
ggc cta gct ggc gtt ccc cta tgc ttt aaa gcg aat att gcg aca ggc 240
Gly Leu Ala Gly Val Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly
65 70 75 80
agg ttc gcc gcg acc gct ggt acg cca ggc tta cag aac cac aaa ccc 288
Arg Phe Ala Ala Thr Ala Gly Thr Pro Gly Leu Gln Asn His Lys Pro
85 90 95
aag acg cct gcc gga gtt gca cga caa ctt ctc gcg gct ggg gca ctg 336
Lys Thr Pro Ala Gly Val Ala Arg Gln Leu Leu Ala Ala Gly Ala Leu
100 105 110
cct ggc gct tcg gga aac atg cac gaa ttg tct ttt ggg atc acg agc 384
Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser
115 120 125
aac aac ttc gcc aca ggc gcc gta cga aac ccg tgg aac cct agt ctc 432
Asn Asn Phe Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu
130 135 140
atc cca ggg gga tca agt ggg ggt gtg gcc gcc gcg gtg gcc ggc cga 480
Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Ala Val Ala Gly Arg
145 150 155 160
ttg atg ctg ggc ggc gtc gga act gac acg gga gcg tcg gtc cgt tta 528
Leu Met Leu Gly Gly Val Gly Thr Asp Thr Gly Ala Ser Val Arg Leu
165 170 175
ccg gcc gcc ttg tgc ggc gtg gtg ggg ttt cgt cct acc gtg ggg cga 576
Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Val Gly Arg
180 185 190
tat cca acg gac gga ata gtt ccg gta agc ccc acc cgg gac acc cct 624
Tyr Pro Thr Asp Gly Ile Val Pro Val Ser Pro Thr Arg Asp Thr Pro
195 200 205

```



PF 53790

41

ggc gtt atc gca cag aat gtt ccg gac gtg att ctt ctt gac ggt atc Gly Val Ile Ala Gln Asn Val Pro Asp Val Ile Leu Leu Asp Gly Ile 210 215 220	672
att tgc ggg aga ccg ccg gtt aat caa acg gtc cgc ctg aag ggg ctg Ile Cys Gly Arg Pro Pro Val Asn Gln Thr Val Arg Leu Lys Gly Leu 225 230 235 240	720
cgt ata ggc ttg cca acc gct tac ttt tac aac gac ctg gag ccc gat Arg Ile Gly Leu Pro Thr Ala Tyr Phe Tyr Asn Asp Leu Glu Pro Asp 245 250 255	768
gtc gcc tta gca gcc gag acg att atc aga gtt ctg gca cgc aaa gat Val Ala Leu Ala Ala Glu Thr Ile Ile Arg Val Leu Ala Arg Lys Asp 260 265 270	816
gtt act ttt gtt gaa gca gat att cct gat tta gcg cat cac aat gaa Val Thr Phe Val Glu Ala Asp Ile Pro Asp Leu Ala His His Asn Glu 275 280 285	864
ggg gtc agc ttt ccg act gcc atc tac gaa ttt ccg ttg tcc ctt gaa Gly Val Ser Phe Pro Thr Ala Ile Tyr Glu Phe Pro Leu Ser Leu Glu 290 295 300	912
cat tat att cag aac ttc gta gag ggt gtt tcc ttt tct gag gtt gtc His Tyr Ile Gln Asn Phe Val Glu Gly Val Ser Phe Ser Glu Val Val 305 310 315 320	960
aga gcg att cgc agt ccg gat gtt gca agt att ctc aat gca caa ctc Arg Ala Ile Arg Ser Pro Asp Val Ala Ser Ile Leu Asn Ala Gln Leu 325 330 335	1008
tcg gat aat ctt att tcc aaa agc gag tat tgt ctg gcg cga cgt ttt Ser Asp Asn Leu Ile Ser Lys Ser Glu Tyr Cys Leu Ala Arg Arg Phe 340 345 350	1056
ttc aga ccg aga ctc caa gcg gcc tac cac agt tac ttc aag gcg cat Phe Arg Pro Arg Leu Gln Ala Ala Tyr His Ser Tyr Phe Lys Ala His 355 360 365	1104
cag cta gat gca att ctt ttc cca aca gct ccg ttg aca gcc aag cca Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Thr Ala Lys Pro 370 375 380	1152
att ggc cat gat cta tcg gtg att cac aat ggc tca atg acc gat acc Ile Gly His Asp Leu Ser Val Ile His Asn Gly Ser Met Thr Asp Thr 385 390 395 400	1200
ttt aaa atc ttc gtg cgg aat gta gat ccc agc agt aat gcg ggc ctg Phe Lys Ile Phe Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu 405 410 415	1248
ccg ggc cta agt ctt ccc gtt tct ctt agt tcc aac ggt ctg cct att Pro Gly Leu Ser Leu Pro Val Ser Leu Ser Ser Asn Gly Leu Pro Ile 420 425 430	1296
ggc atg gaa atc gat ggc tct gca agc tcg gat gaa cgt ctg tta gca Gly Met Glu Ile Asp Gly Ser Ala Ser Ser Asp Glu Arg Leu Leu Ala 435 440 445	1344
att gga cta gcg ata gaa gaa gca ata gac ttt agg cat cgt ccg act Ile Gly Leu Ala Ile Glu Glu Ala Ile Asp Phe Arg His Arg Pro Thr 450 455 460	1392
ctg tcg taa Leu Ser 465	1401

PF 53790

42

&lt;210&gt; 30

&lt;211&gt; 466

&lt;212&gt; PRT

&lt;213&gt; Agrobacterium rhizogenes

&lt;400&gt; 30

Met	Val	Thr	Leu	Ser	Ser	Ile	Thr	Glu	Thr	Leu	Lys	Cys	Leu	Arg	Glu
1				5					10					15	
Arg	Lys	Tyr	Ser	Cys	Phe	Glu	Leu	Ile	Glu	Thr	Ile	Ile	Ala	Arg	Cys
			20					25					30		
Glu	Ala	Ala	Arg	Ser	Leu	Asn	Ala	Phe	Leu	Glu	Thr	Asp	Trp	Ala	His
			35				40					45			
Leu	Arg	Trp	Thr	Ala	Ser	Lys	Ile	Asp	Gln	His	Gly	Gly	Ala	Gly	Val
	50					55					60				
Gly	Leu	Ala	Gly	Val	Pro	Leu	Cys	Phe	Lys	Ala	Asn	Ile	Ala	Thr	Gly
	65				70					75					80
Arg	Phe	Ala	Ala	Thr	Ala	Gly	Thr	Pro	Gly	Leu	Gln	Asn	His	Lys	Pro
				85					90					95	
Lys	Thr	Pro	Ala	Gly	Val	Ala	Arg	Gln	Leu	Leu	Ala	Ala	Gly	Ala	Leu
			100					105					110		
Pro	Gly	Ala	Ser	Gly	Asn	Met	His	Glu	Leu	Ser	Phe	Gly	Ile	Thr	Ser
			115				120					125			
Asn	Asn	Phe	Ala	Thr	Gly	Ala	Val	Arg	Asn	Pro	Trp	Asn	Pro	Ser	Leu
	130					135					140				
Ile	Pro	Gly	Gly	Ser	Ser	Gly	Gly	Val	Ala	Ala	Val	Ala	Gly	Arg	
	145				150					155				160	
Leu	Met	Leu	Gly	Gly	Val	Gly	Thr	Asp	Thr	Gly	Ala	Ser	Val	Arg	Leu
			165					170						175	
Pro	Ala	Ala	Leu	Cys	Gly	Val	Val	Gly	Phe	Arg	Pro	Thr	Val	Gly	Arg
			180					185					190		
Tyr	Pro	Thr	Asp	Gly	Ile	Val	Pro	Val	Ser	Pro	Thr	Arg	Asp	Thr	Pro
			195				200					205			
Gly	Val	Ile	Ala	Gln	Asn	Val	Pro	Asp	Val	Ile	Leu	Leu	Asp	Gly	Ile
	210					215					220				
Ile	Cys	Gly	Arg	Pro	Pro	Val	Asn	Gln	Thr	Val	Arg	Leu	Lys	Gly	Leu
	225				230					235				240	
Arg	Ile	Gly	Leu	Pro	Thr	Ala	Tyr	Phe	Tyr	Asn	Asp	Leu	Glu	Pro	Asp
			245					250						255	
Val	Ala	Leu	Ala	Ala	Glu	Thr	Ile	Ile	Arg	Val	Leu	Ala	Arg	Lys	Asp
			260					265					270		
Val	Thr	Phe	Val	Glu	Ala	Asp	Ile	Pro	Asp	Leu	Ala	His	His	Asn	Glu
			275				280					285			
Gly	Val	Ser	Phe	Pro	Thr	Ala	Ile	Tyr	Glu	Phe	Pro	Leu	Ser	Leu	Glu
	290					295				300					
His	Tyr	Ile	Gln	Asn	Phe	Val	Glu	Gly	Val	Ser	Phe	Ser	Glu	Val	Val
	305				310					315				320	
Arg	Ala	Ile	Arg	Ser	Pro	Asp	Val	Ala	Ser	Ile	Leu	Asn	Ala	Gln	Leu
			325						330					335	

PF 53790

43

Ser Asp Asn Leu Ile Ser Lys Ser Glu Tyr Cys Leu Ala Arg Arg Phe  
 340 345 350  
 Phe Arg Pro Arg Leu Gln Ala Ala Tyr His Ser Tyr Phe Lys Ala His  
 355 360 365  
 Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Thr Ala Lys Pro  
 370 375 380  
 Ile Gly His Asp Leu Ser Val Ile His Asn Gly Ser Met Thr Asp Thr  
 385 390 395 400  
 Phe Lys Ile Phe Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu  
 405 410 415  
 Pro Gly Leu Ser Leu Pro Val Ser Leu Ser Ser Asn Gly Leu Pro Ile  
 420 425 430  
 Gly Met Glu Ile Asp Gly Ser Ala Ser Ser Asp Glu Arg Leu Leu Ala  
 435 440 445  
 Ile Gly Leu Ala Ile Glu Glu Ala Ile Asp Phe Arg His Arg Pro Thr  
 450 455 460  
 Leu Ser  
 465

&lt;210&gt; 31

&lt;211&gt; 2268

&lt;212&gt; DNA

&lt;213&gt; Agrobacterium tumefaciens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(2265)

&lt;223&gt; coding for tryptophan monooxygenase

&lt;400&gt; 31

atg tca gct tca cct ctc ctt gat aac cag tgc gat cat ttc tct acc 48  
 Met Ser Ala Ser Pro Leu Leu Asp Asn Gln Cys Asp His Phe Ser Thr  
 1 5 10 15  
 aaa atg gtg gat ctg ata atg gtc gat aag gct gat gaa ttg gac cgc 96  
 Lys Met Val Asp Leu Ile Met Val Asp Lys Ala Asp Glu Leu Asp Arg  
 20 25 30  
 agg gtt tcc gat gcc ttc tca gaa cgt gaa gct tct agg gga agg agg 144  
 Arg Val Ser Asp Ala Phe Ser Glu Arg Glu Ala Ser Arg Gly Arg Arg  
 35 40 45  
 att act caa atc tcc ggc gag tgc agc gct ggg tta gct tgc aaa agg 192  
 Ile Thr Gln Ile Ser Gly Glu Cys Ser Ala Gly Leu Ala Cys Lys Arg  
 50 55 60  
 ctg gcc gac ggt cgc ttt ccc gag atc tca act ggt gag aag gta gca 240  
 Leu Ala Asp Gly Arg Phe Pro Glu Ile Ser Thr Gly Glu Lys Val Ala  
 65 70 75 80  
 gcc ctc tcc gct tac atc tat gtt ggc aag gaa att ctg ggg cgg ata 288  
 Ala Leu Ser Ala Tyr Ile Tyr Val Gly Lys Glu Ile Leu Gly Arg Ile  
 85 90 95  
 ctt gaa tcg gaa cct tgg gcg cga gca aga gtg agt ggt ctc gtt gcc 336  
 Leu Glu Ser Glu Pro Trp Ala Arg Ala Arg Val Ser Gly Leu Val Ala  
 100 105 110

PF 53790

44

atc gac ctt gca cca ttt tgt atg gat ttc tcc gaa gca caa ctt ctc Ile Asp Leu Ala Pro Phe Cys Met Asp Phe Ser Glu Ala Gln Leu Leu 115 120 125	384
caa acc ctg ttt ttg ctg agc ggt aaa aga tgt gca tcc agc gat ctt Gln Thr Leu Phe Leu Leu Ser Gly Lys Arg Cys Ala Ser Ser Asp Leu 130 135 140	432
agt cat ttc gtg gcc att tca atc tct aag act gcc cgc tcc cga acc Ser His Phe Val Ala Ile Ser Ile Ser Lys Thr Ala Arg Ser Arg Thr 145 150 155 160	480
ctg caa atg ccg ccg tac gag aaa ggc acg acg aaa cgc gtt acc ggg Leu Gln Met Pro Pro Tyr Glu Lys Gly Thr Thr Lys Arg Val Thr Gly 165 170 175	528
ttt acc ctg acc ctt gaa gag gcc gta cca ttt gac atg gta gct tat Phe Thr Leu Thr Leu Glu Glu Ala Val Pro Phe Asp Met Val Ala Tyr 180 185 190	576
ggt cga aac ctg atg ctg aag gct tcg gca ggt tcc ttt cca aca att Gly Arg Asn Leu Met Leu Lys Ala Ser Ala Gly Ser Phe Pro Thr Ile 195 200 205	624
gac ttg ctc tat gac tac aga tcg ttt ttt gac caa tgt tcc gat att Asp Leu Leu Tyr Asp Tyr Arg Ser Phe Phe Asp Gln Cys Ser Asp Ile 210 215 220	672
gga cgg atc ggc ttc ttt ccg gaa gat gtt cct aag ccg aaa gtg gcg Gly Arg Ile Gly Phe Phe Pro Glu Asp Val Pro Lys Pro Lys Val Ala 225 230 235 240	720
atc att ggc gct ggc att tcc gga ctc gtg gta gca agc gaa ctg ctt Ile Ile Gly Ala Gly Ile Ser Gly Leu Val Val Ala Ser Glu Leu Leu 245 250 255	768
cat gct ggt gta gac gat gtt aca ata tat gaa gca agt gat cgg gtt His Ala Gly Val Asp Asp Val Thr Ile Tyr Glu Ala Ser Asp Arg Val 260 265 270	816
gga ggc aag ctt tgg tca cat gct ttc aag gat gct ccc agc gtg gtg Gly Gly Lys Leu Trp Ser His Ala Phe Lys Asp Ala Pro Ser Val Val 275 280 285	864
gcc gaa atg ggg gcg atg cga ttt cct cct gct gca tcg tgc ttg ttt Ala Glu Met Gly Ala Met Arg Phe Pro Pro Ala Ala Ser Cys Leu Phe 290 295 300	912
ttc ttc ctc gag ccg tac ggc ctg tct tcg atg agg ccg ttc cca aat Phe Phe Leu Glu Arg Tyr Gly Leu Ser Ser Met Arg Pro Phe Pro Asn 305 310 315 320	960
ccc ggc aca gtc gac act aac ttg gtc tac caa ggc ctc cga tac gtg Pro Gly Thr Val Asp Thr Asn Leu Val Tyr Gln Gly Leu Arg Tyr Val 325 330 335	1008
tgg aaa gcc ggg cag cag cca ccg aag ctg ttc cat cgc gtt tac agc Trp Lys Ala Gly Gln Gln Pro Pro Lys Leu Phe His Arg Val Tyr Ser 340 345 350	1056
ggt tgg cgt gcg ttc ttg agg gac ggt ttc cat gag gga gat att gtg Gly Trp Arg Ala Phe Leu Arg Asp Gly Phe His Glu Gly Asp Ile Val 355 360 365	1104
ttg gct tcg cct gtt gtt att act caa gcc ttg aaa tca gga gac att Leu Ala Ser Pro Val Val Ile Thr Gln Ala Leu Lys Ser Gly Asp Ile 370 375 380	1152

PF 53790

45

agg cgg gct cat gac tcc tgg caa act tgg ctg aac cgt ttc ggg agg	1200
Arg Arg Ala His Asp Ser Trp Gln Thr Trp Leu Asn Arg Phe Gly Arg	
385 390 395 400	
gag tcc ttc tct tca gcg ata gag agg atc ttt ctg ggc acg cat cct	1248
Glu Ser Phe Ser Ser Ala Ile Glu Arg Ile Phe Leu Gly Thr His Pro	
405 410 415	
cct ggt ggt gaa aca tgg agt ttc cct cat gat tgg gac cta ttc aag	1296
Pro Gly Gly Glu Thr Trp Ser Phe Pro His Asp Trp Asp Leu Phe Lys	
420 425 430	
cta atg gga ata gga tct ggc ggg ttt ggt cca gtt ttt gaa agc ggg	1344
Leu Met Gly Ile Gly Ser Gly Gly Phe Gly Pro Val Phe Glu Ser Gly	
435 440 445	
ttt att gag atc ctt cgc ttg gtc ata aac gga tat gaa gaa aat cag	1392
Phe Ile Glu Ile Leu Arg Leu Val Ile Asn Gly Tyr Glu Glu Asn Gln	
450 455 460	
cgg atg tgc tct gaa gga atc tca gaa ctt cca cgt cga ata gcc tct	1440
Arg Met Cys Ser Glu Gly Ile Ser Glu Leu Pro Arg Arg Ile Ala Ser	
465 470 475 480	
caa gtg gtt aac ggt gtg tct gta agc cag cgt ata cgc cat gtt caa	1488
Gln Val Val Asn Gly Val Ser Val Ser Gln Arg Ile Arg His Val Gln	
485 490 495	
gtc agg gcg att gag aag gaa aag aca aaa ata aag ata agg ctt aag	1536
Val Arg Ala Ile Glu Lys Glu Lys Thr Lys Ile Lys Ile Arg Leu Lys	
500 505 510	
agc ggg ata tct gaa ctt tat gat aag gtg gtg gtt aca tct gga ctc	1584
Ser Gly Ile Ser Glu Leu Tyr Asp Lys Val Val Val Thr Ser Gly Leu	
515 520 525	
gca aat atc caa ctc agg cat tgt ctg aca tgc gat acc acc att ttt	1632
Ala Asn Ile Gln Leu Arg His Cys Leu Thr Cys Asp Thr Thr Ile Phe	
530 535 540	
cgt gca cca gtg aac caa gcg gtt gat aac agc cat atg aca ggc tcg	1680
Arg Ala Pro Val Asn Gln Ala Val Asp Asn Ser His Met Thr Gly Ser	
545 550 555 560	
tca aaa ctc ttt ctg ctg act gaa cga aaa ttt tgg tta gac cat atc	1728
Ser Lys Leu Phe Leu Leu Thr Glu Arg Lys Phe Trp Leu Asp His Ile	
565 570 575	
ctc ccg tcc tgt gtc ctc atg gac ggg atc gca aaa gca gtg tac tgc	1776
Leu Pro Ser Cys Val Leu Met Asp Gly Ile Ala Lys Ala Val Tyr Cys	
580 585 590	
ttg gac tat gag ccg cag gat ccg aat ggt aaa ggt ctg gtg ccc ccc	1824
Leu Asp Tyr Glu Pro Gln Asp Pro Asn Gly Lys Gly Leu Val Pro Pro	
595 600 605	
act tat aca tgg gag gac gac tcc cac aag ctg ttg gcg gtt ccc gac	1872
Thr Tyr Thr Trp Glu Asp Asp Ser His Lys Leu Leu Ala Val Pro Asp	
610 615 620	
aaa aaa gag cga ttc tgt ctg ctg ccg gac gca att tcg aga tct ttc	1920
Lys Lys Glu Arg Phe Cys Leu Leu Arg Asp Ala Ile Ser Arg Ser Phe	
625 630 635 640	
ccg gcg ttt gcc cag cat cta gtt cct gcc tgc gct gat tac gac caa	1968
Pro Ala Phe Ala Gln His Leu Val Pro Ala Cys Ala Asp Tyr Asp Gln	
645 650 655	

PF 53790

46

aat gtt gtt caa cat gat tgg ctt aca gac gag aat gcc ggg gga gct	2016
Asn Val Val Gln His Asp Trp Leu Thr Asp Glu Asn Ala Gly Gly Ala	
660 665 670	
ttc aaa ctc aac cgg cgt ggc gag gat ttt tat tct gaa gaa ctt ttc	2064
Phe Lys Leu Asn Arg Arg Gly Glu Asp Phe Tyr Ser Glu Glu Leu Phe	
675 680 685	
ttt caa gcg ctg gac atg cct aat gat acc gga gtt tac ttg gcg ggt	2112
Phe Gln Ala Leu Asp Met Pro Asn Asp Thr Gly Val Tyr Leu Ala Gly	
690 695 700	
tgc agt tgt tcc ttc acc ggt gga tgg gtg gag ggc gct att cag acc	2160
Cys Ser Cys Ser Phe Thr Gly Gly Trp Val Glu Gly Ala Ile Gln Thr	
705 710 715 720	
gcg tgt aac gcc gtc tgt gca att atc cac aat tgt gga ggt att ttg	2208
Ala Cys Asn Ala Val Cys Ala Ile Ile His Asn Cys Gly Gly Ile Leu	
725 730 735	
gca aag gac aat cct ctc gaa cac tct tgg aag aga tat aac tac cgc	2256
Ala Lys Asp Asn Pro Leu Glu His Ser Trp Lys Arg Tyr Asn Tyr Arg	
740 745 750	
aat aga aat taa	2268
Asn Arg Asn	
755	

&lt;210&gt; 32

&lt;211&gt; 755

&lt;212&gt; PRT

&lt;213&gt; Agrobacterium tumefaciens

&lt;400&gt; 32

Met Ser Ala Ser Pro Leu Leu Asp Asn Gln Cys Asp His Phe Ser Thr	
1 5 10 15	
Lys Met Val Asp Leu Ile Met Val Asp Lys Ala Asp Glu Leu Asp Arg	
20 25 30	
Arg Val Ser Asp Ala Phe Ser Glu Arg Glu Ala Ser Arg Gly Arg Arg	
35 40 45	
Ile Thr Gln Ile Ser Gly Glu Cys Ser Ala Gly Leu Ala Cys Lys Arg	
50 55 60	
Leu Ala Asp Gly Arg Phe Pro Glu Ile Ser Thr Gly Glu Lys Val Ala	
65 70 75 80	
Ala Leu Ser Ala Tyr Ile Tyr Val Gly Lys Glu Ile Leu Gly Arg Ile	
85 90 95	
Leu Glu Ser Glu Pro Trp Ala Arg Ala Arg Val Ser Gly Leu Val Ala	
100 105 110	
Ile Asp Leu Ala Pro Phe Cys Met Asp Phe Ser Glu Ala Gln Leu Leu	
115 120 125	
Gln Thr Leu Phe Leu Leu Ser Gly Lys Arg Cys Ala Ser Ser Asp Leu	
130 135 140	
Ser His Phe Val Ala Ile Ser Ile Ser Lys Thr Ala Arg Ser Arg Thr	
145 150 155 160	
Leu Gln Met Pro Pro Tyr Glu Lys Gly Thr Thr Lys Arg Val Thr Gly	
165 170 175	

PF 53790

47

Phe	Thr	Leu	Thr	Leu	Glu	Glu	Ala	Val	Pro	Phe	Asp	Met	Val	Ala	Tyr	180	185	190
Gly	Arg	Asn	Leu	Met	Leu	Lys	Ala	Ser	Ala	Gly	Ser	Phe	Pro	Thr	Ile	195	200	205
Asp	Leu	Leu	Tyr	Asp	Tyr	Arg	Ser	Phe	Phe	Asp	Gln	Cys	Ser	Asp	Ile	210	215	220
Gly	Arg	Ile	Gly	Phe	Phe	Pro	Glu	Asp	Val	Pro	Lys	Pro	Lys	Val	Ala	225	230	235
Ile	Ile	Gly	Ala	Gly	Ile	Ser	Gly	Leu	Val	Val	Ala	Ser	Glu	Leu	Leu	245	250	255
His	Ala	Gly	Val	Asp	Asp	Val	Thr	Ile	Tyr	Glu	Ala	Ser	Asp	Arg	Val	260	265	270
Gly	Gly	Lys	Leu	Trp	Ser	His	Ala	Phe	Lys	Asp	Ala	Pro	Ser	Val	Val	275	280	285
Ala	Glu	Met	Gly	Ala	Met	Arg	Phe	Pro	Pro	Ala	Ala	Ser	Cys	Leu	Phe	290	295	300
Phe	Phe	Leu	Glu	Arg	Tyr	Gly	Leu	Ser	Ser	Met	Arg	Pro	Phe	Pro	Asn	305	310	315
Pro	Gly	Thr	Val	Asp	Thr	Asn	Leu	Val	Tyr	Gln	Gly	Leu	Arg	Tyr	Val	325	330	335
Trp	Lys	Ala	Gly	Gln	Gln	Pro	Pro	Lys	Leu	Phe	His	Arg	Val	Tyr	Ser	340	345	350
Gly	Trp	Arg	Ala	Phe	Leu	Arg	Asp	Gly	Phe	His	Glu	Gly	Asp	Ile	Val	355	360	365
Leu	Ala	Ser	Pro	Val	Val	Ile	Thr	Gln	Ala	Leu	Lys	Ser	Gly	Asp	Ile	370	375	380
Arg	Arg	Ala	His	Asp	Ser	Trp	Gln	Thr	Trp	Leu	Asn	Arg	Phe	Gly	Arg	385	390	395
Glu	Ser	Phe	Ser	Ser	Ala	Ile	Glu	Arg	Ile	Phe	Leu	Gly	Thr	His	Pro	405	410	415
Pro	Gly	Gly	Glu	Thr	Trp	Ser	Phe	Pro	His	Asp	Trp	Asp	Leu	Phe	Lys	420	425	430
Leu	Met	Gly	Ile	Gly	Ser	Gly	Gly	Phe	Gly	Pro	Val	Phe	Glu	Ser	Gly	435	440	445
Phe	Ile	Glu	Ile	Leu	Arg	Leu	Val	Ile	Asn	Gly	Tyr	Glu	Glu	Asn	Gln	450	455	460
Arg	Met	Cys	Ser	Glu	Gly	Ile	Ser	Glu	Leu	Pro	Arg	Arg	Ile	Ala	Ser	465	470	475
Gln	Val	Val	Asn	Gly	Val	Ser	Val	Ser	Gln	Arg	Ile	Arg	His	Val	Gln	485	490	495
Val	Arg	Ala	Ile	Glu	Lys	Glu	Lys	Thr	Lys	Ile	Lys	Ile	Arg	Leu	Lys	500	505	510
Ser	Gly	Ile	Ser	Glu	Leu	Tyr	Asp	Lys	Val	Val	Val	Thr	Ser	Gly	Leu	515	520	525
Ala	Asn	Ile	Gln	Leu	Arg	His	Cys	Leu	Thr	Cys	Asp	Thr	Thr	Ile	Phe	530	535	540

PF 53790

48

Arg Ala Pro Val Asn Gln Ala Val Asp Asn Ser His Met Thr Gly Ser  
 545 550 555 560  
 Ser Lys Leu Phe Leu Leu Thr Glu Arg Lys Phe Trp Leu Asp His Ile  
 565 570 575  
 Leu Pro Ser Cys Val Leu Met Asp Gly Ile Ala Lys Ala Val Tyr Cys  
 580 585 590  
 Leu Asp Tyr Glu Pro Gln Asp Pro Asn Gly Lys Gly Leu Val Pro Pro  
 595 600 605  
 Thr Tyr Thr Trp Glu Asp Asp Ser His Lys Leu Leu Ala Val Pro Asp  
 610 615 620  
 Lys Lys Glu Arg Phe Cys Leu Leu Arg Asp Ala Ile Ser Arg Ser Phe  
 625 630 635 640  
 Pro Ala Phe Ala Gln His Leu Val Pro Ala Cys Ala Asp Tyr Asp Gln  
 645 650 655  
 Asn Val Val Gln His Asp Trp Leu Thr Asp Glu Asn Ala Gly Gly Ala  
 660 665 670  
 Phe Lys Leu Asn Arg Arg Gly Glu Asp Phe Tyr Ser Glu Glu Leu Phe  
 675 680 685  
 Phe Gln Ala Leu Asp Met Pro Asn Asp Thr Gly Val Tyr Leu Ala Gly  
 690 695 700  
 Cys Ser Cys Ser Phe Thr Gly Gly Trp Val Glu Gly Ala Ile Gln Thr  
 705 710 715 720  
 Ala Cys Asn Ala Val Cys Ala Ile Ile His Asn Cys Gly Gly Ile Leu  
 725 730 735  
 Ala Lys Asp Asn Pro Leu Glu His Ser Trp Lys Arg Tyr Asn Tyr Arg  
 740 745 750  
 Asn Arg Asn  
 755

&lt;210&gt; 33

&lt;211&gt; 1404

&lt;212&gt; DNA

&lt;213&gt; Agrobacterium tumefaciens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1401)

&lt;223&gt; coding for indoleacetamide hydrolase

&lt;400&gt; 33

atg	gtg	ccc	att	acc	tgc	tta	gca	caa	acc	cta	gaa	cgc	ctg	aga	cgg	48
Met	Val	Pro	Ile	Thr	Ser	Leu	Ala	Gln	Thr	Leu	Glu	Arg	Leu	Arg	Arg	
1				5					10					15		
aaa	gac	tac	tcc	tgc	tta	gaa	cta	gta	gaa	act	ctg	ata	gcg	cgt	tgc	96
Lys	Asp	Tyr	Ser	Cys	Leu	Glu	Leu	Val	Glu	Thr	Leu	Ile	Ala	Arg	Cys	
			20					25					30			
caa	gct	gca	aaa	cca	tta	aat	gcc	ctt	ctg	gct	aca	gac	tgg	gat	ggc	144
Gln	Ala	Ala	Lys	Pro	Leu	Asn	Ala	Leu	Leu	Ala	Thr	Asp	Trp	Asp	Gly	
			35				40						45			



PF 53790

49

ttg cgg cga agc gcc aaa aaa aat gat cgt cat gga aac gcc gga tta	192
Leu Arg Arg Ser Ala Lys Lys Asn Asp Arg His Gly Asn Ala Gly Leu	
50 55 60	
ggt ctt tgc ggc att cca ctc tgt ttt aag gcg aac atc gcg acc ggc	240
Gly Leu Cys Gly Ile Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly	
65 70 75 80	
gta ttt cct aca agc gct gct act ccg gcg ctg ata aac cac ttg cca	288
Val Phe Pro Thr Ser Ala Ala Thr Pro Ala Leu Ile Asn His Leu Pro	
85 90 95	
aag ata cca tcc cgc gtc gca gaa aga ctt ttt tca gct gga gca ctg	336
Lys Ile Pro Ser Arg Val Ala Glu Arg Leu Phe Ser Ala Gly Ala Leu	
100 105 110	
ccg ggt gcc tcg gga aac atg cat gag tta tcg ttt gga att acg agc	384
Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser	
115 120 125	
aac aac tat gcc acc ggt gcg gtg cgg aac ccg tgg aat cca agt ctg	432
Asn Asn Tyr Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu	
130 135 140	
ata cca ggg ggt tca agc ggt ggt gtg gct gct gcg gtg gca agc cga	480
Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Val Ala Ser Arg	
145 150 155 160	
ttg atg tta ggc ggc ata ggc acg gat acc ggt gca tct gtt cgc cta	528
Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu	
165 170 175	
ccg gca gcc ctg tgt ggc gta gta gga ttt cga ccg acg ctt ggt cga	576
Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Leu Gly Arg	
180 185 190	
tat cca aga gat cgg ata ata ccg ttc agc ccc acc cgg gac acc gcc	624
Tyr Pro Arg Asp Arg Ile Ile Pro Phe Ser Pro Thr Arg Asp Thr Ala	
195 200 205	
gga atc ata gcg cag tgc gta gcc gat gtt ata atc ctc gac cag gtg	672
Gly Ile Ile Ala Gln Cys Val Ala Asp Val Ile Ile Leu Asp Gln Val	
210 215 220	
att tcc gga cgg tcg gcg aaa att tca ccc atg ccg ctg aag ggg ctt	720
Ile Ser Gly Arg Ser Ala Lys Ile Ser Pro Met Pro Leu Lys Gly Leu	
225 230 235 240	
cgg atc ggc ctc ccc act acc tac ttt tac gat gac ctt gat gct gat	768
Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp	
245 250 255	
gtg gcc ttc gca gct gaa acg acg att cgc ttg cta gcc aac aga ggc	816
Val Ala Phe Ala Ala Glu Thr Thr Ile Arg Leu Leu Ala Asn Arg Gly	
260 265 270	
gta acc ttt gtt gaa gcc gac atc ccc cac cta gag gaa ttg aac agt	864
Val Thr Phe Val Glu Ala Asp Ile Pro His Leu Glu Glu Leu Asn Ser	
275 280 285	
ggg gca agt ttg cca att gcg ctt tac gaa ttt cca cac gct cta aaa	912
Gly Ala Ser Leu Pro Ile Ala Leu Tyr Glu Phe Pro His Ala Leu Lys	
290 295 300	
aag tat ctc gac gat ttt gtg gga aca gtt tct ttt tct gac gtt atc	960
Lys Tyr Leu Asp Asp Phe Val Gly Thr Val Ser Phe Ser Asp Val Ile	
305 310 315 320	

PF 53790

50

```

aaa gga att cgt agc ccc gat gta gcg aac att gtc agt gcg caa att 1008
Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Val Ser Ala Gln Ile
      325      330      335

gat ggg cat caa att tcc aac gat gaa tat gaa ctg gcg cgt caa tcc 1056
Asp Gly His Gln Ile Ser Asn Asp Glu Tyr Glu Leu Ala Arg Gln Ser
      340      345      350

ttc agg cca agg ctc cag gcc act tat cgg aat tac ttc aga ctc tat 1104
Phe Arg Pro Arg Leu Gln Ala Thr Tyr Arg Asn Tyr Phe Arg Leu Tyr
      355      360      365

cag tta gat gca atc ctt ttc cca act gca ccc tta gcg gcc aaa gcc 1152
Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Ala Ala Lys Ala
      370      375      380

ata ggt cag gag tcg tca gtc atc cac aat ggc tca atg atg aac act 1200
Ile Gly Gln Glu Ser Ser Val Ile His Asn Gly Ser Met Met Asn Thr
      385      390      395      400

ttc aag atc tac gtg cga aat gtg gac cca agc agc aac gca ggc cta 1248
Phe Lys Ile Tyr Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu
      405      410      415

cct ggg ttg agc ctt cct gcc tgc ctt aca cct gat cgc ttg cct gtt 1296
Pro Gly Leu Ser Leu Pro Ala Cys Leu Thr Pro Asp Arg Leu Pro Val
      420      425      430

gga atg gaa att gat gga tta gcg ggg tca gac cac cgt ctg tta gca 1344
Gly Met Glu Ile Asp Gly Leu Ala Gly Ser Asp His Arg Leu Leu Ala
      435      440      445

atc ggg gca gca tta gaa aaa gct ata aat ttt tct tcc ttt ccc gat 1392
Ile Gly Ala Ala Leu Glu Lys Ala Ile Asn Phe Ser Ser Phe Pro Asp
      450      455      460

gct ttt aat tag 1404
Ala Phe Asn
465

```

&lt;210&gt; 34

&lt;211&gt; 467

&lt;212&gt; PRT

&lt;213&gt; Agrobacterium tumefaciens

&lt;400&gt; 34

```

Met Val Pro Ile Thr Ser Leu Ala Gln Thr Leu Glu Arg Leu Arg Arg
  1           5           10           15

Lys Asp Tyr Ser Cys Leu Glu Leu Val Glu Thr Leu Ile Ala Arg Cys
      20           25           30

Gln Ala Ala Lys Pro Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Gly
      35           40           45

Leu Arg Arg Ser Ala Lys Lys Asn Asp Arg His Gly Asn Ala Gly Leu
      50           55           60

Gly Leu Cys Gly Ile Pro Leu Cys Phe Lys Ala Asn Ile Ala Thr Gly
      65           70           75           80

Val Phe Pro Thr Ser Ala Ala Thr Pro Ala Leu Ile Asn His Leu Pro
      85           90           95

Lys Ile Pro Ser Arg Val Ala Glu Arg Leu Phe Ser Ala Gly Ala Leu
      100          105          110

```

PF 53790

51

Pro Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser  
 115 120 125  
 Asn Asn Tyr Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu  
 130 135 140  
 Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Val Ala Ser Arg  
 145 150 155 160  
 Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu  
 165 170 175  
 Pro Ala Ala Leu Cys Gly Val Val Gly Phe Arg Pro Thr Leu Gly Arg  
 180 185 190  
 Tyr Pro Arg Asp Arg Ile Ile Pro Phe Ser Pro Thr Arg Asp Thr Ala  
 195 200 205  
 Gly Ile Ile Ala Gln Cys Val Ala Asp Val Ile Ile Leu Asp Gln Val  
 210 215 220  
 Ile Ser Gly Arg Ser Ala Lys Ile Ser Pro Met Pro Leu Lys Gly Leu  
 225 230 235 240  
 Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp  
 245 250 255  
 Val Ala Phe Ala Ala Glu Thr Thr Ile Arg Leu Leu Ala Asn Arg Gly  
 260 265 270  
 Val Thr Phe Val Glu Ala Asp Ile Pro His Leu Glu Glu Leu Asn Ser  
 275 280 285  
 Gly Ala Ser Leu Pro Ile Ala Leu Tyr Glu Phe Pro His Ala Leu Lys  
 290 295 300  
 Lys Tyr Leu Asp Asp Phe Val Gly Thr Val Ser Phe Ser Asp Val Ile  
 305 310 315 320  
 Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Val Ser Ala Gln Ile  
 325 330 335  
 Asp Gly His Gln Ile Ser Asn Asp Glu Tyr Glu Leu Ala Arg Gln Ser  
 340 345 350  
 Phe Arg Pro Arg Leu Gln Ala Thr Tyr Arg Asn Tyr Phe Arg Leu Tyr  
 355 360 365  
 Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Ala Ala Lys Ala  
 370 375 380  
 Ile Gly Gln Glu Ser Ser Val Ile His Asn Gly Ser Met Met Asn Thr  
 385 390 395 400  
 Phe Lys Ile Tyr Val Arg Asn Val Asp Pro Ser Ser Asn Ala Gly Leu  
 405 410 415  
 Pro Gly Leu Ser Leu Pro Ala Cys Leu Thr Pro Asp Arg Leu Pro Val  
 420 425 430  
 Gly Met Glu Ile Asp Gly Leu Ala Gly Ser Asp His Arg Leu Leu Ala  
 435 440 445  
 Ile Gly Ala Ala Leu Glu Lys Ala Ile Asn Phe Ser Ser Phe Pro Asp  
 450 455 460  
 Ala Phe Asn  
 465

PF 53790

52

<210> 35  
 <211> 1419  
 <212> DNA  
 <213> Agrobacterium vitis

<220>  
 <221> CDS  
 <222> (1)..(1416)  
 <223> coding for indoleacetamide hydrolase

<400> 35  
 atg gtg acc cta ggt tca atc aag gaa acc ctg gaa tgt ctc agg ctg 48  
 Met Val Thr Leu Gly Ser Ile Lys Glu Thr Leu Glu Cys Leu Arg Leu  
 1 5 10 15  
 aaa aaa tac tcc tgt tcc gaa ctg gct gaa acc ata ata gcc cgt tgc 96  
 Lys Lys Tyr Ser Cys Ser Glu Leu Ala Glu Thr Ile Ile Ala Arg Cys  
 20 25 30  
 gaa gcc gcg aaa tct ctc aat gct ctt ctg gcg act gac tgg gat tac 144  
 Glu Ala Ala Lys Ser Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Tyr  
 35 40 45  
 ctg cgg cgt aat gcc aag aaa gta gat gaa gat gga agc gcc ggc gag 192  
 Leu Arg Arg Asn Ala Lys Lys Val Asp Glu Asp Gly Ser Ala Gly Glu  
 50 55 60  
 ggt ctt gcc ggc atc ccg ctg tgt tct aaa gcg aac att gca aca ggc 240  
 Gly Leu Ala Gly Ile Pro Leu Cys Ser Lys Ala Asn Ile Ala Thr Gly  
 65 70 75 80  
 ata ttc cca gca agc gcg gcc acg ccg gcg ctt gat gaa cat tta cct 288  
 Ile Phe Pro Ala Ser Ala Ala Thr Pro Ala Leu Asp Glu His Leu Pro  
 85 90 95  
 aca aca cca gcc ggc gtc cgt aaa ccg ctt cta gac gct ggg gca ctg 336  
 Thr Thr Pro Ala Gly Val Arg Lys Pro Leu Leu Asp Ala Gly Ala Leu  
 100 105 110  
 ata ggc gct tcg gga aac atg cat gag tta tcg ttt ggc att acc agt 384  
 Ile Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser  
 115 120 125  
 aac aac cac gcc act ggt gcg gtg aga aac ccc tgg aat ccc agc tta 432  
 Asn Asn His Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu  
 130 135 140  
 ata cca gga ggc tcg agc ggc ggc gtg gct gct gct gta gca tca cgg 480  
 Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Ala Val Ala Ser Arg  
 145 150 155 160  
 tta atg ctc ggc gga att ggc acc gac acg ggg gct tcg gtc cgc cta 528  
 Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu  
 165 170 175  
 cct gca tcc cta tgt ggc gta gtg gga ttc cgc ccg acg atc ggc aga 576  
 Pro Ala Ser Leu Cys Gly Val Val Gly Phe Arg Pro Thr Ile Gly Arg  
 180 185 190  
 tat cct gga gac cga att gtg ccg gtt agc ccc acc cgc gat aca gcc 624  
 Tyr Pro Gly Asp Arg Ile Val Pro Val Ser Pro Thr Arg Asp Thr Ala  
 195 200 205

PF 53790

53

gga att atc gca cag agc gtt cct gat gtg ata ctc ctt gac caa atc	672
Gly Ile Ile Ala Gln Ser Val Pro Asp Val Ile Leu Leu Asp Gln Ile	
210 215 220	
att tgc ggg aag ctc acg acc cac caa cct gta ccc ctg gag gga tta	720
Ile Cys Gly Lys Leu Thr Thr His Gln Pro Val Pro Leu Glu Gly Leu	
225 230 235 240	
cgt atc ggc ttg cca acc act tac ttt tac gat gac ctt gat gct gat	768
Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp	
245 250 255	
gtg gcc ttc gca gct gaa aac ctt atc acg ctg ctg gcc agc aag ggt	816
Val Ala Phe Ala Ala Glu Asn Leu Ile Thr Leu Leu Ala Ser Lys Gly	
260 265 270	
gta acc ttt gtt aag gcc gag att cca gat ctg cag cgt ctg aac atc	864
Val Thr Phe Val Lys Ala Glu Ile Pro Asp Leu Gln Arg Leu Asn Ile	
275 280 285	
ggg gtt agc ttt cct att gcc ctg tac gag ttt ccg ttc gcc cta caa	912
Gly Val Ser Phe Pro Ile Ala Leu Tyr Glu Phe Pro Phe Ala Leu Gln	
290 295 300	
aag tat atc gat gac ttt gtg aag gat gtg tct ttt tct gac gtc atc	960
Lys Tyr Ile Asp Asp Phe Val Lys Asp Val Ser Phe Ser Asp Val Ile	
305 310 315 320	
aaa gga att cgt agc cct gat gta gcc aac att gcc aat gct caa att	1008
Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Ala Asn Ala Gln Ile	
325 330 335	
gat gga cat caa att tcc aaa gct tca tat gaa ctg gcg cga caa tct	1056
Asp Gly His Gln Ile Ser Lys Ala Ser Tyr Glu Leu Ala Arg Gln Ser	
340 345 350	
ttc aga cca aag ctg caa gcc gcc tac cat gat tac ttc aag ctg cac	1104
Phe Arg Pro Lys Leu Gln Ala Ala Tyr His Asp Tyr Phe Lys Leu His	
355 360 365	
cag cta gac gcg atc ctt ttc ccg aca gct ccc ctg aca gcc aaa ccg	1152
Gln Leu Asp Ala Ile Leu Phe Pro Thr Ala Pro Leu Thr Ala Lys Pro	
370 375 380	
atc ggc caa gat tta tcg gtg atg cac aat ggc gta atg gcc gac acg	1200
Ile Gly Gln Asp Leu Ser Val Met His Asn Gly Val Met Ala Asp Thr	
385 390 395 400	
ttt aaa atc ttc gtg cga aat gtg gat ccg ggg agc aac gca ggc ctg	1248
Phe Lys Ile Phe Val Arg Asn Val Asp Pro Gly Ser Asn Ala Gly Leu	
405 410 415	
cca gga tta agc ctt ccc gtt tct ctt act tca aag ggt ttg cct att	1296
Pro Gly Leu Ser Leu Pro Val Ser Leu Thr Ser Lys Gly Leu Pro Ile	
420 425 430	
gga atg gaa atc gat gga tta gcg ggc atg gac gac cgt ttg cta gca	1344
Gly Met Glu Ile Asp Gly Leu Ala Gly Met Asp Asp Arg Leu Leu Ala	
435 440 445	
atc gga gcg gca cta gag gaa gcg ata gct ttt cat aat tta cct gac	1392
Ile Gly Ala Ala Leu Glu Glu Ala Ile Ala Phe His Asn Leu Pro Asp	
450 455 460	
ttc ccg aaa gtc gag aca aac tac tga	1419
Phe Pro Lys Val Glu Thr Asn Tyr	
465 470	

PF 53790

54

&lt;210&gt; 36

&lt;211&gt; 472

&lt;212&gt; PRT

&lt;213&gt; Agrobacterium vitis

&lt;400&gt; 36

```

Met Val Thr Leu Gly Ser Ile Lys Glu Thr Leu Glu Cys Leu Arg Leu
 1           5           10           15
Lys Lys Tyr Ser Cys Ser Glu Leu Ala Glu Thr Ile Ile Ala Arg Cys
 20           25           30
Glu Ala Ala Lys Ser Leu Asn Ala Leu Leu Ala Thr Asp Trp Asp Tyr
 35           40           45
Leu Arg Arg Asn Ala Lys Lys Val Asp Glu Asp Gly Ser Ala Gly Glu
 50           55           60
Gly Leu Ala Gly Ile Pro Leu Cys Ser Lys Ala Asn Ile Ala Thr Gly
 65           70           75           80
Ile Phe Pro Ala Ser Ala Ala Thr Pro Ala Leu Asp Glu His Leu Pro
 85           90           95
Thr Thr Pro Ala Gly Val Arg Lys Pro Leu Leu Asp Ala Gly Ala Leu
100          105          110
Ile Gly Ala Ser Gly Asn Met His Glu Leu Ser Phe Gly Ile Thr Ser
115          120          125
Asn Asn His Ala Thr Gly Ala Val Arg Asn Pro Trp Asn Pro Ser Leu
130          135          140
Ile Pro Gly Gly Ser Ser Gly Gly Val Ala Ala Val Ala Ser Arg
145          150          155          160
Leu Met Leu Gly Gly Ile Gly Thr Asp Thr Gly Ala Ser Val Arg Leu
165          170          175
Pro Ala Ser Leu Cys Gly Val Val Gly Phe Arg Pro Thr Ile Gly Arg
180          185          190
Tyr Pro Gly Asp Arg Ile Val Pro Val Ser Pro Thr Arg Asp Thr Ala
195          200          205
Gly Ile Ile Ala Gln Ser Val Pro Asp Val Ile Leu Leu Asp Gln Ile
210          215          220
Ile Cys Gly Lys Leu Thr Thr His Gln Pro Val Pro Leu Glu Gly Leu
225          230          235          240
Arg Ile Gly Leu Pro Thr Thr Tyr Phe Tyr Asp Asp Leu Asp Ala Asp
245          250          255
Val Ala Phe Ala Ala Glu Asn Leu Ile Thr Leu Leu Ala Ser Lys Gly
260          265          270
Val Thr Phe Val Lys Ala Glu Ile Pro Asp Leu Gln Arg Leu Asn Ile
275          280          285
Gly Val Ser Phe Pro Ile Ala Leu Tyr Glu Phe Pro Phe Ala Leu Gln
290          295          300
Lys Tyr Ile Asp Asp Phe Val Lys Asp Val Ser Phe Ser Asp Val Ile
305          310          315          320
Lys Gly Ile Arg Ser Pro Asp Val Ala Asn Ile Ala Asn Ala Gln Ile
325          330          335

```



PF 53790

56

ttg att gga atg aga tac ctt gag cct cct cat atc att ctc cgc aaa	384
Leu Ile Gly Met Arg Tyr Leu Glu Pro Pro His Ile Ile Leu Arg Lys	
115 120 125	
gga ctc att gct ggg att gag tat cct ttc ctc gca gac cac atg tct	432
Gly Leu Ile Ala Gly Ile Glu Tyr Pro Phe Leu Ala Asp His Met Ser	
130 135 140	
gat tac atg gcg aag act ctc ttc ttc act tct ctc ctc tat cac gat	480
Asp Tyr Met Ala Lys Thr Leu Phe Phe Thr Ser Leu Leu Tyr His Asp	
145 150 155 160	
acc aca gag cac aga aga gca gta acc gaa ttt tgt ggt aat gtg gag	528
Thr Thr Glu His Arg Arg Ala Val Thr Glu Phe Cys Gly Asn Val Glu	
165 170 175	
tta tgc cga tta acg gag caa gtt gtg ttt tcg gac cca tat aga gtt	576
Leu Cys Arg Leu Thr Glu Gln Val Val Phe Ser Asp Pro Tyr Arg Val	
180 185 190	
tcc aca ttt aat cgt tgg act tca cct tat ctt gat gat gat gct aag	624
Ser Thr Phe Asn Arg Trp Thr Ser Pro Tyr Leu Asp Asp Ala Lys	
195 200 205	
gct gtg cgc gaa gac agt gcc ttg aag ctc gaa atc gca gag cta aaa	672
Ala Val Arg Glu Asp Ser Ala Leu Lys Leu Glu Ile Ala Glu Leu Lys	
210 215 220	
tcg atg ttc tgt gaa aga gct caa gct tta ata cat ggt gat ctt cat	720
Ser Met Phe Cys Glu Arg Ala Gln Ala Leu Ile His Gly Asp Leu His	
225 230 235 240	
act ggt tct gtc atg gtt act caa gat tca acg caa gtt ata gat cca	768
Thr Gly Ser Val Met Val Thr Gln Asp Ser Thr Gln Val Ile Asp Pro	
245 250 255	
gag ttt tcg ttc tat gga ccg atg ggt ttc gat att ggc gct tat ctt	816
Glu Phe Ser Phe Tyr Gly Pro Met Gly Phe Asp Ile Gly Ala Tyr Leu	
260 265 270	
ggt aac ttg ata cta gct ttc ttt gca caa gat gga cac gcc act cag	864
Gly Asn Leu Ile Leu Ala Phe Phe Ala Gln Asp Gly His Ala Thr Gln	
275 280 285	
gaa aat gat cga aaa gaa tac aag cag tgg atc ttg aga acc att gag	912
Glu Asn Asp Arg Lys Glu Tyr Lys Gln Trp Ile Leu Arg Thr Ile Glu	
290 295 300	
caa act tgg aat ttg ttt aac aaa agg ttc att gcg cta tgg gat caa	960
Gln Thr Trp Asn Leu Phe Asn Lys Arg Phe Ile Ala Leu Trp Asp Gln	
305 310 315 320	
aac aaa gat gga cca ggc gaa gca tac ctt gca gat atc tat aac aat	1008
Asn Lys Asp Gly Pro Gly Glu Ala Tyr Leu Ala Asp Ile Tyr Asn Asn	
325 330 335	
acc gag gtt ttg aag ttt gtt caa gaa aac tac atg agg aat ttg ttg	1056
Thr Glu Val Leu Lys Phe Val Gln Glu Asn Tyr Met Arg Asn Leu Leu	
340 345 350	
cat gac tca ctc gga ttc ggc gct gca aag atg att agg aga att gtg	1104
His Asp Ser Leu Gly Phe Gly Ala Ala Lys Met Ile Arg Arg Ile Val	
355 360 365	
gga gtg gca cat gtt gag gac ttt gaa tca atc gaa gaa gat aag cga	1152
Gly Val Ala His Val Glu Asp Phe Glu Ser Ile Glu Glu Asp Lys Arg	
370 375 380	



PF 53790

57

```

aga gct att tgc gag aga agt gca ctc gag ttt gcg aag atg ctt ctc 1200
Arg Ala Ile Cys Glu Arg Ser Ala Leu Glu Phe Ala Lys Met Leu Leu
385          390          395          400

aag gaa agg aga aag ttt aag agt atc ggt gaa gtt gtt tca gca att 1248
Lys Glu Arg Arg Lys Phe Lys Ser Ile Gly Glu Val Val Ser Ala Ile
          405          410          415

caa caa caa agc taa 1263
Gln Gln Gln Ser
          420

```

<210> 38  
 <211> 420  
 <212> PRT  
 <213> Arabidopsis thaliana

```

<400> 38
Met Ser Phe Glu Glu Phe Thr Pro Leu Asn Glu Lys Ser Leu Val Asp
 1          5          10          15

Tyr Ile Lys Ser Thr Pro Ala Leu Ser Ser Lys Ile Gly Ala Asp Lys
          20          25          30

Ser Asp Asp Asp Leu Val Ile Lys Glu Val Gly Asp Gly Asn Leu Asn
          35          40          45

Phe Val Phe Ile Val Val Gly Ser Ser Gly Ser Leu Val Ile Lys Gln
          50          55          60

Ala Leu Pro Tyr Ile Arg Cys Ile Gly Glu Ser Trp Pro Met Thr Lys
          65          70          75          80

Glu Arg Ala Tyr Phe Glu Ala Thr Thr Leu Arg Lys His Gly Asn Leu
          85          90          95

Ser Pro Asp His Val Pro Glu Val Tyr His Phe Asp Arg Thr Met Ala
          100          105          110

Leu Ile Gly Met Arg Tyr Leu Glu Pro Pro His Ile Ile Leu Arg Lys
          115          120          125

Gly Leu Ile Ala Gly Ile Glu Tyr Pro Phe Leu Ala Asp His Met Ser
          130          135          140

Asp Tyr Met Ala Lys Thr Leu Phe Phe Thr Ser Leu Leu Tyr His Asp
          145          150          155          160

Thr Thr Glu His Arg Arg Ala Val Thr Glu Phe Cys Gly Asn Val Glu
          165          170          175

Leu Cys Arg Leu Thr Glu Gln Val Val Phe Ser Asp Pro Tyr Arg Val
          180          185          190

Ser Thr Phe Asn Arg Trp Thr Ser Pro Tyr Leu Asp Asp Asp Ala Lys
          195          200          205

Ala Val Arg Glu Asp Ser Ala Leu Lys Leu Glu Ile Ala Glu Leu Lys
          210          215          220

Ser Met Phe Cys Glu Arg Ala Gln Ala Leu Ile His Gly Asp Leu His
          225          230          235          240

Thr Gly Ser Val Met Val Thr Gln Asp Ser Thr Gln Val Ile Asp Pro
          245          250          255

```

PF 53790

58

Glu Phe Ser Phe Tyr Gly Pro Met Gly Phe Asp Ile Gly Ala Tyr Leu  
                   260                  265                  270  
 Gly Asn Leu Ile Leu Ala Phe Phe Ala Gln Asp Gly His Ala Thr Gln  
                   275                  280                  285  
 Glu Asn Asp Arg Lys Glu Tyr Lys Gln Trp Ile Leu Arg Thr Ile Glu  
                   290                  295                  300  
 Gln Thr Trp Asn Leu Phe Asn Lys Arg Phe Ile Ala Leu Trp Asp Gln  
                   305                  310                  315                  320  
 Asn Lys Asp Gly Pro Gly Glu Ala Tyr Leu Ala Asp Ile Tyr Asn Asn  
                   325                  330                  335  
 Thr Glu Val Leu Lys Phe Val Gln Glu Asn Tyr Met Arg Asn Leu Leu  
                   340                  345                  350  
 His Asp Ser Leu Gly Phe Gly Ala Ala Lys Met Ile Arg Arg Ile Val  
                   355                  360                  365  
 Gly Val Ala His Val Glu Asp Phe Glu Ser Ile Glu Glu Asp Lys Arg  
                   370                  375                  380  
 Arg Ala Ile Cys Glu Arg Ser Ala Leu Glu Phe Ala Lys Met Leu Leu  
                   385                  390                  395                  400  
 Lys Glu Arg Arg Lys Phe Lys Ser Ile Gly Glu Val Val Ser Ala Ile  
                   405                  410                  415  
 Gln Gln Gln Ser  
                   420

<210> 39  
 <211> 1200  
 <212> DNA  
 <213> *Klebsiella pneumoniae*  
 <220>  
 <221> CDS  
 <222> (1)..(1197)  
 <223> coding for 5-methylthioribose kinase

<400> 39  
 atg tcg caa tac cat acc ttc acc gcc cac gat gcc gtg gct tac gcg 48  
 Met Ser Gln Tyr His Thr Phe Thr Ala His Asp Ala Val Ala Tyr Ala  
   1                  5                  10                  15  
 caa cag ttc gcc ggc atc gac aac cca tct gag ctg gtc agc gcg cag 96  
 Gln Gln Phe Ala Gly Ile Asp Asn Pro Ser Glu Leu Val Ser Ala Gln  
                   20                  25                  30  
 gaa gtg ggc gat ggc aac ctc aat ctg gtg ttt aaa gtg ttc gat cgt 144  
 Glu Val Gly Asp Gly Asn Leu Asn Leu Val Phe Lys Val Phe Asp Arg  
                   35                  40                  45  
 cag ggc gtc agc cgg gcg atc gtc aaa cag gcc ctg ccc tac gtg cgc 192  
 Gln Gly Val Ser Arg Ala Ile Val Lys Gln Ala Leu Pro Tyr Val Arg  
                   50                  55                  60  
 tgc gtc ggc gaa tcc tgg ccg ctg acc ctc gac cgc gcc cgt ctc gaa 240  
 Cys Val Gly Glu Ser Trp Pro Leu Thr Leu Asp Arg Ala Arg Leu Glu  
   65                  70                  75                  80

PF 53790

59

gcg cag acc ctg gtc gcc cac tat cag cac agc ccg cag cac acg gta	288
Ala Gln Thr Leu Val Ala His Tyr Gln His Ser Pro Gln His Thr Val	
85 90 95	
aaa atc cat cac ttt gat ccc gag ctg gcg gtg atg gtg atg gaa gat	336
Lys Ile His His Phe Asp Pro Glu Leu Ala Val Met Val Met Glu Asp	
100 105 110	
ctt tcc gac cac cgc atc tgg cgc gga gag ctt atc gct aac gtc tac	384
Leu Ser Asp His Arg Ile Trp Arg Gly Glu Leu Ile Ala Asn Val Tyr	
115 120 125	
tat ccc cag gcg gcc cgc cag ctt ggc gac tat ctg gcg cag gtg ttg	432
Tyr Pro Gln Ala Ala Arg Gln Leu Gly Asp Tyr Leu Ala Gln Val Leu	
130 135 140	
ttc cac acc agc gat ttc tac ctc cat ccc cac gag aaa aag gcg cag	480
Phe His Thr Ser Asp Phe Tyr Leu His Pro His Glu Lys Lys Ala Gln	
145 150 155 160	
gtg gcg cag ttt att aac ccg gcg atg tgc gag atc acc gag gat ctg	528
Val Ala Gln Phe Ile Asn Pro Ala Met Cys Glu Ile Thr Glu Asp Leu	
165 170 175	
ttc ttt aac gac ccg tat cag atc cac gag cgc aat aac tac ccg gcg	576
Phe Phe Asn Asp Pro Tyr Gln Ile His Glu Arg Asn Asn Tyr Pro Ala	
180 185 190	
gag ctg gag gcc gat gtc gcc gcc ctg cgc gac gac gcc cag ctt aag	624
Glu Leu Glu Ala Asp Val Ala Ala Leu Arg Asp Asp Ala Gln Leu Lys	
195 200 205	
ctg gcg gtg gcg gcg ctg aag cac cgt ttc ttt gcc cat gcg gaa gcg	672
Leu Ala Val Ala Ala Leu Lys His Arg Phe Phe Ala His Ala Glu Ala	
210 215 220	
ctg ctg cac ggc gat atc cac agc ggg tcg atc ttc gtt gcc gaa ggt	720
Leu Leu His Gly Asp Ile His Ser Gly Ser Ile Phe Val Ala Glu Gly	
225 230 235 240	
agc ctg aag gcc atc gac gcc gag ttc ggc tac ttc ggc ccc atc ggc	768
Ser Leu Lys Ala Ile Asp Ala Glu Phe Gly Tyr Phe Gly Pro Ile Gly	
245 250 255	
ttc gat atc ggc acc gcc atc ggc aac ctg ctg ctg aac tac tgc ggc	816
Phe Asp Ile Gly Thr Ala Ile Gly Asn Leu Leu Leu Asn Tyr Cys Gly	
260 265 270	
ctg ccg ggc cag ctc ggc att cgc gat gcc gcc gcc gcg cgc gag cag	864
Leu Pro Gly Gln Leu Gly Ile Arg Asp Ala Ala Ala Arg Glu Gln	
275 280 285	
cgg ctg aac gac atc cac cag ctg tgg acc acc ttc gcc gag cgc ttc	912
Arg Leu Asn Asp Ile His Gln Leu Trp Thr Thr Phe Ala Glu Arg Phe	
290 295 300	
cag gcg ctg gcg gcg gag aaa acc cgc gac gcg gcg ctg gct tac ccc	960
Gln Ala Leu Ala Ala Glu Lys Thr Arg Asp Ala Ala Leu Ala Tyr Pro	
305 310 315 320	
ggc tac gcc tcc gcc ttt ctg aag aaa gtc tgg gcg gac gcg gtc ggc	1008
Gly Tyr Ala Ser Ala Phe Leu Lys Lys Val Trp Ala Asp Ala Val Gly	
325 330 335	
ttc tgc ggc agc gaa ctg atc cgc cgc agc gtc gga ctg tcg cac gtc	1056
Phe Cys Gly Ser Glu Leu Ile Arg Arg Ser Val Gly Leu Ser His Val	
340 345 350	

<400> 40																
Met	Ser	Gln	Tyr	His	Thr	Phe	Thr	Ala	His	Asp	Ala	Val	Ala	Tyr	Ala	
1				5					10					15		
Gln	Gln	Phe	Ala	Gly	Ile	Asp	Asn	Pro	Ser	Glu	Leu	Val	Ser	Ala	Gln	
			20					25					30			
Glu	Val	Gly	Asp	Gly	Asn	Leu	Asn	Leu	Val	Phe	Lys	Val	Phe	Asp	Arg	
		35					40					45				
Gln	Gly	Val	Ser	Arg	Ala	Ile	Val	Lys	Gln	Ala	Leu	Pro	Tyr	Val	Arg	
	50					55					60					
Cys	Val	Gly	Glu	Ser	Trp	Pro	Leu	Thr	Leu	Asp	Arg	Ala	Arg	Leu	Glu	
65					70					75					80	
Ala	Gln	Thr	Leu	Val	Ala	His	Tyr	Gln	His	Ser	Pro	Gln	His	Thr	Val	
			85						90					95		
Lys	Ile	His	His	Phe	Asp	Pro	Glu	Leu	Ala	Val	Met	Val	Met	Glu	Asp	
		100						105					110			
Leu	Ser	Asp	His	Arg	Ile	Trp	Arg	Gly	Glu	Leu	Ile	Ala	Asn	Val	Tyr	
	115					120						125				
Tyr	Pro	Gln	Ala	Ala	Arg	Gln	Leu	Gly	Asp	Tyr	Leu	Ala	Gln	Val	Leu	
	130					135					140					
Phe	His	Thr	Ser	Asp	Phe	Tyr	Leu	His	Pro	His	Glu	Lys	Lys	Ala	Gln	
145					150					155					160	
Val	Ala	Gln	Phe	Ile	Asn	Pro	Ala	Met	Cys	Glu	Ile	Thr	Glu	Asp	Leu	
			165						170					175		
Phe	Phe	Asn	Asp	Pro	Tyr	Gln	Ile	His	Glu	Arg	Asn	Asn	Tyr	Pro	Ala	
		180						185					190			
Glu	Leu	Glu	Ala	Asp	Val	Ala	Ala	Leu	Arg	Asp	Asp	Ala	Gln	Leu	Lys	
	195					200						205				
Leu	Ala	Val	Ala	Ala	Leu	Lys	His	Arg	Phe	Phe	Ala	His	Ala	Glu	Ala	
	210				215					220						
Leu	Leu	His	Gly	Asp	Ile	His	Ser	Gly	Ser	Ile	Phe	Val	Ala	Glu	Gly	
225					230					235					240	
Ser	Leu	Lys	Ala	Ile	Asp	Ala	Glu	Phe	Gly	Tyr	Phe	Gly	Pro	Ile	Gly	
			245						250					255		
Phe	Asp	Ile	Gly	Thr	Ala	Ile	Gly	Asn	Leu	Leu	Leu	Asn	Tyr	Cys	Gly	
			260					265					270			

PF 53790

61

Leu Pro Gly Gln Leu Gly Ile Arg Asp Ala Ala Ala Arg Glu Gln  
 275 280 285  
 Arg Leu Asn Asp Ile His Gln Leu Trp Thr Thr Phe Ala Glu Arg Phe  
 290 295 300  
 Gln Ala Leu Ala Ala Glu Lys Thr Arg Asp Ala Ala Leu Ala Tyr Pro  
 305 310 315 320  
 Gly Tyr Ala Ser Ala Phe Leu Lys Lys Val Trp Ala Asp Ala Val Gly  
 325 330 335  
 Phe Cys Gly Ser Glu Leu Ile Arg Arg Ser Val Gly Leu Ser His Val  
 340 345 350  
 Ala Asp Ile Asp Thr Ile Gln Asp Asp Ala Met Arg His Glu Cys Leu  
 355 360 365  
 Arg His Ala Ile Thr Leu Gly Arg Ala Leu Ile Val Leu Ala Glu Arg  
 370 375 380  
 Ile Asp Ser Val Asp Glu Leu Leu Ala Arg Val Arg Gln Tyr Ser  
 385 390 395

&lt;210&gt; 41

&lt;211&gt; 1140

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1137)

&lt;223&gt; coding for alcohol dehydrogenase

&lt;400&gt; 41

atg tct acc acc gga cag att att cga tgc aaa gct gct gtg gca tgg 48  
 Met Ser Thr Thr Gly Gln Ile Ile Arg Cys Lys Ala Ala Val Ala Trp  
 1 5 10 15  
 gaa gcc gga aag cca ctg gtg atc gag gaa gtg gag gtt gct cca ccg 96  
 Glu Ala Gly Lys Pro Leu Val Ile Glu Glu Val Glu Val Ala Pro Pro  
 20 25 30  
 cag aaa cac gaa gtt cgt atc aag att ctc ttc act tct ctc tgt cac 144  
 Gln Lys His Glu Val Arg Ile Lys Ile Leu Phe Thr Ser Leu Cys His  
 35 40 45  
 acc gat gtt tac ttc tgg gaa gct aag gga caa aca ccg ttg ttt cca 192  
 Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Leu Phe Pro  
 50 55 60  
 cgt atc ttc ggc cat gaa gct gga ggg att gtt gag agt gtt gga gaa 240  
 Arg Ile Phe Gly His Glu Ala Gly Gly Ile Val Glu Ser Val Gly Glu  
 65 70 75 80  
 gga gtg act gat ctt cag cca gga gat cat gtg ttg ccg atc ttt acc 288  
 Gly Val Thr Asp Leu Gln Pro Gly Asp His Val Leu Pro Ile Phe Thr  
 85 90 95  
 gga gaa tgt gga gat tgt cgt cat tgc cag tcg gag gaa tca aac atg 336  
 Gly Glu Cys Gly Asp Cys Arg His Cys Gln Ser Glu Glu Ser Asn Met  
 100 105 110  
 tgt gat ctt ctc agg atc aac aca gag cga gga ggt atg att cac gat 384  
 Cys Asp Leu Leu Arg Ile Asn Thr Glu Arg Gly Gly Met Ile His Asp  
 115 120 125

PF 53790

62

ggt gaa tct aga ttc tcc att aat ggc aaa cca atc tac cat ttc ctt	432
Gly Glu Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Tyr His Phe Leu	
130 135 140	
ggg acg tcc acg ttc agt gag tac act gtg gtt cac tct ggt cag gtc	480
Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Val His Ser Gly Gln Val	
145 150 155 160	
gct aag atc aat ccg gat gct cct ctt gac aag gtc tgt att gtc agt	528
Ala Lys Ile Asn Pro Asp Ala Pro Leu Asp Lys Val Cys Ile Val Ser	
165 170 175	
tgt ggt ttg tct act ggg tta gga gca act ttg aat gtg gct aaa ccc	576
Cys Gly Leu Ser Thr Gly Leu Gly Ala Thr Leu Asn Val Ala Lys Pro	
180 185 190	
aag aaa ggt caa agt gtt gcc att ttt ggt ctt ggt gct gtt ggt tta	624
Lys Lys Gly Gln Ser Val Ala Ile Phe Gly Leu Gly Ala Val Gly Leu	
195 200 205	
ggc gct gca gaa ggt gct aga atc gct ggt gct tct agg atc atc ggt	672
Gly Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly	
210 215 220	
gtt gat ttt aac tct aaa aga ttc gac caa gct aag gaa ttc ggt gtg	720
Val Asp Phe Asn Ser Lys Arg Phe Asp Gln Ala Lys Glu Phe Gly Val	
225 230 235 240	
acc gag tgt gtg aac ccg aaa gac cat gac aag cca att caa cag gtg	768
Thr Glu Cys Val Asn Pro Lys Asp His Asp Lys Pro Ile Gln Gln Val	
245 250 255	
atc gct gag atg acg gat ggt ggg gtg gac agg agt gtg gaa tgc acc	816
Ile Ala Glu Met Thr Asp Gly Gly Val Asp Arg Ser Val Glu Cys Thr	
260 265 270	
gga agc gtt cag gcc atg att caa gca ttt gaa tgt gtc cac gat ggc	864
Gly Ser Val Gln Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly	
275 280 285	
tgg ggt gtt gca gtg ctg gtg ggt gtg cca agc aaa gac gat gcc ttc	912
Trp Gly Val Ala Val Leu Val Gly Val Pro Ser Lys Asp Asp Ala Phe	
290 295 300	
aag act cat ccg atg aat ttc ttg aat gag agg act ctt aag ggt act	960
Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr	
305 310 315 320	
ttc ttc ggg aac tac aaa ccc aaa act gac att ccc ggg gtt gtg gaa	1008
Phe Phe Gly Asn Tyr Lys Pro Lys Thr Asp Ile Pro Gly Val Val Glu	
325 330 335	
aag tac atg aac aag gag ctg gag ctt gag aaa ttc atc act cac aca	1056
Lys Tyr Met Asn Lys Glu Leu Glu Leu Glu Lys Phe Ile Thr His Thr	
340 345 350	
gtg cca ttc tcg gaa atc aac aag gcc ttt gat tac atg ctg aag gga	1104
Val Pro Phe Ser Glu Ile Asn Lys Ala Phe Asp Tyr Met Leu Lys Gly	
355 360 365	
gag agt att cgt tgc atc atc acc atg ggt gct tga	1140
Glu Ser Ile Arg Cys Ile Ile Thr Met Gly Ala	
370 375	

&lt;210&gt; 42

&lt;211&gt; 379

PF 53790

63

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 42

```

Met Ser Thr Thr Gly Gln Ile Ile Arg Cys Lys Ala Ala Val Ala Trp
 1           5           10           15
Glu Ala Gly Lys Pro Leu Val Ile Glu Glu Val Glu Val Ala Pro Pro
 20           25           30
Gln Lys His Glu Val Arg Ile Lys Ile Leu Phe Thr Ser Leu Cys His
 35           40           45
Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Leu Phe Pro
 50           55           60
Arg Ile Phe Gly His Glu Ala Gly Gly Ile Val Glu Ser Val Gly Glu
 65           70           75           80
Gly Val Thr Asp Leu Gln Pro Gly Asp His Val Leu Pro Ile Phe Thr
 85           90           95
Gly Glu Cys Gly Asp Cys Arg His Cys Gln Ser Glu Glu Ser Asn Met
100          105          110
Cys Asp Leu Leu Arg Ile Asn Thr Glu Arg Gly Gly Met Ile His Asp
115          120          125
Gly Glu Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Tyr His Phe Leu
130          135          140
Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Val His Ser Gly Gln Val
145          150          155          160
Ala Lys Ile Asn Pro Asp Ala Pro Leu Asp Lys Val Cys Ile Val Ser
165          170          175
Cys Gly Leu Ser Thr Gly Leu Gly Ala Thr Leu Asn Val Ala Lys Pro
180          185          190
Lys Lys Gly Gln Ser Val Ala Ile Phe Gly Leu Gly Ala Val Gly Leu
195          200          205
Gly Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly
210          215          220
Val Asp Phe Asn Ser Lys Arg Phe Asp Gln Ala Lys Glu Phe Gly Val
225          230          235          240
Thr Glu Cys Val Asn Pro Lys Asp His Asp Lys Pro Ile Gln Gln Val
245          250          255
Ile Ala Glu Met Thr Asp Gly Gly Val Asp Arg Ser Val Glu Cys Thr
260          265          270
Gly Ser Val Gln Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly
275          280          285
Trp Gly Val Ala Val Leu Val Gly Val Pro Ser Lys Asp Asp Ala Phe
290          295          300
Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr
305          310          315          320
Phe Phe Gly Asn Tyr Lys Pro Lys Thr Asp Ile Pro Gly Val Val Glu
325          330          335
Lys Tyr Met Asn Lys Glu Leu Glu Leu Glu Lys Phe Ile Thr His Thr
340          345          350

```

PF 53790

64

Val Pro Phe Ser Glu Ile Asn Lys Ala Phe Asp Tyr Met Leu Lys Gly  
 355 360 365  
 Glu Ser Ile Arg Cys Ile Ile Thr Met Gly Ala  
 370 375

&lt;210&gt; 43

&lt;211&gt; 1140

&lt;212&gt; DNA

&lt;213&gt; Hordeum vulgare

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1137)

&lt;223&gt; coding for alcohol dehydrogenase

&lt;400&gt; 43

atg gcg acg gcc ggc aag gtg atc aag tgc aaa gcc gcg gtg gcg tgg	48
Met Ala Thr Ala Gly Lys Val Ile Lys Cys Lys Ala Ala Val Ala Trp	
1 5 10 15	
gag gcc ggg aag ccg ctg acc atg gag gag gtg gag gtg gcg ccg ccg	96
Glu Ala Gly Lys Pro Leu Thr Met Glu Glu Val Glu Val Ala Pro Pro	
20 25 30	
cag gcc atg gag gtg cgc gtc aag atc ctc ttc acc tcc ctc tgc cac	144
Gln Ala Met Glu Val Arg Val Lys Ile Leu Phe Thr Ser Leu Cys His	
35 40 45	
acc gac gtc tac ttc tgg gag gcc aag ggg cag acc ccc atg ttc cct	192
Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Met Phe Pro	
50 55 60	
cgg atc ttc ggc cat gaa gct gga ggc ata gtg gag agt gtt gga gag	240
Arg Ile Phe Gly His Glu Ala Gly Gly Ile Val Glu Ser Val Gly Glu	
65 70 75 80	
ggc gtg act gat gtt gcc cct ggt gac cac gtc ctc cct gtg ttc act	288
Gly Val Thr Asp Val Ala Pro Gly Asp His Val Leu Pro Val Phe Thr	
85 90 95	
ggg gag tgt aag gaa tgc cca cat tgc aag tct gcg gag agc aac atg	336
Gly Glu Cys Lys Glu Cys Pro His Cys Lys Ser Ala Glu Ser Asn Met	
100 105 110	
tgt gat ctg ctc agg atc aac acc gac aga ggt gtg atg atc ggg gat	384
Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Ile Gly Asp	
115 120 125	
ggc aag tcg cgc ttc tct att ggc ggc aag ccg att tac cat ttc gta	432
Gly Lys Ser Arg Phe Ser Ile Gly Gly Lys Pro Ile Tyr His Phe Val	
130 135 140	
ggg act tcc acc ttc agt gag tac act gtc atg cat gtc ggt tgt gtt	480
Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Met His Val Gly Cys Val	
145 150 155 160	
gcc aag atc aac cct gag gct ccc ctt gat aaa gtc tgt gtt ctt agc	528
Ala Lys Ile Asn Pro Glu Ala Pro Leu Asp Lys Val Cys Val Leu Ser	
165 170 175	
tgt ggt att tgc act ggt ctt ggc gcg tca att aat gtt gca aaa cca	576
Cys Gly Ile Cys Thr Gly Leu Gly Ala Ser Ile Asn Val Ala Lys Pro	
180 185 190	



PF 53790

65

```

cca aag ggt tcc aca gtg gcg ata ttt ggg cta gga gct gtt ggc ctt 624
Pro Lys Gly Ser Thr Val Ala Ile Phe Gly Leu Gly Ala Val Gly Leu
      195                200                205

gct gct gca gaa ggt gca agg att gca ggt gca tca agg atc att ggt 672
Ala Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly
      210                215                220

gtt gac ctg aac gcc agc aga ttt gaa gag gct agg aag ttt ggc tgc 720
Val Asp Leu Asn Ala Ser Arg Phe Glu Glu Ala Arg Lys Phe Gly Cys
      225                230                235                240

acg gaa ttt gtg aac ccg aaa gat cac acc aag cca gtt cag cag gtg 768
Thr Glu Phe Val Asn Pro Lys Asp His Thr Lys Pro Val Gln Gln Val
      245                250                255

ctc gct gac atg aca aat ggc gga gtt gac cgc agt gtt gag tgc act 816
Leu Ala Asp Met Thr Asn Gly Gly Val Asp Arg Ser Val Glu Cys Thr
      260                265                270

ggc aac gtc aat gct atg ata caa gca ttt gaa tgt gtt cat gat ggc 864
Gly Asn Val Asn Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly
      275                280                285

tgg ggt gta gct gtg ctg gtg ggt gtg cca cac aag gac gct gaa ttc 912
Trp Gly Val Ala Val Leu Val Gly Val Pro His Lys Asp Ala Glu Phe
      290                295                300

aag acc cac ccg atg aac ttc ctg aat gag agg acc ctg aag ggc acc 960
Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr
      305                310                315                320

ttc ttc ggt aac ttc aag ccg cgc act gac ctg ccc aat gtc gtg gag 1008
Phe Phe Gly Asn Phe Lys Pro Arg Thr Asp Leu Pro Asn Val Val Glu
      325                330                335

atg tac atg aag aag gag ctg gag gtg gag aag ttc atc aca cac agc 1056
Met Tyr Met Lys Lys Glu Leu Glu Val Glu Lys Phe Ile Thr His Ser
      340                345                350

gtg ccg ttc tcg gag ata aac aag gcc ttc gac ctt atg gcg aag ggg 1104
Val Pro Phe Ser Glu Ile Asn Lys Ala Phe Asp Leu Met Ala Lys Gly
      355                360                365

gag ggc atc cgt tgc atc atc cgc atg gac aac tag 1140
Glu Gly Ile Arg Cys Ile Ile Arg Met Asp Asn
      370                375

```

&lt;210&gt; 44

&lt;211&gt; 379

&lt;212&gt; PRT

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 44

```

Met Ala Thr Ala Gly Lys Val Ile Lys Cys Lys Ala Ala Val Ala Trp
  1                5                10                15

Glu Ala Gly Lys Pro Leu Thr Met Glu Glu Val Glu Val Ala Pro Pro
  20                25                30

Gln Ala Met Glu Val Arg Val Lys Ile Leu Phe Thr Ser Leu Cys His
  35                40                45

Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Met Phe Pro
  50                55                60

```

PF 53790

66

Arg Ile Phe Gly His Glu Ala Gly Gly Ile Val Glu Ser Val Gly Glu  
 65 70 75 80  
 Gly Val Thr Asp Val Ala Pro Gly Asp His Val Leu Pro Val Phe Thr  
 85 90 95  
 Gly Glu Cys Lys Glu Cys Pro His Cys Lys Ser Ala Glu Ser Asn Met  
 100 105 110  
 Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Ile Gly Asp  
 115 120 125  
 Gly Lys Ser Arg Phe Ser Ile Gly Gly Lys Pro Ile Tyr His Phe Val  
 130 135 140  
 Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Met His Val Gly Cys Val  
 145 150 155 160  
 Ala Lys Ile Asn Pro Glu Ala Pro Leu Asp Lys Val Cys Val Leu Ser  
 165 170 175  
 Cys Gly Ile Cys Thr Gly Leu Gly Ala Ser Ile Asn Val Ala Lys Pro  
 180 185 190  
 Pro Lys Gly Ser Thr Val Ala Ile Phe Gly Leu Gly Ala Val Gly Leu  
 195 200 205  
 Ala Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly  
 210 215 220  
 Val Asp Leu Asn Ala Ser Arg Phe Glu Glu Ala Arg Lys Phe Gly Cys  
 225 230 235 240  
 Thr Glu Phe Val Asn Pro Lys Asp His Thr Lys Pro Val Gln Gln Val  
 245 250 255  
 Leu Ala Asp Met Thr Asn Gly Gly Val Asp Arg Ser Val Glu Cys Thr  
 260 265 270  
 Gly Asn Val Asn Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly  
 275 280 285  
 Trp Gly Val Ala Val Leu Val Gly Val Pro His Lys Asp Ala Glu Phe  
 290 295 300  
 Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr  
 305 310 315 320  
 Phe Phe Gly Asn Phe Lys Pro Arg Thr Asp Leu Pro Asn Val Val Glu  
 325 330 335  
 Met Tyr Met Lys Lys Glu Leu Glu Val Glu Lys Phe Ile Thr His Ser  
 340 345 350  
 Val Pro Phe Ser Glu Ile Asn Lys Ala Phe Asp Leu Met Ala Lys Gly  
 355 360 365  
 Glu Gly Ile Arg Cys Ile Ile Arg Met Asp Asn  
 370 375

&lt;210&gt; 45

&lt;211&gt; 1140

&lt;212&gt; DNA

&lt;213&gt; Oryza sativa

&lt;220&gt;

&lt;221&gt; CDS

PF 53790

67

&lt;222&gt; (1)..(1137)

&lt;223&gt; coding for alcohol dehydrogenase

&lt;400&gt; 45

atg gcg acc gca ggg aag gtg atc aag tgc aaa gcg gcg gtg gca tgg	48
Met Ala Thr Ala Gly Lys Val Ile Lys Cys Lys Ala Ala Val Ala Trp	
1 5 10 15	
gag gcc gcg aag ccg ctg gtg atc gag gag gtg gag gtg gcg ccg ccg	96
Glu Ala Ala Lys Pro Leu Val Ile Glu Glu Val Glu Val Ala Pro Pro	
20 25 30	
cag gcc atg gag gtg cgc gtc aag atc ctc ttc acc tcg ctc tgc cac	144
Gln Ala Met Glu Val Arg Val Lys Ile Leu Phe Thr Ser Leu Cys His	
35 40 45	
acc gac gtc tac ttc tgg gag gcc aag gga cag act ccc gtg ttc cct	192
Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Val Phe Pro	
50 55 60	
cgg atc ttc ggc cat gaa gct gga ggt att gtg gag agt gtt gga gag	240
Arg Ile Phe Gly His Glu Ala Gly Gly Ile Val Glu Ser Val Gly Glu	
65 70 75 80	
ggt gtg act gat ctt gcc cct ggt gac cat gtt ctc cct gtg ttc act	288
Gly Val Thr Asp Leu Ala Pro Gly Asp His Val Leu Pro Val Phe Thr	
85 90 95	
ggg gag tgc aag gag tgt gcc cac tgc aag tca gca gag agc aac atg	336
Gly Glu Cys Lys Glu Cys Ala His Cys Lys Ser Ala Glu Ser Asn Met	
100 105 110	
tgt gat ctg ctc agg atc aac act gac agg ggt gtg atg att ggt gat	384
Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Ile Gly Asp	
115 120 125	
ggc aaa tca cgc ttt tcc atc aac ggg aag ccc att tac cat ttc gtc	432
Gly Lys Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Tyr His Phe Val	
130 135 140	
ggg act tcg acc ttc agc gag tac act gtc atg cat gtt ggt tgc gtt	480
Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Met His Val Gly Cys Val	
145 150 155 160	
gcg aag atc aac ccg gca gct cca ctt gat aaa gtt tgc gtt ctt agc	528
Ala Lys Ile Asn Pro Ala Ala Pro Leu Asp Lys Val Cys Val Leu Ser	
165 170 175	
tgt ggt att tct act ggt ctt ggt gct aca atc aat gtg gca aag cca	576
Cys Gly Ile Ser Thr Gly Leu Gly Ala Thr Ile Asn Val Ala Lys Pro	
180 185 190	
cca aag ggt tcg acg gtg gcg ata ttt ggt cta gga gct gta ggc ctt	624
Pro Lys Gly Ser Thr Val Ala Ile Phe Gly Leu Gly Ala Val Gly Leu	
195 200 205	
gct gcc gca gaa ggt gca agg att gca gga gcg tca agg atc att ggc	672
Ala Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly	
210 215 220	
att gac ctg aac gcc aac aga ttt gaa gaa gct agg aaa ttt ggt tgc	720
Ile Asp Leu Asn Ala Asn Arg Phe Glu Glu Ala Arg Lys Phe Gly Cys	
225 230 235 240	
act gaa ttt gtg aac cca aag gac cat gac aag cca gtt cag cag gta	768
Thr Glu Phe Val Asn Pro Lys Asp His Asp Lys Pro Val Gln Gln Val	
245 250 255	

PF 53790

68

ctt gct gag atg acc aat ggc gga gtt gac cgc agc gtt gaa tgc act 816  
 Leu Ala Glu Met Thr Asn Gly Gly Val Asp Arg Ser Val Glu Cys Thr  
 260 265 270

ggc aac atc aac gcc atg atc caa gca ttt gaa tgt gtt cat gat ggc 864  
 Gly Asn Ile Asn Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly  
 275 280 285

tgg ggt gtt gct gtt ttg gtc ggc gtg cca cac aag gac gcc gag ttc 912  
 Trp Gly Val Ala Val Leu Val Gly Val Pro His Lys Asp Ala Glu Phe  
 290 295 300

aag acc cac ccg atg aac ttc ctg aac gag agg act ctc aag gga acc 960  
 Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr  
 305 310 315 320

ttc ttc ggc aac tac aag cca cgc acc gat ctg ccc aac gtc gtc gag 1008  
 Phe Phe Gly Asn Tyr Lys Pro Arg Thr Asp Leu Pro Asn Val Val Glu  
 325 330 335

ctc tac atg aag aag gag ctg gag gtg gag aag ttc atc aca cac agc 1056  
 Leu Tyr Met Lys Lys Glu Leu Glu Val Glu Lys Phe Ile Thr His Ser  
 340 345 350

gtg ccg ttc tcg gag atc aac acg gcg ttc gac ctg atg cac aag ggc 1104  
 Val Pro Phe Ser Glu Ile Asn Thr Ala Phe Asp Leu Met His Lys Gly  
 355 360 365

gag ggc atc cgc tgc atc atc cgc atg gag aac tga 1140  
 Glu Gly Ile Arg Cys Ile Ile Arg Met Glu Asn  
 370 375

<210> 46  
 <211> 379  
 <212> PRT  
 <213> *Oryza sativa*  
 <400> 46

Met Ala Thr Ala Gly Lys Val Ile Lys Cys Lys Ala Ala Val Ala Trp  
 1 5 10 15

Glu Ala Ala Lys Pro Leu Val Ile Glu Glu Val Glu Val Ala Pro Pro  
 20 25 30

Gln Ala Met Glu Val Arg Val Lys Ile Leu Phe Thr Ser Leu Cys His  
 35 40 45

Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Val Phe Pro  
 50 55 60

Arg Ile Phe Gly His Glu Ala Gly Gly Ile Val Glu Ser Val Gly Glu  
 65 70 75 80

Gly Val Thr Asp Leu Ala Pro Gly Asp His Val Leu Pro Val Phe Thr  
 85 90 95

Gly Glu Cys Lys Glu Cys Ala His Cys Lys Ser Ala Glu Ser Asn Met  
 100 105 110

Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Ile Gly Asp  
 115 120 125

Gly Lys Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Tyr His Phe Val  
 130 135 140

Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Met His Val Gly Cys Val  
 145 150 155 160

PF 53790

69

Ala Lys Ile Asn Pro Ala Ala Pro Leu Asp Lys Val Cys Val Leu Ser  
 165 170 175  
 Cys Gly Ile Ser Thr Gly Leu Gly Ala Thr Ile Asn Val Ala Lys Pro  
 180 185 190  
 Pro Lys Gly Ser Thr Val Ala Ile Phe Gly Leu Gly Ala Val Gly Leu  
 195 200 205  
 Ala Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly  
 210 215 220  
 Ile Asp Leu Asn Ala Asn Arg Phe Glu Glu Ala Arg Lys Phe Gly Cys  
 225 230 235 240  
 Thr Glu Phe Val Asn Pro Lys Asp His Asp Lys Pro Val Gln Gln Val  
 245 250 255  
 Leu Ala Glu Met Thr Asn Gly Gly Val Asp Arg Ser Val Glu Cys Thr  
 260 265 270  
 Gly Asn Ile Asn Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly  
 275 280 285  
 Trp Gly Val Ala Val Leu Val Gly Val Pro His Lys Asp Ala Glu Phe  
 290 295 300  
 Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr  
 305 310 315 320  
 Phe Phe Gly Asn Tyr Lys Pro Arg Thr Asp Leu Pro Asn Val Val Glu  
 325 330 335  
 Leu Tyr Met Lys Lys Glu Leu Glu Val Glu Lys Phe Ile Thr His Ser  
 340 345 350  
 Val Pro Phe Ser Glu Ile Asn Thr Ala Phe Asp Leu Met His Lys Gly  
 355 360 365  
 Glu Gly Ile Arg Cys Ile Ile Arg Met Glu Asn  
 370 375

&lt;210&gt; 47

&lt;211&gt; 1140

&lt;212&gt; DNA

&lt;213&gt; Zea mays

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1137)

&lt;223&gt; coding for alcohol dehydrogenase

&lt;400&gt; 47

atg gcg acc gcg ggg aag gtg atc aag tgc aaa gct gcg gtg gca tgg 48  
 Met Ala Thr Ala Gly Lys Val Ile Lys Cys Lys Ala Ala Val Ala Trp  
 1 5 10 15  
 gag gcc ggc aag cca ctg tcg atc gag gag gtg gag gta gcg cct ccg 96  
 Glu Ala Gly Lys Pro Leu Ser Ile Glu Glu Val Glu Val Ala Pro Pro  
 20 25 30  
 cag gcc atg gag gtg cgc gtc aag atc ctc ttc acc tcg ctc tgc cac 144  
 Gln Ala Met Glu Val Arg Val Lys Ile Leu Phe Thr Ser Leu Cys His  
 35 40 45

PF 53790

70

acc gac gtc tac ttc tgg gag gcc aag ggg cag act ccc gtg ttc cct	192
Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Val Phe Pro	
50 55 60	
cgg atc ttt ggc cat gag gct gga ggt atc ata gag agt gtt gga gag	240
Arg Ile Phe Gly His Glu Ala Gly Gly Ile Ile Glu Ser Val Gly Glu	
65 70 75 80	
ggt gtg act gac gta gct ccg ggc gac cat gtc ctt cct gtg ttc act	288
Gly Val Thr Asp Val Ala Pro Gly Asp His Val Leu Pro Val Phe Thr	
85 90 95	
ggg gag tgc aag gag tgc gcc cac tgc aag tgc gca gag agc aac atg	336
Gly Glu Cys Lys Glu Cys Ala His Cys Lys Ser Ala Glu Ser Asn Met	
100 105 110	
tgt gat ttg ctc agg atc aac act gac cgc ggt gtg atg att ggc gat	384
Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Ile Gly Asp	
115 120 125	
ggc aag tgc cgg ttt tca atc aat ggg aag cct atc tac cac ttt gtt	432
Gly Lys Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Tyr His Phe Val	
130 135 140	
ggg act tcc acc ttc agc gag tac acc gtc atg cat gtc ggt tgt gtt	480
Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Met His Val Gly Cys Val	
145 150 155 160	
gca aag atc aac cct cag gct ccc ctt gat aaa gtt tgc gtc ctt agc	528
Ala Lys Ile Asn Pro Gln Ala Pro Leu Asp Lys Val Cys Val Leu Ser	
165 170 175	
tgt ggt att tct act ggt ctt ggt gca tca att aat gtt gca aaa cct	576
Cys Gly Ile Ser Thr Gly Leu Gly Ala Ser Ile Asn Val Ala Lys Pro	
180 185 190	
ccg aag ggt tgc aca gtg gct gtt ttc ggt tta gga gcc gtt ggt ctt	624
Pro Lys Gly Ser Thr Val Ala Val Phe Gly Leu Gly Ala Val Gly Leu	
195 200 205	
gcc gct gca gaa ggt gca agg att gct gga gcg tca agg atc att ggt	672
Ala Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly	
210 215 220	
gtc gac ctg aac ccc agc aga ttc gaa gaa gct agg aag ttc ggt tgc	720
Val Asp Leu Asn Pro Ser Arg Phe Glu Glu Ala Arg Lys Phe Gly Cys	
225 230 235 240	
act gaa ttt gtg aac cca aaa gac cac aac aag ccg gtg cag gag gta	768
Thr Glu Phe Val Asn Pro Lys Asp His Asn Lys Pro Val Gln Glu Val	
245 250 255	
ctt gct gag atg acc aac gga ggg gtc gac cgc agc gtg gaa tgc act	816
Leu Ala Glu Met Thr Asn Gly Gly Val Asp Arg Ser Val Glu Cys Thr	
260 265 270	
ggc aac atc aat gct atg atc caa gct ttc gaa tgt gtt cat gat ggc	864
Gly Asn Ile Asn Ala Met Ile Gln Ala Phe Glu Cys Val His Asp Gly	
275 280 285	
tgg ggt gtt gcc gtg ctg gtg ggt gtg ccg cat aag gac gct gag ttc	912
Trp Gly Val Ala Val Leu Val Gly Val Pro His Lys Asp Ala Glu Phe	
290 295 300	
aag acc cac ccg atg aac ttc ctg aac gaa agg acc ctg aag ggg acc	960
Lys Thr His Pro Met Asn Phe Leu Asn Glu Arg Thr Leu Lys Gly Thr	
305 310 315 320	

PF 53790

71

```

ttc ttt ggc aac tat aag cca cgc act gat ctg cca aat gtg gtg gag 1008
Phe Phe Gly Asn Tyr Lys Pro Arg Thr Asp Leu Pro Asn Val Val Glu
      325                      330                      335

ctg tac atg aaa aag gag ctg gag gtg gag aag ttc atc acg cac agc 1056
Leu Tyr Met Lys Lys Glu Leu Glu Val Glu Lys Phe Ile Thr His Ser
      340                      345                      350

gtc ccg ttc gcg gag atc aac aag gcg ttc aac ctg atg gcc aag ggg 1104
Val Pro Phe Ala Glu Ile Asn Lys Ala Phe Asn Leu Met Ala Lys Gly
      355                      360                      365

gag ggc atc cgc tgc atc atc cgc atg gag aac tag 1140
Glu Gly Ile Arg Cys Ile Ile Arg Met Glu Asn
      370                      375

```

&lt;210&gt; 48

&lt;211&gt; 379

&lt;212&gt; PRT

&lt;213&gt; Zea mays

&lt;400&gt; 48

```

Met Ala Thr Ala Gly Lys Val Ile Lys Cys Lys Ala Ala Val Ala Trp
  1          5          10          15

Glu Ala Gly Lys Pro Leu Ser Ile Glu Glu Val Glu Val Ala Pro Pro
      20          25          30

Gln Ala Met Glu Val Arg Val Lys Ile Leu Phe Thr Ser Leu Cys His
      35          40          45

Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Thr Pro Val Phe Pro
      50          55          60

Arg Ile Phe Gly His Glu Ala Gly Gly Ile Ile Glu Ser Val Gly Glu
      65          70          75          80

Gly Val Thr Asp Val Ala Pro Gly Asp His Val Leu Pro Val Phe Thr
      85          90          95

Gly Glu Cys Lys Glu Cys Ala His Cys Lys Ser Ala Glu Ser Asn Met
      100          105          110

Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Ile Gly Asp
      115          120          125

Gly Lys Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Tyr His Phe Val
      130          135          140

Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Met His Val Gly Cys Val
      145          150          155          160

Ala Lys Ile Asn Pro Gln Ala Pro Leu Asp Lys Val Cys Val Leu Ser
      165          170          175

Cys Gly Ile Ser Thr Gly Leu Gly Ala Ser Ile Asn Val Ala Lys Pro
      180          185          190

Pro Lys Gly Ser Thr Val Ala Val Phe Gly Leu Gly Ala Val Gly Leu
      195          200          205

Ala Ala Ala Glu Gly Ala Arg Ile Ala Gly Ala Ser Arg Ile Ile Gly
      210          215          220

Val Asp Leu Asn Pro Ser Arg Phe Glu Glu Ala Arg Lys Phe Gly Cys
      225          230          235          240

```

PF 53790

72

Thr	Glu	Phe	Val	Asn	Pro	Lys	Asp	His	Asn	Lys	Pro	Val	Gln	Glu	Val
				245					250					255	
Leu	Ala	Glu	Met	Thr	Asn	Gly	Gly	Val	Asp	Arg	Ser	Val	Glu	Cys	Thr
			260					265					270		
Gly	Asn	Ile	Asn	Ala	Met	Ile	Gln	Ala	Phe	Glu	Cys	Val	His	Asp	Gly
		275					280					285			
Trp	Gly	Val	Ala	Val	Leu	Val	Gly	Val	Pro	His	Lys	Asp	Ala	Glu	Phe
	290					295					300				
Lys	Thr	His	Pro	Met	Asn	Phe	Leu	Asn	Glu	Arg	Thr	Leu	Lys	Gly	Thr
305					310					315					320
Phe	Phe	Gly	Asn	Tyr	Lys	Pro	Arg	Thr	Asp	Leu	Pro	Asn	Val	Val	Glu
			325						330					335	
Leu	Tyr	Met	Lys	Lys	Glu	Leu	Glu	Val	Glu	Lys	Phe	Ile	Thr	His	Ser
		340						345					350		
Val	Pro	Phe	Ala	Glu	Ile	Asn	Lys	Ala	Phe	Asn	Leu	Met	Ala	Lys	Gly
		355					360					365			
Glu	Gly	Ile	Arg	Cys	Ile	Ile	Arg	Met	Glu	Asn					
	370					375									

&lt;210&gt; 49

&lt;211&gt; 505

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; Description of the artificial sequence: coding for sense RNA-fragment of E.coli codA gene

&lt;400&gt; 49

```

aagcttggt aacagtgtcg aataacgctt tacaaacaat tattaacgcc cggttaccag 60
gcgaagagg gctgtggcag attcatctgc aggacggaaa aatcagcgcc attgatgcgc 120
aatccggcgt gatgcccata actgaaaaca gcctggatgc cgaacaagggt ttagttatac 180
cgccggtttgt ggagccacat attcacctgg acaccacgca aaccgccgga caaccgaact 240
ggaatcagtc cggcacgctg tttgaaggca ttgaacgctg ggccgagcgc aaagcgttat 300
taacccatga cgatgtgaaa caacgcgcat ggcaaacgct gaaatggcag attgccaacg 360
gcattcagca tgtgcgtacc catgtcgatg ttccggatgc aacgctaact gcgctgaaag 420
caatgctgga agtgaagcag gaagtcgcgc cgtggattga tctgcaaadc gtcgccttcc 480
ctcaggaagg gattttgtcg tcgac                                     505

```

&lt;210&gt; 50

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; Description of the artificial sequence: oligonucleotide primer

&lt;400&gt; 50

cgtgaatacg gcgtggagtc g

21

&lt;210&gt; 51

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence



PF 53790

73

&lt;220&gt;

<223> Description of the artificial sequence:  
oligonucleotide primer

&lt;400&gt; 51

cggcaggata atcagggttg

20

&lt;210&gt; 52

&lt;211&gt; 505

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence: coding for  
antisense RNA-fragment of E.coli codA gene

&lt;400&gt; 52

gaattcggct aacagtgtcg aataacgctt tacaacaat tattaacgcc cggttaccag 60  
gcgaagaggg gctgtggcag attcatctgc aggacgaaa aatcagcgcc attgatgcgc 120  
aatccggcgt gatgccata actgaaaaca gcctggatgc cgaacaagggt ttagttatac 180  
cgccgtttgt ggagccacat attcacctgg acaccacgca aaccgccgga caaccgaact 240  
ggaatcagtc cggcacgctg tttgaaggca ttgaacgctg ggccgagcgc aaagcggtat 300  
taacccatga cgatgtgaaa caacgcgcgt ggcaaacgct gaaatggcag attgccaacg 360  
gcattcagca tgtgcgtacc catgtcgatg tttcggatgc aacgctaact gcgctgaaag 420  
caatgctgga agtgaagcag gaagtgcgcg cgtggattga tctgcaaata gtcgccttcc 480  
ctcaggaagg gattttgtcg gatcc 505

&lt;210&gt; 53

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence:  
oligonucleotide primer

&lt;400&gt; 53

gtcaacgtaa ccaaccctgc

20

&lt;210&gt; 54

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence:  
oligonucleotide primer

&lt;400&gt; 54

ggatccgaca aaatccccttc ctgagg

26

&lt;210&gt; 55

&lt;211&gt; 5674

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence: vector  
construct pBluKS-nitP-STLS1-35S-T

&lt;400&gt; 55

ccagcttttg ttcccttttag tgagggttaa tttcgagctt ggcgtaatca tggatcatagc 60  
tgtttcctgt gtgaaattgt tatccgctca caattccaca caacatacga gccggaagca 120

PF 53790

74

taaagtgtaa	agcctggggg	gcctaagtga	tgagctaact	cacattaatt	gcgttgcgct	180
cactgccccg	tttccagtcg	ggaaacctgt	cgtgccagct	gcattaatga	atcggccaac	240
gcgcggggag	aggcggtttg	cgtattgggc	gotcttccgc	ttctcgcctc	actgactcgc	300
tgcgctcggg	cgttcggctg	cggcgagcgg	tatcagctca	ctcaaaggcg	gtaatacggg	360
tatccacaga	atcaggggat	aacgcaggaa	agaacatgtg	agcaaaaggc	cagcaaaagg	420
ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	taggctccgc	ccccctgacg	480
agcatcacaa	aaatcgacgc	tcaagtcaga	ggtggcgaaa	cccgcagga	ctataaagat	540
accaggcggt	ttccccctga	agctccctcg	tgcgctctcc	tggtccgacc	ctgccgctta	600
ccggatacct	gtccgccttt	ctcccttcgg	gaagcgtggc	gctttctcat	agctcacgct	660
gtaggtatct	cagttcgggtg	taggtcgttc	gtccaaagct	gggctgtgtg	cacgaacccc	720
ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtcc	aaccgggtaa	780
gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	gcgaggtatg	840
taggcgggtg	tacagagttc	ttgaagtggg	ggcctaacta	cggctacact	agaaggacag	900
tatttggtat	ctgcgctctg	ctgaagccag	ttaccttcgg	aaaaagaggt	ggtagctctt	960
gatccggcaa	acaaaccacc	gctggtagcg	gtgggttttt	tgtttgcaag	cagcagatta	1020
cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	1080
agtggaaacg	aaactcacgt	taagggattt	tggtcatgag	attatcaaaa	aggatcttca	1140
cctagatcct	tttaaattaa	aatgaagtt	ttaaatcaat	ctaaagtata	tatgagtaaa	1200
cttggtctga	cagttaccaaa	tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	1260
ttcgttcac	catagttgcc	tgactccccg	tcgtgtagat	aactacgata	cgggaggggct	1320
taccatctgg	ccccagtgtc	gcaatgatac	cgcgagaccc	acgtcacccg	gctccagatt	1380
tatcagcaat	aaaccagcca	gccggaagg	ccgagcgcag	aagtggctct	gcaactttat	1440
ccgcctccat	ccagtctatt	aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	1500
atagtttgcg	caacgttggt	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	1560
gtatggcttc	attcagctcc	ggttcccaac	gatcaaggcg	agttacatga	tccccatgt	1620
tgtgcaaaaa	agcgggttagc	tccttcgggtc	ctccgatcgt	tgtcagaagt	aagttggccg	1680
cagtgttatc	actcatgggt	atggcagcac	tgcataattc	tcttactgtc	atgccatccg	1740
taagatgctt	ttctgtgact	ggtgagtact	caaccaagtc	attctgagaa	tagtgtatgc	1800
ggcgaccgag	ttgctcttgc	ccggcgtaaa	tacgggataa	taccgcgcca	catagcagaa	1860
ctttaaaagt	gctcatcatt	ggaaaacgtt	cttcggggcg	aaaactctca	aggatcttac	1920
cgctgttgag	atccagttcg	atgtaaccca	ctcgtgcacc	caactgatct	tcagcatctt	1980
ttactttcac	cagcggtttct	gggtgagcaa	aaacaggaag	gcaaaatgcc	gcaaaaaagg	2040
gaataagggc	gacacggaaa	tggtgaatac	tcatactctt	cctttttcaa	tattattgaa	2100
gcatttatca	gggttattgt	ctcatgagcg	gatacatatt	tgaaatgtatt	tagaaaaata	2160
aacaaatagg	ggttccgcgc	acatttcccc	gaaaagtgcc	acctgacgcg	ccctgtagcg	2220
gcgcattaag	cgcggcgggg	gtggtgggta	cgcgcagcgt	gaccgctaca	cttgccagcg	2280
ccctagcgcc	cgtctcttcc	gcttcttctc	cttcccttct	cgccacgttc	gocgggttcc	2340
cccgtaaac	tctaaatcgg	gggctccctt	taggggtccg	atttagtgct	ttacggcacc	2400
tcgaccccaa	aaaacttgat	taggggtgatg	gttcacgtag	tgggccatcg	ccctgataga	2460
cgggtttttcg	ccctttgacg	ttggagtcca	cgttctttta	tagtggaactc	ttgttccaaa	2520
ctggaacaac	actcaaccct	atctcgggtct	attcttttga	tttataaggg	attttgccga	2580
tttcggccta	ttggttaaaa	aatgagctga	tttaacaaaa	atttaacgcg	aattttaaca	2640
aaatattaac	gcttacaatt	tccattcgcc	attcaggctg	cgcaactgtt	gggaaggggcg	2700
atcgggtgcg	gcctcttcgc	tattacgcca	gctggcgaaa	gggggatgtg	ctgcaaggcg	2760
attaagttgg	gtaacgccag	ggttttccca	gtcacgacgt	tgtaaaacga	cggccagtg	2820
attgtaatac	gactcactat	agggcggaatt	ggagctcgtc	gagaccagat	gttttacact	2880
tgaccgtaaa	tgagcaccgc	aagaaaccgc	tcacattcat	ttcgaagggtg	gagaaagcgg	2940
aagatgactc	aaacaagtga	tcggttggtg	ttcgtcagtt	catgtcactc	ctatgaagg	3000
gtcaagttca	aaatgttatg	ttgagtttca	aacttttatg	ctaaactttt	tttctttatt	3060
ttcgttaata	atggaagaga	accaattctc	ttgtatctaa	agattatcca	tctatcatcc	3120
aatttgagtg	ttcaattctg	gatgttggtg	tacctacat	tctacaacca	tgtagccaat	3180
tattatgaat	ctggctttga	tttcagttgt	gttcttttct	tttttttctt	tgcatatttg	3240
catttagaat	gtttaataat	taagttactg	tatttcacac	tacattagtt	ccaagaatat	3300
acataatta	atttattttt	cttaaaaaatg	ttttggaatg	actaatattg	acaacgaaaa	3360
tagaagctat	gctaaaccat	tacgtatatg	tgacttcaca	tggtgtgtgt	ttacattccc	3420
tatatatatg	gatggctgtc	acaatcagaa	acgtgatcga	aaaaagacaa	acagtgtttg	3480

PF 53790

75

```

cataaaaaaga ctatttcggtt tcattgacaa tttgtgttta tttgtaaaga aaagtggcaa 3540
agtggaaattt gagttcctgc aagtaagaaa gatgaaataa aagacttgag tgtgtgtttt 3600
tttcttttat ctgaaagctg caatgaaata ttcctacca gcccgtttga ttattaattg 3660
gggtttgggtt ttcttgatgc gaactaattg gttatataag aaactataca atccatgta 3720
attcaaaaat tttgatttct cttgtaggaa tatgatttac tatatgagac tttcttttcg 3780
ccaataatag taaatccaaa gatatttgac cggaccaaaa cacattgato ttttttttag 3840
tttatttaat ccagtttctc tgagataatt cattaaggaa aacttagtat taacccatcc 3900
taagattaaa taggagccaa actcacattt caaatattaa ataacataaa atggatttaa 3960
aaaaatctata cgtcaaattt tatttatgac atttcttatt taaatttata tttaatgaaa 4020
tacagctaag acaaaccaaa aaaaaaatatc tttctaagtg gtccaaaaca tcaattccgt 4080
tcaatattat taggtagaat cgtacgacca aaaaaaggta ggtaataacg aattagaaac 4140
atatctataa catagtatat attattacct attatgagga atcaaaatgc atcaaatatg 4200
gatttaagga atccataaaa gaataaattc tacgggaaaa aaaatggaat aaattctttt 4260
aagttttttt tttgtttttt atttggtagt tctccatttt gttttatttc gtttggattt 4320
attgtgtcca aatactttgt aaaccaccgt tgtaattctt aaacgggggtt ttcacttctt 4380
ttttatattc agacataaag catcggctgg tttaatcaat caatagattt tatttttctt 4440
ctcaattatt agtaggtttg atgtgaactt tacaacaaaa acaaaaaacaa atcaatgcag 4500
agaaaagaaa ccacgtgggc tagtcccacc ttgtttcatt tccaccacag gttcgatctt 4560
cgttaccgtc tccaatagga aataaaacgt gaccacaaaa aaaaaacaaa aaaaagtcta 4620
tatattgctt ctctcaagtc tctgagtgtc atgaacaaaa gtaaaaaaca aagactcgac 4680
ctgcaggcat gcaagcttat cgtcgactac gtaagtttct gcttctacct ttgatata 4740
tataataatt atcattaatt agtagtaata taatatttca aatatttttt tcaaaataaa 4800
agaatgtagt atatagcaat tgcttttctg tagtttataa gtgtgtatat tttaatttat 4860
aacttttcta atatatgacc aaaatttgtt gatgtgcagg tatcaccgga tccatcgaat 4920
tcggtacgct gaaatcacca gtctctctct acaaactctat ctctctctat tttctccata 4980
aataatgtgt gagtagtttc ccgataaggg gaanttaggg ttcttatagg gtttcgctca 5040
tgtgttgagc atataagaaa cccttagtat gtatttgtat ttgtaaaata cttctatcaa 5100
taaaatttct aattcctaaa accaaaatcc agtactaaaa tccagatctc cttaaagtccc 5160
tatagatctt tgctcgtgaat ataaaccaga cacgagacga ctaaaccctgg agcccagacg 5220
ccgttcgaag ctgagaagta cgcttaggca ggaggccgtt agggaaaaaga tgctaaggca 5280
gggttggtta cgttgactcc ccgtaggtt tggtttaaat atgatgaagt ggacggaagg 5340
aaggaggaag acaagggaagg ataaggttgc aggccctgtg caaggtaaga agatggaaat 5400
ttgatagagg tacgtacta tacttatact atacgctaag ggaatgcttg tatttatacc 5460
ctataccccc taataacccc ttatcaattt aagaaataat ccgcataagc ccccgttaa 5520
aaattggtat cagagccatg aataggtcta tgacaaaaac tcaagaggat aaaacctcac 5580
caaaatacga aagagttctt aactctaaag ataaaagatc tttcaagatc aaaactagtt 5640
ccctcacacc ggtgacgggg atcgcgatgg gtac 5674

```

&lt;210&gt; 56

&lt;211&gt; 6046

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence: binary  
vector pSUN1

&lt;400&gt; 56

```

ttccatggac atacaaatgg acgaacggat aaaccttttc acgccctttt aaatatccga 60
ttattctaatt aaacgctctt ttctcttagg ttaccgcc aatatatcct gtcaaacact 120
gatagtttaa actgaaggcg ggaacgcaca atcagatcta gtaggaaaca gctatgacca 180
tgattacgcc aagcttgcat gcctgcaggt cgactctaga ctagtggatc cgatatcgcc 240
cgggctcgag gtaccgagct cgaattcact ggccgtcgtt ttacaacgac tcagctgctt 300
ggtataaatt gtcattagat tgtttttatg catagatgca ctcgaaatca gccaatttta 360
gacaagtatc aaacggatgt taattcagta cattaaagac gtccgcaatg tgttattaag 420
ttgtctaacg gtcattttgt ttacaccaca atatatcctg ccaccagcca gccaacagct 480
ccccgaccgg cagctcggca caaaatcacc acgcgttacc accacgccgg ccggccgcat 540
ggtgttgacc gtgttcgccg gcattgccga gttcagagct tccctaatac tcgaccgcac 600

```

PF 53790

76

```

ccggagcggg cgcgaggccg ccaaggcccg aggcgtgaag tttggccccc gccctaccct 660
caccgccgga cagatcgcg cgcgcccgga gctgatcgac caggaaggcc gcaccgtgaa 720
agaggcggct gcaactgcttg gcggtgcatcg ctcgaccctg taccgcgcac ttgagcgag 780
cgaggaagtg acgcccaccg agggccaggcg gcgcggtgcc ttccgtgagg acgcattgac 840
cgaggccgac gccctggcgg ccgcccagaa tgaacgcaa gaggaacaag catgaaaccg 900
caccaggacg gccaggacga accgtttttc attaccgaag agatcgaggc ggagatgatc 960
gcggccgggt acgtgttcga gccgcccgcg cactgtctca ccgtgcggct gcatgaaatc 1020
ctggccgggt tgtctgatgc caagctggcg gcctggccgg ccagcttggc cgctgaagaa 1080
accgagcgcc gccgtctaaa aaggtgatgt gtattttgagt aaaacagctt gcgtcatgcy 1140
gtcgctgcgt atatgatgc atgagtaaat aaacaaatc gcaaggggaa cgcataag 1200
ttatcgctgt acttaaccag aaaggcgggt caggcaagac gaccatcgca accatctag 1260
cccgccccct gcaactcgcc ggggcccgat ttctgttagt cgattccgat ccccgaggca 1320
gtgcccgcga ttgggcggcc gtgcgggaag atcaaccgct aaccgttgtc ggcacgacc 1380
gcccgcagat tgaccgcgac gtgaaggcca tcggccggcg cgacttcgta gtgatcgac 1440
gagcgcccca ggcggcggac ttggtgtgt ccgcgatcaa ggcagccgac ttctgtctga 1500
ttccggtgca gccaaagccct tacgacatat gggccaccgc cgacctggtg gagctggtta 1560
agcagcgcat tgaggtcacg gatggaaggc tacaagcggc ctttgcctg tcgcggcgca 1620
tcaaaggcac gcgcacgccc ggtgaggttg ccgaggcgct ggccgggtac gagctgccc 1680
ttcttgagtc ccgtatcacg cagcgctga gctaccagg cactgccgc gccggcaca 1740
ccgttcttga atcagaaccc gagggcgacg ctgcccgcga ggtccaggcg ctggccgctg 1800
aaattaaatc aaaactcatt tgagttaatg aggtaaagag aaaatgagca aaagcaca 1860
cacgctaagt gccggccgtc cgagcgacag gctgcaacgt tggccagcc 1920
ggcagacacg ccagccatga agcgggtcaa ctttcagttg ccggcgagg atcacacca 1980
gctgaagatg tacgcggtac gccaaaggca gaccattacc gagctgctat ctgaatacat 2040
cgcgagccta ccagagtaaa tgagcaaatg aataaatgag tagatgaatt ttagcggcta 2100
aaggaggcgg catggaaaat caagaacaac caggcaccga cgcctggaa tgccccatgt 2160
gtggaggaac gggcggttgg ccaggcgtaa gcgctgggt tgtctgccg ccctgcaatg 2220
gcactggaac ccccaagccc gaggaatcgg cgtgagcggt cgcaaacat ccggcccggt 2280
acaaatcggc gggcgcttgg gtgatgacct ggtggagaag ttgaaggccg cgcaggccgc 2340
ccagcgcaa cgcatcgagg cagaagcacg ccccggtgaa tcgtggcaag cggccgctga 2400
tcgaatccgc aaagaatccc ggcaaccgcc ggcagccggt gcgcgctga ttaggaagcc 2460
gcccagggc gacgagcaac cagatttttt cgttccgatg ctctatgacg tggccaccgc 2520
cgatagtcgc agcatcatgg acgtggccgt tttccgtctg tcgaagcgtg accgacgagc 2580
tggcgaggtg atccgctacg agcttccaga cgggcacgta gaggtttccg cagggcgggc 2640
cggcattggc agtgtgtggg attacgacct ggtactgatg gcggtttccc atctaaccga 2700
atccatgaac cgataccggg aagggaagg agacaagccc ggcgcgctgt tccgtccaca 2760
cgttgcggac gtactcaagt tctgccggcg agccgatggc ggaaagcaga aagacgacct 2820
ggtagaaacc tgcatcgggt taaacaccac gcacgttgcc atgcagcgta cgaagaaggc 2880
caagaacggc cgcttggta cggtatccga ggggtgaagc ttgattagcc gctacaagat 2940
cgtaaagagc gaaaccgggc ggccggagta catcgagatc gagctagctg attggatgta 3000
ccgcgagatc acagaaggca agaaccggga cgtgctgacg gttcaccocg attacttttt 3060
gatcgatccc ggcacggcc gttttctcta ccgctggca cgcgcgccc caggcaaggc 3120
agaagccaga tggttgttca agacgatcta ggaacgcagt ggcagcgccg gagagttaa 3180
gaagttctgt ttcaccgtgc gcaagctgat cgggtcaaat gacctgccg agtacgattt 3240
gaaggaggag gcggggcagg ctggcccgat cctagtcatg cgctaccgca acctgatcga 3300
gggcgaagca tccgcccgtt cctaattgtac ggagcagatg ctagggcaaa ttgccctagc 3360
aggggaaaaa ggtcgaaaaa gtotctttcc tgtggatagc acgtacattg ggaacccaaa 3420
gccgtacatt gggaaccgga acccgtaac tgggaaacca aagccgtaca ttgggaaccg 3480
gtcacacatg taagtactg atataaaaga gaaaaaaggc gatttttccg cctaaaactc 3540
tttaaaactt attaaaactc ttaaaacccg cctggcctgt gcataactgt ctggccagcg 3600
cacagccgaa gagctgcaa aagcgctac ccttcggctc ctgcgctccc tacgccccgc 3660
cgcttcgctg cggcctatcg cgcccgctgg ccgctcaaaa atggctggcc tacggccagg 3720
caatctacca gggcgcgac aagccgcgc gtcgccactc gaccgcccgc gccacatca 3780
aggcaccctg cctcgcgctg ttggtgatg accgtgaaaa cctctgacac atgcagctcc 3840
cggagacggt cacagcttgt ctgtaagcgg atgcgggag cagacaagcc cgtcaggcg 3900
cgtcagcggg tgttggcggg tgtcggggcg cagccatgac ccagtcacgt agcgatagcg 3960

```

PF 53790

77

```

gagtgtatac tggcttaact atgcggcatt agagcagatt gtactgagag tgcaccatat 4020
gcgggtgtgaa ataccgcaca gatgcgtaag gagaaaatac cgcatacaggc gctcttccgc 4080
ttcctcgtctc actgactcgc tgcgctcggg cggttcggctg cggcgagcgg tatcagctca 4140
ctcaaaggcg gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacatgtg 4200
agcaaaaggc cagcaaaagg ccaggaaccg taaaaaggcc gcgttgctgg cgtttttcca 4260
taggctccgc cccctgacg agcatcacaa aaatcgacgc tcaagtcaga ggtggcgaaa 4320
cccgacagga ctataaagat accaggcgtt tccccctgga agctccctcg tgcgctctcc 4380
tgttccgacc ctgccgctta ccggatacct gtccgccttt ctcccttcgg gaagcgtggc 4440
gctttctcat agctcacgct gtaggtatct cagttcgggtg taggtcgttc gctccaagct 4500
gggctgtgtg cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg 4560
tcttgagtc caccgggtaa gacacgactt atcgccactg gcagcagcca ctggtaacag 4620
gattagcaga gcgaggtatg taggcgggtg tacagagttc ttgaagtggg ggcctaacta 4680
cggctacact agaaggacag tatttgggat ctgcgctctg ctgaagccag ttaccttcgg 4740
aaaaagagtt ggtagctctt gatccggcaa acaaacacc gctggtagcg gtggtttttt 4800
tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt 4860
ttctacgggg tctgacgctc agtggaaacga aaactcactg taagggattt tggtcatgca 4920
tgatatatct cccaatttgt gtagggctta ttatgcacgc ttaaaaaataa taaaagcaga 4980
cttgacctga tagtttggtg gtgagcaatt atgtgcttag tgcactaac gcttgagtta 5040
agccgcggcg cgaagcggcg tcggcttgaa cgaatttcta gctagacatt atttgccgac 5100
taccttggtg atctcgcctt tcacgtagtg gacaaattct tccaactgat ctgcgcgcga 5160
ggccaagcga tcttcttctt gtccaagata agcctgtcta gcttcaagta tgacgggctg 5220
atactgggccc ggcaggcgcct ccattgccc aatgcgggga caacgtaagc actacatttc gctcatcgcc 5340
gcccggttact gcgctgtacc aaatgcggga taagggttca tttagcgcct caaatagatc 5400
agcccagtcg ggcggcaggt agagttcttc cgccgctgga cctaccaagg caacgctatg 5460
ctgttcagga accggatcaa agatagccag atcaatgtcg atcgtggctg gctcgaagat 5520
ttctcttgct tttgtcagca gctgccattc tccaaattgc agttcgcgct tagctggata 5580
acctgcaaga atgtcattgc cgtgcacaac aatggtgact tctacagcgc ggagaatctc 5640
acgccacgga atgatgtcgt aagtttccaa naggtcgttg atcaaagctc gccgcgttgt 5700
gtctcttcca ggggaagccg ccgtaaccag caaatcaata tcaactgtgtg gcttcaggcc 5760
ttcatcaagc cttacgggtc acaaatgtac ggccagcaac gtcgggttga gatggcgctc 5820
gccatccact gcggagccgt atagttgagt cgatacttcg gcgatcaccg cttcccccat 5880
gatgtttaac tttgttttag ggcgactgcc ctgctgcgta acatcggtgc tgctccataa 5940
catcaaacat cgaccacgg cgtaacgcgc ttgctgcttg gatgcccag gcatagactg 6000
taccacaaaa aaacagtcac aacaagccat gaaaaccgcc actgcg 6046

```

&lt;210&gt; 57

&lt;211&gt; 9838

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence: Transgenic  
expression vector for codA dsRNA pSUN1-codA-RNAi

&lt;400&gt; 57

```

cgaattcact ggccgtcgtt ttacaacgac tcagctgctt ggtaataatt gtcattagat 60
tgtttttatg catagatgca ctcgaaatca gccaatttta gacaagtatc aaacggatgt 120
taattcagta cattaaagac gtccgcaatg tggtattaaag ttgtctaaag gtcaatttgt 180
ttacaccaca atatatcctg ccaccagcca gccaacagct ccccgaccgg cagctcggca 240
caaaatcacc acgcgttacc accacgcccg ccggccgcat ggtgttgacc gtgttcgccg 300
gcattgccga gttcgagcgt tccctaatac tcgaccgcac ccggagcggg cgcgaggccg 360
ccaaggcccc aggcgtgaag tttggcccc gccctaccct cccccggca cagatcgcg 420
acgcccgcga gctgatcgac caggaaggcc gcaccgtgaa agaggcggct gactgcttg 480
gcgtgcacgc ctcgaccctg taccgcgcac ttgagcgcag cgagggaagt acgcccagc 540
aggccaggcg gcgcggtgcc ttccgtgagg acgcattgac cgaggccgac gccctggcg 600
ccgccgagaa tgaacgccaa gaggaacaag catgaaaccg caccaggacg gccaggacga 660
accgtttttc attaccgaag agatcgaggc ggagatgatc gcggccgggt acgtgttcga 720

```

PF 53790

78

```

gccgcccgcg cactgtctcaa ccgtgcggct gcatgaaatc ctggccgggt tgtctgatgc 780
caagctggcg gcctggccgg ccagcttggc cgctgaagaa accgagcgcc gccgtctaaa 840
aaggtgatgt gtatttgagt aaaacagctt gcgtcatgcg gtcgctgcgt atatgatgcg 900
atgagtaaat aaacaaatac gcaaggggaa cgcatgaagg ttatcgctgt acttaaccag 960
aaagggcggt caggcaagac gaccatcgca acccatctag ccgcgcgcct gcaactcgcc 1020
ggggccgatg ttctgttagt cgattccgat cccaggggca gtgccgcga ttggggcgcc 1080
gtgcgggaag atcaaccgct aaccgttgct ggcatcgacc gcccgacgat tgaccgcgac 1140
gtgaaggcca tcggccggcg cgacttcgta gtgatcgacg gagcgcccca ggccggcgac 1200
ttggctgtgt ccgcgatcaa ggcagccgac ttctgtgctga ttccggtgca gccaaagccct 1260
tacgacatat gggccaccgc cgacctggtg gagctggtta agcagcgcat tgaggtcacg 1320
gatggaaggc tacaagcggc ctttgtcgtg tcgcggcgca tcaaaggcac gcgcatcgcc 1380
ggtgagggtt ccgagcgct ggcgggttac gagctgcca ttcttgagtc ccgtatcacg 1440
cagcgcgtag gctaccagg cactgccgcc ccgggcacaa ccgttcttga atcagaacct 1500
gagggcgacg ctgccgcga ggtccaggcg ctggccgctg aaattaaatc aaaactcatt 1560
tgagttaatg aggtaaagag aaaatgagca aaagcacaaa cagcctaagt gccggccgct 1620
cgagcgcacg cagcagcaag gctgcaacgt tggccagcct ggcagacacg ccagccatga 1680
agcgggtcaa ctttcagttg ccggcgagg atcacaccaa gctgaagatg tacgggtac 1740
gccaaggcaa gaccattacc gagctgctat ctgaatacat cgcgcagcta ccagagtaa 1800
tgagcaaatg aataaatgag tagatgaatt tttagcgcta aaggaggcgg catggaaaat 1860
caagaacaac caggcaccga cgccgtggaa tgcccatgt gtggaggaaac gggcggttgg 1920
ccaggcgtaa gcggtgggt tgtctgccgg ccctgcaatg gcaactcggc cccaagccc 1980
gaggaatcgg cgtgagcggg cgcaaaccat ccggcccggg acaaatcggc gcggcgctgg 2040
gtgatgacct ggtgagaag ttgaaggccg cgaggccgc ccagcgcaa cgcatcgag 2100
cagaagcacg ccccggtgaa tcgtggcaag cgcgcgctga tccaatccgc aaagaatccc 2160
ggcaaccgcc ggcagccggg gcgcgctcga ttagggaagc gcccaagggc gacgagcaac 2220
cagatttttt cgttcgatg ctctatgacg tgggcacccg cgatagtcgc agcatcatgg 2280
acgtggccgt tttccgtctg tcgaagcgtg accgacgagc tggcgagggtg atccgctacg 2340
agcttccaga cgggcacgta gaggtttccc cagggccggc cgcatggcc agtgtgtggg 2400
attacgacct ggtactgatg cgggtttccc atctaaccga atccatgaac cgataccggg 2460
aagggaaggg agacaagccc ggcgcgctgt tccgtccaca cgttgcggac gtactcaagt 2520
tctgccggcg agccgatggc ggaaagcaga aagacgacct ggtagaaacc tgcattcggg 2580
taaacaccac gcacgttgcc atgcagcgtg cgaagaaggc caagaacggc cgctcgggta 2640
cggtatccga gggtagaagg gctacaagat cgtaaagagc gaaaccgggc 2700
ggccggagta catcgagtc gagctagtg ttggatgta ccgcgagatc acagaaggca 2760
agaaccggga cgtgctgacg gttcaccccc attacttttt gatcgatccc ggcacggcc 2820
gttttctcta ccgcctggca cgccgcgccg caggcaaggc agaagccaga tggttgttca 2880
agacgatcta cgaacgcagt ggcagcgccg gagagttcaa gaagttctgt ttcaccgtgc 2940
gcaagctgat cgggtcaaat gacctgccgg agtacgattt gaaggaggag gcggggcagg 3000
ctggcccgat cctagtcatg cgctaccgca acctgatcga ggcgaagca tccgccgggt 3060
ctaattgtac ggagcagatg ctagggcaaa ttgccctagc aggggaaaaa ggtcgaaaag 3120
gtctctttcc tgtggatagc acgtacattg ggaacccaaa gccgtacatt gggaaccgga 3180
acccgtacat tgggaaccca aagccgtaca ttgggaaccg gtcacacatg taagtgaact 3240
atataaaaaga gaaaaaaggc gattttttcc cctaaaactc tttaaaactt attaaaactc 3300
ttaaaccctg cctggcctgt gcataactgt ctggccagcg cacagccgaa gagctgcaaa 3360
aagcgcttac ccttcggteg ctgcctccc tacgcccgc cgcttcgctg cggcctatcg 3420
cggccgctgg ccgctcaaaa atggctggcc tacggccagg caatctacca ggcgcgggac 3480
aagccgcgcc gtcgccactc gaccgcggc gccacatca aggcaccctg cctcgcgcgt 3540
ttcggtgatg acggtgaaaa cctctgacac atgcagctcc cggagacggt cacagcttgt 3600
ctgtaagcgg atgccgggag cagacaagcc cgtcaggcg cgtcagcggg tgttggcggg 3660
tgtcggggcg cagccatgac ccagtcacgt agcgatagcg gagtgtatac tggcttaact 3720
atgcggcatc agagcagatt gtactgagag tgcaccatat gcggtgtgaa ataccgcaca 3780
gatgcgtaag gagaaaaata cgcatacagg gctcttccgc ttcctcgctc actgactcgc 3840
tgcgctcggt cgttcggctg cggcgagcgg tatcagctca ctcaaaggcg gtaatacggg 3900
tatccacaga atcaggggat aacgcaggaa agaacatgtg agcaaaaggc cagcaaaagg 3960
ccaggaaccg taaaaaggcc gcgttgctgg cgtttttcca taggctcgcg cccctgacg 4020
agcatcacia aaatcgacgc tcaagtcaga ggtggcgaaa cccgacagga ctataaagat 4080

```

PF 53790

79

accaggcggtt	tccccctgga	agctccctcg	tgcgctctcc	tgttccgacc	ctgccgctta	4140
ccggataacct	gtccgccttt	ctcccttcgg	gaagcggtggc	gctttctcat	agctcacgct	4200
gtaggatatct	cagttcggtg	taggtcggtc	gtcccaagct	gggctgtgtg	cacgaacccc	4260
ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtcc	aacccggtaa	4320
gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	gcgaggatg	4380
tagggcggtgc	tacagagttc	ttgaagtggg	ggcctaacta	cggctacact	agaaggacag	4440
tatttggtat	ctgcgcctcg	ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	4500
gatccggcaa	acaaaccacc	gctggtagcg	gtggtttttt	tgtttgcaag	cagcagatta	4560
cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	4620
agtggaaacga	aaactcacgt	taagggattt	tggatgca	tgatatatct	cccaatttgt	4680
gtagggctta	ttatgcacgc	ttaaaaataa	taaaagcaga	cttgacctga	tagtttggct	4740
gtgagcaatt	atgtgcttag	tgcattctaac	gcttgagtta	agccgcgcgc	cgaagcggcg	4800
toggcttgaa	cgaatttcta	gctagacatt	atttgccgac	taccttggtg	atctcgcctt	4860
tcacgtagt	gacaaattct	tccaactgat	ctgcgcgcga	ggccaagcga	tcttcttctt	4920
gtccaagata	agcctgtcta	gcttcaagta	tgacgggctg	atactgggccc	ggcaggcgct	4980
ccattgcccc	gtcggcagcg	acatccttcg	gcgcgatttt	gccggttact	gcgctgtacc	5040
aaatgcggga	caacgtaagc	actacatttc	gctcatcgcc	agcccagtcg	ggcggcgagt	5100
tccatagcgt	taaggtttca	tttagcgccct	caaatagatc	ctgttcagga	accggatcaa	5160
agagttcctc	cgccgctgga	cctaccaagg	caacgctatg	ttctcttgct	tttgctcagca	5220
agatagccag	atcaatgtcg	atcgtggctg	gctcgaagat	acctgcaaga	atgtcattgc	5280
gctgccattc	tccaaattgc	agttcgcgct	tagctggata	acgccacgga	atgatgtcgt	5340
cgtgcacaa	aatggtgact	tctacagcgc	ggagaatctc	gctctctcca	ggggaagccg	5400
aagtttccaa	aaggctcgtg	atcaaagctc	gccgcgttgt	ttcatcaagc	cttacgggtca	5460
ccgtaaccag	caaatcaata	tactgtgtg	gcttoaggcc	gccatccact	gcggagccgt	5520
acaaatgtac	ggccagcaac	gtcggttcga	gatggcgctc	gatgacgcca	actacctctg	5580
atagttgagt	cgatacttcg	gcgatcaccg	cttcccccat	gatgtttaac	tttgtttttag	5640
ggcgactgcc	ctgctgcgta	acatcgttgc	tgctccataa	catcaaacat	cgaccacagg	5700
cgtaacgcgc	ttgctgcttg	gatgcccgag	gcatagactg	tacccccaaa	aaacagtcct	5760
aacaagccat	gaaaaccgcc	actgcgttcc	atggacatac	aaatggacga	acggataaac	5820
cttttcacgc	ccttttaaat	atccgattat	tctaataaac	gctcttttct	cttaggttta	5880
cccgcgaata	tatcctgtca	aacactgata	gtttaaactg	aaggcgggaa	acgacaatca	5940
gatctagtag	gaaacagcta	tgaccatgat	tacggcaagc	ttgcatgcct	gcaggctcga	6000
tctagactag	tggatccgat	atcgcccggg	ctcgaggtag	ccatcgcgat	ccccgtcacc	6060
ggtgtgaggg	aactagtttt	gatcttgaaa	gatcttttat	cttttagagt	aagaactctt	6120
tcgtattttg	gtgaggtttt	atcctcttga	gttttggtca	tagacctatt	catggctctg	6180
ataccaattt	ttaagcgggg	gcttatgcgg	attatttctt	aaattgataa	ggggttatta	6240
gggggtatag	cgataaata	caagcattcc	cttagcgtag	agtataagta	tagtagcgta	6300
cccttatcaa	atttccatct	tcttaccttg	cacagggcct	gcaaccttat	ccttccttgt	6360
cttctctctt	ccttcctctc	acttcatcat	atttaaacca	aacctacggg	ggagtcaacg	6420
taaccaaccc	tgccttagca	tcttttccct	aacggcctcc	tgccaaagcg	gtacttctag	6480
cttcgaacgg	cgtctgggct	ccaggtttag	tcgtctcggt	tctggtttat	attcacgaca	6540
aagatctata	gggacttttag	gagatctgga	tttttagtact	ggatttttgt	tttaggaatt	6600
agaaatttta	ttgatagaag	tattttacaa	atacaaatac	atactaaggg	tttctttatc	6660
gctcaacaca	tgagcgaaac	cctataagaa	ccctaanttc	cccttatcgg	gaaactactc	6720
acacattatt	tatggagaaa	atagagagag	atagatttgt	agagagagac	tggtgatttc	6780
agcgtagccga	attcggctaa	cagtgtcgaa	taacgcttta	caaacaatta	ttaacgcccg	6840
gttaccaggc	gaagaggggc	tgtggcagat	tcatctgcag	gacggaaaaa	tcagcgccat	6900
tgatgcgcaa	tccggcggtg	tgcccataac	tgaaaacagc	ctggatgcgc	aacaaggttt	6960
agttataccg	cggtttctgg	agccacatat	tcacctggac	accacgcaaa	ccgcgggaca	7020
accgaactgg	aatcagtcgc	gcacgctgtt	tgaaggcatt	gaacgctggg	ccgagcgcaa	7080
agcggtatta	acccatgacg	atgtgaaaca	acgcgcgatg	caaacgctga	aatggcagat	7140
tgccaacggc	attcagcatg	tgcgtaccca	tgctcgatgt	tcggatgcaa	cgctaactgc	7200
gctgaaagca	atgctggaag	tgaagcagga	agtcgcgcgc	tggattgata	tgcaaatcgt	7260
cgcttccct	cagggaaggga	ttttgtcggg	tccggtgata	cctgcacatc	aacaaatttt	7320
ggtcatatat	tagaaaagtt	ataaattaaa	atatacacac	ttataaacta	cagaaaagca	7380
attgctatat	actacattct	tttattttga	aaaaaatatt	tgaatatatta	tattactact	7440

PF 53790

80

```

aattaatgat aattattata tatatatcaa aggtagaagc agaaacttac gtagtcgacg 7500
acaaaatccc ttcctgaggg aaggcgacga tttgcagatc aatccacggc gcgacttcc 7560
gcttcacttc cagcattgct ttcagcgagc ttagcggttg atccgaaaca tcgacatggg 7620
tacgcacatg ctgaatgccg ttggcaatct gccatttcag cgtttgccat gcgcgttgtt 7680
tcacatcgtc atgggttaat aacgctttgc gctcgccca gcgttcaatg cttcaaaca 7740
gcgtgccgga ctgattccag ttcggttgtc cggcggtttg cgtggtgtcc aggtgaatat 7800
gtggctccac aaacggcggt ataactaaac cttgttcggc atccaggctg tttcagtta 7860
tgggcatcac gccggattgc gcatcaatgg cgtgattttt tccgtcctgc agatgaatct 7920
gccacagccc ctcttcgcct gtaaccggg cgttaataat tgtttgtaa gcgttattcg 7980
acactgttag ccaagcttgc atgcctgcag gtcgagtctt tgttttttac tttggttcat 8040
gacactcaga gacttgagag aagcaatata tagacttttt tttgtttttt tttgtggtc 8100
acgtttatatt tccattggga gacggtaacg aagatcgaac ctgtggtgga aatgaaaca 8160
gggtgggacta gccacgtgg tttcttttct ctgcattgat ttgtttttgt tttttttgta 8220
aagttcacat caaacctact aataattgag aagaaaaata aaatctattg attgattaaa 8280
ccagccgatg ctttatgtct gaataaaaaa aagaagtga aaccccggtt aagaattaca 8340
acggtgggtt acaaagtatt tggacacaat aaatccaaac gaaataaaac aaaatggaga 8400
actaccaaata aaaaaacaaa taaaaaactt aaaagaattt attccatttt tttcccgta 8460
gaatttatct ttttatggat tccttaaatc catatttgat gcattttgat tcctcataat 8520
aggtataaat atatactatg ttatagatat gtttctaatt cgtattaacc tacctttttt 8580
tggtcgtagc attctaccta ataatttga acggaattga tgttttgga cacttagaaa 8640
gtattttttt tttggtttgt cttagctgta tttcattaaa tataaattta aataagaaat 8700
gtcataaata aaatttgacg tatagatttt ttaaattccat tttatgttat ttaattttg 8760
aaatgtgagt ttggtccta tttaatctta ggatgggtta atactaagtt ttccttaatg 8820
aattatctca gagaaactgg attaaataaa ctaaaaaata gatcaatgtg ttttggtccg 8880
gtcaaataatc tttggattta ctattattgg cgaaaagaaa gtctcatata gtaaatcata 8940
ttcctacaag agaaatcaaa atttttgaat taacatggat tgtatagttt cttatataac 9000
caattagttc gcatcaagaa aaccaaacc caattaataa tcaaacgggc ttggtaggaa 9060
tatttcattg cagctttcag ataaaagaaa aaacacaca ctcaagtctt ttatttcac 9120
tttcttactt gcaggaactc aaattccact ttgccacttt tctttacaaa taaacacaaa 9180
ttgtcaatga aacgaaatag tctttttatg caaacactgt ttgtcttttt tcatcacgt 9240
ttctgattgt gacagccatc catatatata gggaaatgta aacaacaaca tgtgaagtca 9300
catatacgta atggtttagc atagcttcta ttttcgttgt caatattagt cattccaaaa 9360
catttttaag aaaaataaat taatatatgt atattcttgg aactaatgta tgtggaata 9420
cagtaactta attattaaac attctaattg caaatatgca aagaaaaaaa agaaaagaa 9480
acaactgaaa tcaaagccag attcataata attggctaca tggttgtaga atgtagggt 9540
acacaacatc cagaattgaa cactcaaatt ggatgataga tggataatct ttagatacaa 9600
gagaattggt tctcttccat tattaacgaa aataaagaaa aaagtttag cataaaagt 9660
tgaaactcaa cataacattt tgaacttgac tccttcatag gagtgcacg aactgacgaa 9720
tcacaaccga ttacttggtt gagtcactct ccgctttctc cacttcgaa atgaatgtga 9780
ccggtttctt cgggtgctca tttacggtca agtgtaaaac atctggtctc gacgagct 9838

```

&lt;210&gt; 58

&lt;211&gt; 14184

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence: Expression  
vector pSUN1-codA-RNAi-At.Act.-2-At.Als-R-ocsT

&lt;400&gt; 58

```

ctgcttggtg ataattgtca ttagattggt tttatgcata gatgcactcg aaatcagcca 60
attttagaca agtatcaaac ggatgttaat tcagtacatt aaagacgtcc gcaatgtgtt 120
attaagttgt ctaagcgta atttgtttac accacaatat atcctgccac cagccagcca 180
acagctcccc gaccggcagc tcggcacaaa atcaccacgc gttaccacca cgccggccgg 240
ccgcatgggtg ttgaccgtgt tcgccggcat tgccgagttc gagcgttccc taatcatcga 300
ccgcacccgg agcggggcgg aggcggccaa ggcccagggc gtgaagtgtt gcccccgccc 360
taccctcacc ccggcacaga tcgcgcacgc ccgcgagctg atcgaccagg aaggccgcac 420

```



PF 53790

81

cgtgaaagag	gcggtgac	tgcttgccgt	gcacgcctcg	accctgtacc	gcgcacttga	480
gcgcagcgag	gaagtgcgc	ccaccgaggc	caggcgccgc	ggtgccttcc	gtgaggacgc	540
attgaccgag	gccgacgccc	tgccgcccgc	cgagaatgaa	cgccaagagg	aacaagcatg	600
aaaccgcacc	aggacggcca	ggacgaaccg	tttttcatta	ccgaagagat	cgaggcgagg	660
atgatcgccg	ccgggtacgt	gttcgagccg	cccgcgcacg	tctcaaccgt	gcggctgcat	720
gaaatcctgg	ccggtttgtc	tgatgccaa	ctggcgccct	ggccggccag	cttgcccgct	780
gaagaaaccg	agcgccgccc	tctaaaaagg	tgatgtgtat	ttgagtaaaa	cagcttgcgt	840
catgcggtcg	ctgcgtatat	gatgcgatga	gtaaataaac	aaatacgcaa	ggggaacgca	900
tgaaggttat	cgctgtactt	aaccagaaa	gcgggtcagg	caagacgacc	atcgcaaccc	960
atctagcccc	cgccctgcaa	ctcgccgggg	ccgatgttct	gttagtgcgt	tccgatcccc	1020
agggcagtg	ccgcgattgg	cgccgctg	gggaagatca	accgctaacc	gttgcctggca	1080
tcgaccgccc	gacgattgac	cgcgacgtga	agcccatcgg	ccggcgcgac	ttcgtagtga	1140
tcgacggagc	gccccaggcg	gcggacttgg	ctgtgtccgc	gatcaaggca	gocgacttcg	1200
tgctgattcc	ggtgcagcca	agcccttacg	acatatgggc	caccgcccgc	ctgggtggagc	1260
tggttaagca	gcgcattgag	gtcacggatg	gaaggctaca	agcggccctt	gtcgtgtcgc	1320
ggcgcatcaa	aggcacgcgc	atcgccggtg	aggttgccga	ggcgctggcc	gggtacgagc	1380
tgccattctt	tgagtcctgt	atcacgcagc	cgctgagcta	cccaggcact	gcccgcgcgc	1440
gcacaaccgt	tcttgaatca	gaacccgagg	gcgacgctgc	ccgcgaggtc	caggcgctgg	1500
ccgctgaaat	taaatacaaaa	ctcatttgag	ttaatgaggt	aaagagaaaa	tgagcaaaa	1560
cacaaacacg	ctaagtgcgc	gccgtccgag	cgacgcagc	agcaaggctg	caacgttggc	1620
cagcctggca	gacacgccag	ccatgaagcg	ggtcaacttt	cagttgccc	cgagggatca	1680
caccaagctg	aagatgtacg	cggtacgcca	aggcaagacc	attaccgagc	tgctatctga	1740
atacatcgcg	cagctaccag	agtaaatgag	caaatgaata	aatgagtaga	tgaattttag	1800
cggtataagg	aggcgccatg	gaaaatcaag	aacaaccagg	caccgacgcc	gtggaatgcc	1860
ccatgtgtgg	aggaacgggc	ggttgccag	gcgtaagcgg	ctgggttgtc	tgccggccct	1920
gcaatggcac	tggaaccccc	aagcccgagg	aatcgccgtg	agcggtcgca	aaccatccgg	1980
cccgttacaa	atcgccgccc	cgctgggtga	tgacctgggt	gagaagttag	aggccgcgca	2040
ggccgcccag	cgcaacgca	tcgaggcaga	agcacgcccc	ggtgaatcgt	ggcaagcggc	2100
cgctgatoga	atccgcaaa	aatcccgcca	accgcccggca	gccggtgcgo	cgctgattag	2160
gaagccgccc	aagggcgacg	agcaaccaga	tttttctggt	ccgatgctct	atgacgtggg	2220
caccgcgcgat	agtcgcagca	tcattggacgt	ggccgttttc	cgctctgtcg	agcgtgaccg	2280
acgagctggc	gaggtgatcc	gtacgagct	tccagacggg	cacgtagagg	tttccgcagg	2340
gcccggccgg	atggccagtg	tgtgggatta	cgacctggta	ctgatggcgg	tttcccatct	2400
aaccgaatcc	atgaaccgat	accgggaagg	gaagggagac	aagcccggcc	gcgtgttccg	2460
tccacacgtt	gcggacgtac	tcaagttctg	ccggcgagcc	gatggcgga	agcagaaaga	2520
cgacctggta	gaaacctgca	ttcgggttaa	caccacgcac	ggtgccatgc	agcgtacgaa	2580
gaaggccaag	aacggccgccc	tggtgacggt	atccgaggg	gaagccttga	ttagccgcta	2640
caagatcgta	aagagcgaaa	ccggcgggcc	ggagtacatc	gagatcgagc	tagctgattg	2700
gatgtaccgc	gagatcacag	aaggcaagaa	cccggacgtg	ctgacggttc	accccgatta	2760
ctttttgatc	gatcccgcca	tcggccggtt	tctctaccgc	ctggcacgcc	gcgcgcgagg	2820
caaggcgaga	gccagatgg	tgttcaagac	gatctacgaa	cgagtgga	gcgcgcgaga	2880
gttcaagaag	ttctgtttca	ccgtgcgcaa	gctgatcggg	tcaaatgacc	tgccggagta	2940
cgatttgaag	gaggaggcgg	ggcaggctgg	cccgatcccta	gtcatgcgct	accgcaacct	3000
gacgagggc	gaagcatccg	ccggttccta	atgtacggag	cagatgctag	ggcaaatg	3060
cctagcagg	gaaaaagg	gaaaaaggt	ctttcctgtg	gatagcacgt	acattgggaa	3120
cccaaagccg	tacattggga	accggaaccc	gtacattggg	aacccaaagc	cgtacattgg	3180
gaaccgggtca	cacatgtaag	tgactgat	aaaagagaaa	aaaggcgatt	tttccgccta	3240
aaactcttta	aaacttatta	aaactcttaa	aacccgcctg	gcctgtgcat	aactgtctgg	3300
ccagcgca	gccgaagagc	tgcaaaaagc	gcctaccctt	cggtcgctgc	gtccctacg	3360
ccccgcgct	tcgcgtcgcc	ctatcgccgc	cgctggccgc	tcaaaaatgg	ctggcctacg	3420
gccaggcaat	ctaccagggc	gcggacaagc	cgcccgctcg	ccactcgacc	gccggcgccc	3480
acatcaaggc	accctgcctc	gcgcgtttcg	gtgatgacgg	tgaaaacctc	tgacacatgc	3540
agctcccggg	gacggtcaca	gcttgtctgt	aagcggatgc	cgggagcaga	caagcccgtc	3600
agggcgcgtc	agcgggtgtt	ggcgggtgtc	ggggcgccgc	catgacccag	tcacgtagcg	3660
atagcggag	gtatactggc	ttactatgc	ggcatcagag	cagattgtac	tgagagtgca	3720
ccatatgcgg	tgtgaaatac	cgcacagatg	cgtaaggaga	aaataccgca	tcaggcgctc	3780

PF 53790

82

ttccgcttcc	tcgctcactg	actcgctgcg	ctcggtcggt	cggtcgcggc	gagcggtatc	3840
agctcactca	aaggcggtaa	tacggttatc	cacagaatca	ggggataacg	caggaaagaa	3900
catgtgagca	aaaggccagc	aaaaggccag	gaaccgtaaa	aaggccgcgt	tgctggcggt	3960
tttccatagg	ctccgcccc	ctgacgagca	tcacaaaaat	cgacgctcaa	gtcagagggtg	4020
gcgaaacccg	acaggactat	aaagatacca	ggcgtttccc	cctggaagct	ccctcgtagc	4080
ctctcctggt	ccgaccctgc	cgcttaccgg	atacctgtcc	gcctttctcc	cttcgggaag	4140
cgtggcgctt	tctcatagct	cacgctgtag	gtatctcagt	tcggtgtagg	tcgttcgctc	4200
caagctgggc	tgtgtgcacg	aacccccgt	tcagcccagc	cgctgcgcct	tatccggtaa	4260
ctatcgtctt	gagtccaacc	cggtaagaca	cgacttatcg	ccactggcag	cagccactgg	4320
taacaggatt	agcagagcga	ggtagttagg	cggtagctaca	gagttcttga	agtgggtggcc	4380
taactacggc	tacactagaa	ggacagtatt	tggtatctgc	gctctgctga	agccagttac	4440
cttcggaaaa	agagttggta	gctcttgatc	cggcaaaaaa	accaccgctg	gtagcggtgg	4500
tttttttgg	tgcaagcagc	agattacgcg	cagaaaaaaa	ggatctcaag	aagatccctt	4560
gatcttttct	acgggggtctg	acgctcagtg	gaacgaaaac	tcacgttaag	ggattttggt	4620
catgcatgat	atatctccca	atttgtgtag	ggcttattat	gcacgcttaa	aaataataaa	4680
agcagacttg	acctgatagt	ttggctgtga	gcaattatgt	gcttagtgca	tctaacgctt	4740
gagttaaagg	gcgcgcgcaa	gcggcgctcg	cttgaacgaa	tttctagcta	gacattatct	4800
gccgactacc	ttggtgatct	cgcttttcac	gtagtggaca	aattcttcca	actgatctgc	4860
gcgcgaggcc	aagcgatctt	cttcttgctc	aagataagcc	tgtctagctt	caagtatgac	4920
gggctgatac	tgggcgcgca	ggcgctccat	tgcccagtcg	gcagcgacat	ccttcggcgc	4980
gattttgccc	gttactgcgc	tgtaccaaata	gcgggacaac	gtaagcacta	catttcgctc	5040
atcgccagcc	cagtcggggc	gcgagttcca	tgcggttaag	gtttcattta	gcgcctcaaa	5100
tagatcctgt	tcaggaaccg	gatcaaagag	ttcctccgcc	gctggaccta	ccaaggcaac	5160
gctatgttct	cttgcttttg	tcagcaagat	agccagatca	atgtcgatcg	tggctggctc	5220
gaagataacct	gcaagaatgt	cattgcgctg	ccattctcca	aattgcagtt	cgcgcttagc	5280
tggataacgc	cacggaatga	tgtcgtcgtg	cacaacaatg	gtgacttcta	cagcgcgag	5340
aatctcgctc	tctccagggg	aagccgaagt	ttccaaaagg	tcggtgatca	aagctcgccg	5400
cgttggttca	tcaagcctta	cggtcaccgt	aaccagcaaa	tcaatatcac	tgtgtggctt	5460
caggcogcca	tccactgcgg	agccgtacaa	atgtagcgcc	agcaacgtcg	gttcgagatg	5520
gcgctcgatg	acgccaacta	cctctgatag	ttgagtcgat	acttcggcga	tcaccgcttc	5580
ccccatgatg	tttaactttg	ttttagggcg	actgccctgc	tgcgtaacat	cgttgctgct	5640
ccataacatc	aaacatcgac	ccacggcgta	acgcgcttgc	tgcttggtatg	cccgaggcat	5700
agactgtacc	ccaaaaaaac	agtcataaca	agccatgaaa	accgccactg	cgttccatgg	5760
acatacaaat	ggacgaacgg	ataaaccttt	tcacgcccct	ttaaatatcc	gattattcta	5820
ataaacgctc	ttttctctta	ggtttaccgc	ccaatatatc	ctgtcaaaca	ctgatagttt	5880
aaactgaagg	cgggaaacga	caatcagatc	tagtaggaaa	cagctatgac	catgattacg	5940
ccaagcttgc	atgcctgcag	gtcgactcta	gactagtggg	tccgatatcg	cccgggctcg	6000
aggtacccat	cgcgatcccc	gtcaccgggtg	tgagggaact	agttttgatc	ttgaaagatc	6060
ttttatcttt	agagtttaaga	actctttcgt	attttgggtga	ggttttatcc	tcttgagttt	6120
tggtcataga	cctattcatg	gctctgatac	caatttttaa	gcgggggctt	atgcggatta	6180
tttcttaaat	tgataagggg	ttattagggg	gtatagggta	taaatacaag	cattccctta	6240
gcgtatagta	taagtatagt	agcgtacctc	tatcaaattt	ccatcttctt	accttgacac	6300
gggcctgcaa	ccttatcctt	ccttgtcttc	ctccttcctt	ccgtccactt	catcatatct	6360
aaacctaaac	tacggggggag	tcaacgtaac	caaccctgcc	ttagcatctt	ttccctaacg	6420
gcttcctgcc	taagcggtag	ttctagcttc	gaacggcgtc	tgggctccag	gtttagtcgt	6480
ctcgtgtctg	gtttatatct	acgacaaaga	tctataggga	ctttaggaga	tctggatttt	6540
agtactggat	tttggtttta	ggaattagaa	attttattga	tagaagtatt	ttacaaatac	6600
aaatacatat	taagggtttc	ttatatgctc	aacacatgag	cgaacccta	taagaaccct	6660
aatttccctt	atcgggaaac	tactcacaca	ttattttatg	agaaaataga	gagagataga	6720
tttgtagaga	gagactgggt	atttcagcgt	accgaattcg	attttcgggt	aacagtgtcg	6780
aataacgctt	tacaacaac	tattaacgcc	cggttaccag	gcgaagagg	gctgtggcag	6840
attcatctgc	aggacggaaa	aatcagcgcc	attgatgcgc	aatccggcgt	gatgcccata	6900
actgaaaaca	gcctggatgc	cgaacaaggt	ttagttatatac	cgccgtttgt	ggagccacat	6960
attcacctgg	acaccacgca	aaccgcccga	caaccgaact	ggaatcagtc	cggcacgctg	7020
tttgaaggca	ttgaacgctg	ggccgagcgc	aaagcggtat	taaccatga	cgatgtgaaa	7080
caacgcgcat	ggcaaacgct	gaaatggcag	attgccaacg	gcattcagca	tgtgcgtacc	7140

PF 53790

83

catgtcgatg	tttcggatgc	aacgctaact	gcgctgaaag	caatgctgga	agtgaagcag	7200
gaagtcgcgc	cgtggattga	tctgcaaate	gtcgccttcc	ctcaggaagg	gattttgtcg	7260
gatccggtga	tacctgcaca	tcaacaaatt	ttggtcatat	attagaaaag	ttataaatta	7320
aatatacac	acttataaac	tacagaaaag	caattgctat	atactacatt	cttttatttt	7380
gaaaaaata	tttgaaatat	tatattacta	ctaattaatg	ataattatta	tatatatata	7440
aaaggtagaa	gcagaaactt	acgtagtcga	cgacaaaate	ccgtcctgag	ggaaggcgac	7500
gatttgacga	tcaatccacg	gcgcgacttc	ctgcttcact	tccagcattg	ctttcagcgc	7560
agttagcgtt	gcacccgaaa	catcgacatg	ggtacgcaca	tgctgaatgc	cgttggcaat	7620
ctgccatttc	agcgtttgcc	atgcgcgttg	tttcacatcg	tcatgggtta	ataacgcttt	7680
gcgctcggcc	cagcgttcaa	tgccctcaaa	cagcgtgccc	gactgattcc	agttcgggtg	7740
tccggcggtt	tgcggtggtg	ccagggtgaat	atgtggctcc	acaaacggcg	gtataactaa	7800
acctgtttcg	gcattccagg	tgttttcagt	tatgggcata	acgccggatt	gcgcatcaat	7860
ggcgctgatt	tttccgtcct	gcagatgaat	ctgccacagc	ccctcttcgc	ctggtaaccg	7920
ggcggttaata	attgtttgta	aagcgttatt	cgacactgtt	agccaagctt	gcatgcctgc	7980
aggtcgactc	tagaggatcc	ccgatccact	cgagtctttg	ttttttactt	tggttcatga	8040
cactcagaga	cttgagagaa	gcaatatata	gacttttttt	tgtttttttt	ttgtggtcac	8100
gtttattttc	ctattggaga	cggtaacgaa	gatcgaacct	gtggtggaag	tgaaacmagg	8160
tgggactagc	ccacgtgggt	tcttttctct	gcatgtgatt	gtttttgttt	tttytgtaaa	8220
gttcacatca	aacctactaa	taattgagaa	gaaaaataaa	atctattgat	tgattaaacc	8280
agccgatgct	ttatgtctga	atataaaaaa	gaagtgaaaa	ccccgtttta	gaattacaac	8340
ggtggtttac	aaagtatttt	gacacaataa	atccaaacga	aataaaaaca	aatggagAAC	8400
taccaataaa	aaaacaataa	aaaaacttaa	aagaatttat	tccatttttt	ttcccgtaga	8460
atttattctt	ttatggattc	cttaaatcca	tatttgatgc	attttgattc	ctcataatag	8520
gtaataatat	atactatgtt	atagatatgt	ttctaattcg	tattaacctc	cctttttttg	8580
gtcgtacgat	tctacctaat	aatattgaac	ggaattgatg	ttttggacca	cttagaaagt	8640
attttttttt	tggtttgtct	tagctgtatt	tcattaaata	taaattttaa	taagaaatgt	8700
cataaataaa	atttgacgta	tagatttttt	aaatccattt	tatgtttatt	aatatttgaa	8760
atgtgagttt	ggctcctatt	taatcttagg	atgggttaat	actaagtttt	ccttaatgaa	8820
ttatctcaga	gaaactggat	taaataaact	aaaaaataga	tcaatgtgtt	ttggtccggt	8880
caaatatctt	tggttttact	attattggcg	aaaagaaagt	ctcatatagt	aaatcatatt	8940
cctacaagag	aatcaaaaat	ttttgaatta	acatggattg	tatagtttct	tatataacca	9000
attagttcgc	atcaagaaaa	ccaaacccca	attaataatc	aaacgggctt	ggtaggaata	9060
tttcatttgc	gctttcagat	aaaagaaaaa	aacacacact	caagtctttt	atttcatctt	9120
tcttacttgc	aggaactcaa	attocacttt	gccacttttc	tttacaataa	aacacaaatt	9180
gtcaatgaaa	cgaaatagtc	tttttatgca	aacactgttt	gtcttttttc	gatcacgttt	9240
ctgattgtga	cagccatcca	tatatatagg	gaatgtaaaa	caacaacatg	tgaagtcaca	9300
tatacgtaat	ggttttagcat	agcttctatt	ttcggtgtca	atattagtca	ttccaaaaaca	9360
tttttaagaa	aaataaatta	atatatgtat	attcttggaa	ctaattgtatg	tggaaatata	9420
gtaacttaat	tattaaacat	tctaaatgca	aatatgcaaa	gaaaaaaaag	aaaaagaacac	9480
aactgaaatc	aaagccagat	tcataataat	tggctacatg	gttgtagaat	gtagggtaac	9540
acaacatcca	gaattgaaca	ctcaaattgg	atgatagatg	gataatcttt	agatacaaga	9600
gaattgggtc	tcttccatta	ttaacgaaaa	taaagaaaaa	aagtttagca	taaaagtttg	9660
aaactcaaca	taacattttg	aacttgactc	cttcatagga	gtgacatgaa	ctgacgaatc	9720
acaaccgatt	acttgtttga	gtcatcttcc	gctttctcca	ccttcgaaat	gaatgtgacc	9780
ggtttcttcg	ggtgctcatt	tacggtcaag	tgtaaaacat	ctggctctga	gtaattgtcca	9840
accgaatcga	agtacaactt	agctcttgct	acatcaccaa	gatcttgatg	ggggatcggy	9900
taccgagctc	gaattcactg	gcggtcgttt	tacaacgact	cagcacgcgt	tggtttcgac	9960
aaaattttaga	acgaacttaa	ttatgatctc	aaatacattg	atacatatct	catctagatc	10020
taggttatca	ttatgtaaga	aagttttgac	gaatatggca	cgacaaaatg	gctagactcg	10080
atgtaattgg	tatctcaact	caacattata	cttataccac	acattagtta	gacaaaattt	10140
aaacaactat	tttttatgta	tgcaagagtc	agcatatgta	taattgattc	agaatcggtt	10200
tgacgagttc	ggatgtagta	gtagccatta	tttaatgtac	atactaatac	tgaatagtga	10260
atatgatgaa	acattgtatc	ttattgtata	aatatccata	aacacatcat	gaaagacact	10320
ttctttcacg	gtctgaatta	attatgatac	aattcttaata	gaaaacgaat	taaattacgt	10380
tgaattgtat	gaaatctaata	tgaacaagcc	aaccacgacg	acgactaacg	ttgcctggat	10440
tgactcgggt	taagttaacc	actaaaaaaa	cggagctgtc	atgtaaacac	cggatcgagc	10500

PF 53790

84

aggtcacagt	catgaagcca	tcaaagcaaa	agaactaate	caagggctga	gatgattaat	10560
tagtttaaaa	attagttaac	acgagggaaa	aggctgtctg	acagccaggt	cacgttatct	10620
ttacctgtgg	tcgaaatgat	tcgtgtctgt	cgattttaat	tatttttttg	aaaggccgaa	10680
aataaagttg	taagagataa	accgcctat	ataaatccat	atattttcct	ctccgctttg	10740
aattgtctcg	ttgtccctcct	cactttcatc	agccgttttg	aatctccggc	gacttgacag	10800
agaagaacaa	ggaagaagac	taagagagaa	agtaagagat	aatccaggag	attcattctc	10860
cgttttgaat	cttcctcaat	ctcatcttct	tccgctcttt	ctttccaagg	taataggaa	10920
tttctggatc	tactttattt	gctggatctc	gatcttgttt	tctcaatttc	cttgagatct	10980
ggaattcggt	taatttggat	ctgtgaacct	ccactaaatc	ttttggtttt	actagaatcg	11040
atctaagttg	accgatcagt	tagctcgatt	atagctacca	gaatttggct	tgaccttgat	11100
ggagagatcc	atgttcatgt	tacctgggaa	atgatttgta	tatgtgaatt	gaaatctgaa	11160
ctgttgaagt	tagattgaat	ctgaacactg	tcaattgtag	attgaatctg	aacactgttt	11220
aaggttagat	gaagtttgtg	tatagattct	tcgaaacttt	aggatttgta	gtgtcgtagc	11280
ttgaacagaa	agctatttct	gattcaatca	gggtttattt	gactgtattg	aactcttttt	11340
gtgtgtttgc	agctcataaa	aaaaacgcga	acctgcaggc	atggcggcgg	caacaacaac	11400
aacaacaaca	tcttcttcga	tctccttctc	caccaaacca	tctccttctc	cctccaaaatc	11460
accattacca	atctccagat	tctcctctcc	attctcccta	aaacccaaca	aatcatctct	11520
ctcctcccg	cgcgcggta	tcaaatccag	ctctccctcc	tccatctccg	ccgtgctcaa	11580
cacaaccacc	aatgtcacaa	ccactccctc	tccaaccaaa	cctaccaaac	ccgaaacatt	11640
catctcccg	ttcgctccag	atcaaccctg	caaaggcgct	gatatactcg	togaagcttt	11700
agaacgtcaa	ggcgtagaaa	ccgtattcgc	ttaccctgga	ggtagcatcaa	tggagattca	11760
ccaagcctta	accgcctctt	cctcaatccg	taacgtcctt	cctcgctcag	aacaaggagg	11820
tgtattcgca	gcagaaggat	acgctcgatc	ctcaggtaaa	ccaggatatc	gtatagccac	11880
ttcagggtccc	ggagctacaa	atctcgttag	cggattagcc	gatgcgttgt	tagatagttg	11940
tcctcttgta	gcaatcacag	gacaagtccc	tcgtcgtagt	attggtacag	atgcgtttca	12000
agagactccg	attgttgagg	taacgcgttc	gattacgaag	cataactatc	ttgtgatgga	12060
tgttgaagat	atccctagga	ttattgagga	agctttcttt	ttagctactt	ctggtagacc	12120
tggacctggt	ttggttgatg	ttcctaaga	tattcaacaa	cagcttgoga	ttcctaattg	12180
ggaacaggct	atgagattac	ctggttatat	gtctaggatg	cctaaccctc	cggaagattc	12240
tcatttgagg	cagattgtta	gggtgatttc	tgagtctaag	aagcctgtgt	tgtatggttg	12300
tgggtggtgt	ttgaattcta	gcgatgaatt	gggtaggttt	gttgagctta	cggggatccc	12360
tgttgcgagt	acgttgatgg	ggctgggac	ttatccttgt	gatgatgagt	tgctggtaca	12420
tatgcttgga	atgcatggga	ctgtgtatgc	aaattacgct	gtggagcata	gtgatttggt	12480
gttgccgttt	ggggtagggt	ttgatgatcg	tgtaacgggt	aagcttgagg	cttttgctag	12540
tagggctaag	attgttcata	ttgatattga	ctcggctgag	attgggaaga	ataagactcc	12600
tcatgtgtct	gtgtgtggtg	atgttaagct	ggctttgcaa	gggatgaata	aggttccttg	12660
gaaccgagcg	gaggagctta	agcttgattt	tggagtttg	aggatgagt	tgaacgtaca	12720
gaaacagaag	tttccgttga	gctttaagac	gtttggggaa	gctattcctc	cacagtatgc	12780
gattaaggtc	cttgatgagt	tgactgatgg	aaaagccata	ataagtactg	gtgtcgggca	12840
acatcaaatg	tgggcggcgc	agttctacaa	ttacaagaaa	ccaaggcagt	ggctatcatc	12900
aggaggcctt	ggagctatgg	gatttggact	tcctgtctgc	attggagcgt	ctgttgctaa	12960
ccctgatgcg	atagttgtgg	atattgacgg	agatggaagc	tttataatga	atgtgcaaga	13020
gctagccact	attcgtgtag	agaatcttcc	agtgaaggta	cttttattaa	acaaccagca	13080
tcttgccatg	gttatgcaat	gggaagatcg	gttctacaaa	gctaaccgag	ctcacacatt	13140
tctcggggat	ccggctcagg	aggacgagat	attcccgaac	atgttgctgt	ttgcagcagc	13200
ttgcgggatt	ccagcggcga	gggtgacaaa	gaaagcagat	ctccgagaag	ctattcagac	13260
aatgctggat	acaccaggac	cttacctggt	ggatgtgatt	tgtccgcacc	aagaacatgt	13320
gttgccgatg	atcccgaatg	gtggcacttt	caacgatgtc	ataacggaag	gagatggccg	13380
gattaaatag	tgagagatga	aaccggcctg	gcccggcccg	agtggggagg	cacgatggcc	13440
gctttggtcg	atcgacggga	tcgatcctgc	tttaatgaga	tatgcgagac	gcctatgatc	13500
gcattgatatt	tgttttcaat	tctgttgtgc	acgttgtaaa	aaacctgagc	atgtgtagct	13560
cagatcctta	ccgcgggttt	cggttcattc	taatgaatat	atcaccctgt	actatcgtat	13620
ttttatgaat	aatattctcc	gttcaattta	ctgattgtac	cctactactt	atattgtaca	13680
tattaaaatg	aaaacaatat	attgtgctga	ataggtttat	agcgacatct	atgatagagc	13740
gccacaataa	caaacaattg	cgttttatta	ttacaaatcc	aatttttaaaa	aaagcggcag	13800
aaccgggtcaa	acctaaaaga	ctgattacat	aatctttatt	caaatttcaa	aaggccccag	13860

PF 53790

85

```

gggctagtat ctacgacaca ccgagcggcg aactaataac gttcactgaa gggaactccg 13920
gttccccgcc ggcgcgcgatg ggtgagattc cttgaagttg agtattggcc gtccgctcta 13980
ccgaaagtta cgggcacccat tcaaccocggc ccagcacggc ggccgggtaa ccgacttgct 14040
gccccgagaa ttatgcagca tttttttggt gtatgtgggc cccaaatgaa gtgcaggtca 14100
aaccttgaca gtgacgacaa atcgttgggc ggggtccaggc cgaattttgc gacaacatgt 14160
cgaggctcag caggatgggc ccag                                     14184

```

&lt;210&gt; 59

&lt;211&gt; 1011

&lt;212&gt; DNA

&lt;213&gt; Zea mays

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(981)

&lt;223&gt; coding for 5-methylthioribose kinase

&lt;400&gt; 59

```

gca cga gca ctc ctc tcc tct cct ctc gcc ggc gca tcg ccc gac tgt 48
Ala Arg Ala Leu Leu Ser Ser Pro Leu Ala Gly Ala Ser Pro Asp Cys
1 5 10 15

cag tca gcc tca gcc atg gcc gcg gag gag gag cag ggc ttc cgc ccg 96
Gln Ser Ala Ser Ala Met Ala Ala Glu Glu Glu Gln Gly Phe Arg Pro
20 25 30

ctg gac gag tcg tcc ctg ctc gcc tac atc aag gcc acg ccg gcg ctc 144
Leu Asp Glu Ser Ser Leu Leu Ala Tyr Ile Lys Ala Thr Pro Ala Leu
35 40 45

gcc tcc cgc ctc ggc ggc ggt ggc agt cta gac tcc atc gag atc aag 192
Ala Ser Arg Leu Gly Gly Gly Gly Ser Leu Asp Ser Ile Glu Ile Lys
50 55 60

gag gtc ggc gac ggc aac ctc aac ttc gtc tac atc gtg cag tcc gag 240
Glu Val Gly Asp Gly Asn Leu Asn Phe Val Tyr Ile Val Gln Ser Glu
65 70 75 80

gcc ggc gcc atc gtc gtc aag cag gcg ctc ccg tac gtg cgc tgc gtg 288
Ala Gly Ala Ile Val Val Lys Gln Ala Leu Pro Tyr Val Arg Cys Val
85 90 95

ggg gat tcg tgg ccc atg acg cgg gag cgc gcc tac ttc gag gcc tcc 336
Gly Asp Ser Trp Pro Met Thr Arg Glu Arg Ala Tyr Phe Glu Ala Ser
100 105 110

acg ctg cgg gag cac ggc cgc ctg tgc ccg gag cac acc ccc gag gtg 384
Thr Leu Arg Glu His Gly Arg Leu Cys Pro Glu His Thr Pro Glu Val
115 120 125

tac cac ttc gac cgg acc ttg tcg ctg atg ggg atg cgc tac atc gag 432
Tyr His Phe Asp Arg Thr Leu Ser Leu Met Gly Met Arg Tyr Ile Glu
130 135 140

ccc ccg cac atc atc ctc cgc aag ggc ctc gtc gcc ggt gtc gag tac 480
Pro Pro His Ile Ile Leu Arg Lys Gly Leu Val Ala Gly Val Glu Tyr
145 150 155 160

ccg ctg ctc gcc gac cac atg tcc gat tac atg gcc aag acg ctc ttc 528
Pro Leu Leu Ala Asp His Met Ser Asp Tyr Met Ala Lys Thr Leu Phe
165 170 175

ttc acc tcc ctc ctc tat aac aat acc acg gat cat aag aac gga gtt 576
Phe Thr Ser Leu Leu Tyr Asn Asn Thr Thr Asp His Lys Asn Gly Val
180 185 190

```

PF 53790

86

```

gct aag tac tct gcg aac gtg gag atg tgt agg ctc acg gag caa gtt 624
Ala Lys Tyr Ser Ala Asn Val Glu Met Cys Arg Leu Thr Glu Gln Val
      195                200                205

gtg ttc tcg gac cca tac cgt gtt tcc aaa ttt aat cgg tgg acc tcg 672
Val Phe Ser Asp Pro Tyr Arg Val Ser Lys Phe Asn Arg Trp Thr Ser
      210                215                220

cct tat ctc gac aaa gat gct gag gca gtt cgc gag gat gat gag ctc 720
Pro Tyr Leu Asp Lys Asp Ala Glu Ala Val Arg Glu Asp Asp Glu Leu
      225                230                235                240

aag ttg gaa gta gct ggg ctg aaa tcg atg ttt atc gag aga gct caa 768
Lys Leu Glu Val Ala Gly Leu Lys Ser Met Phe Ile Glu Arg Ala Gln
      245                250                255

gct ctg att cat gga gat ctc cac act ggt tct atc atg gtg acc gaa 816
Ala Leu Ile His Gly Asp Leu His Thr Gly Ser Ile Met Val Thr Glu
      260                265                270

gtt caa ctc aag tca ttg atc cag aat ttg ggt tct atg ggg cca atg 864
Val Gln Leu Lys Ser Leu Ile Gln Asn Leu Gly Ser Met Gly Pro Met
      275                280                285

ggg ttt gat att ggg agc ctt cct tgg aaa cct gat ttt ggg cat act 912
Gly Phe Asp Ile Gly Ser Leu Pro Trp Lys Pro Asp Phe Gly His Thr
      290                295                300

atg cac aga atg ggc atg ctg atc aag cga atg atc gta agg ctt aca 960
Met His Arg Met Gly Met Leu Ile Lys Arg Met Ile Val Arg Leu Thr
      305                310                315                320

aga atg gat ctt gaa gac aat tgaagagtcg tggaatttgt tccacaaaaa 1011
Arg Met Asp Leu Glu Asp Asn                                325

```

&lt;210&gt; 60

&lt;211&gt; 327

&lt;212&gt; PRT

&lt;213&gt; Zea mays

&lt;400&gt; 60

```

Ala Arg Ala Leu Leu Ser Ser Pro Leu Ala Gly Ala Ser Pro Asp Cys
  1                5                10                15

Gln Ser Ala Ser Ala Met Ala Ala Glu Glu Gln Gly Phe Arg Pro
      20                25                30

Leu Asp Glu Ser Ser Leu Leu Ala Tyr Ile Lys Ala Thr Pro Ala Leu
      35                40                45

Ala Ser Arg Leu Gly Gly Gly Gly Ser Leu Asp Ser Ile Glu Ile Lys
      50                55                60

Glu Val Gly Asp Gly Asn Leu Asn Phe Val Tyr Ile Val Gln Ser Glu
      65                70                75                80

Ala Gly Ala Ile Val Val Lys Gln Ala Leu Pro Tyr Val Arg Cys Val
      85                90                95

Gly Asp Ser Trp Pro Met Thr Arg Glu Arg Ala Tyr Phe Glu Ala Ser
      100                105                110

Thr Leu Arg Glu His Gly Arg Leu Cys Pro Glu His Thr Pro Glu Val
      115                120                125

```

PF 53790

87

Tyr His Phe Asp Arg Thr Leu Ser Leu Met Gly Met Arg Tyr Ile Glu  
 130 135 140  
 Pro Pro His Ile Ile Leu Arg Lys Gly Leu Val Ala Gly Val Glu Tyr  
 145 150 155 160  
 Pro Leu Leu Ala Asp His Met Ser Asp Tyr Met Ala Lys Thr Leu Phe  
 165 170 175  
 Phe Thr Ser Leu Leu Tyr Asn Asn Thr Thr Asp His Lys Asn Gly Val  
 180 185 190  
 Ala Lys Tyr Ser Ala Asn Val Glu Met Cys Arg Leu Thr Glu Gln Val  
 195 200 205  
 Val Phe Ser Asp Pro Tyr Arg Val Ser Lys Phe Asn Arg Trp Thr Ser  
 210 215 220  
 Pro Tyr Leu Asp Lys Asp Ala Glu Ala Val Arg Glu Asp Asp Glu Leu  
 225 230 235 240  
 Lys Leu Glu Val Ala Gly Leu Lys Ser Met Phe Ile Glu Arg Ala Gln  
 245 250 255  
 Ala Leu Ile His Gly Asp Leu His Thr Gly Ser Ile Met Val Thr Glu  
 260 265 270  
 Val Gln Leu Lys Ser Leu Ile Gln Asn Leu Gly Ser Met Gly Pro Met  
 275 280 285  
 Gly Phe Asp Ile Gly Ser Leu Pro Trp Lys Pro Asp Phe Gly His Thr  
 290 295 300  
 Met His Arg Met Gly Met Leu Ile Lys Arg Met Ile Val Arg Leu Thr  
 305 310 315 320  
 Arg Met Asp Leu Glu Asp Asn  
 325

&lt;210&gt; 61

&lt;211&gt; 471

&lt;212&gt; DNA

&lt;213&gt; Brassica napus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (2)..(469)

&lt;223&gt; coding for 5-methylthioribose kinase

&lt;400&gt; 61

a ttt ccg ggt cga cga ttt cgt ggc aat ctc aac ttc gtt ttc atc gtc 49  
 Phe Pro Gly Arg Arg Phe Arg Gly Asn Leu Asn Phe Val Phe Ile Val  
 1 5 10 15

atc gga tcc act ggc tca ctc gtc atc aaa cag gcg ctt ccg tat ata 97  
 Ile Gly Ser Thr Gly Ser Leu Val Ile Lys Gln Ala Leu Pro Tyr Ile  
 20 25 30

cgt tgt att ggg gag tct tgg cca atg acg aaa gaa aga gct tac ttt 145  
 Arg Cys Ile Gly Glu Ser Trp Pro Met Thr Lys Glu Arg Ala Tyr Phe  
 35 40 45

gaa gct aca act ctg aga aag cac gga gct ttg tct cct gat cat gtt 193  
 Glu Ala Thr Thr Leu Arg Lys His Gly Ala Leu Ser Pro Asp His Val  
 50 55 60

PF 53790

88

```

cct gaa gtc tac cat ttt gac agg acc atg gct ttg att gga atg agg 241
Pro Glu Val Tyr His Phe Asp Arg Thr Met Ala Leu Ile Gly Met Arg
 65              70              75              80

tat ctg gag cct cct cac atc atc ctc cgc aaa gga ctc gtt gct gga 289
Tyr Leu Glu Pro Pro His Ile Ile Leu Arg Lys Gly Leu Val Ala Gly
              85              90              95

atc cag tac cct ttc ctt gca gaa cac atg gct gat tac atg gcc aaa 337
Ile Gln Tyr Pro Phe Leu Ala Glu His Met Ala Asp Tyr Met Ala Lys
              100              105              110

acc ctc ttc ttc act tcg ctc ctc tat cat gat acc aca gag cac aaa 385
Thr Leu Phe Phe Thr Ser Leu Leu Tyr His Asp Thr Thr Glu His Lys
              115              120              125

aga gca gta acc gag ttt tgt ggt aat gtg gag tta tgc cgg tta acg 433
Arg Ala Val Thr Glu Phe Cys Gly Asn Val Glu Leu Cys Arg Leu Thr
              130              135              140

gag caa gta gtg ttc tct gac ccg tat aga gtt tct ag 471
Glu Gln Val Val Phe Ser Asp Pro Tyr Arg Val Ser
145              150              155

```

&lt;210&gt; 62

&lt;211&gt; 156

&lt;212&gt; PRT

&lt;213&gt; Brassica napus

&lt;400&gt; 62

```

Phe Pro Gly Arg Arg Phe Arg Gly Asn Leu Asn Phe Val Phe Ile Val
 1              5              10              15

Ile Gly Ser Thr Gly Ser Leu Val Ile Lys Gln Ala Leu Pro Tyr Ile
              20              25              30

Arg Cys Ile Gly Glu Ser Trp Pro Met Thr Lys Glu Arg Ala Tyr Phe
              35              40              45

Glu Ala Thr Thr Leu Arg Lys His Gly Ala Leu Ser Pro Asp His Val
              50              55              60

Pro Glu Val Tyr His Phe Asp Arg Thr Met Ala Leu Ile Gly Met Arg
              65              70              75              80

Tyr Leu Glu Pro Pro His Ile Ile Leu Arg Lys Gly Leu Val Ala Gly
              85              90              95

Ile Gln Tyr Pro Phe Leu Ala Glu His Met Ala Asp Tyr Met Ala Lys
              100              105              110

Thr Leu Phe Phe Thr Ser Leu Leu Tyr His Asp Thr Thr Glu His Lys
              115              120              125

Arg Ala Val Thr Glu Phe Cys Gly Asn Val Glu Leu Cys Arg Leu Thr
              130              135              140

Glu Gln Val Val Phe Ser Asp Pro Tyr Arg Val Ser
145              150              155

```

&lt;210&gt; 63

&lt;211&gt; 415

&lt;212&gt; DNA

&lt;213&gt; Brassica napus



PF 53790

89

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (3)..(413)

&lt;223&gt; coding for 5-methylthioribose kinase

&lt;400&gt; 63

```

gg gtc gac gat ttc gtg ctg aga gca aaa gag atg tcg ttc gat gag      47
  Val Asp Asp Phe Val Leu Arg Ala Lys Glu Met Ser Phe Asp Glu
    1             5             10             15

ttc aag ccg ttg aac gag aaa tct cta gta gag tac ata aag gca acg      95
Phe Lys Pro Leu Asn Glu Lys Ser Leu Val Glu Tyr Ile Lys Ala Thr
              20             25             30

cct gcc ctc tcc tcc agg ctc gga gac aag tac gat gat ctg gtc atc     143
Pro Ala Leu Ser Ser Arg Leu Gly Asp Lys Tyr Asp Asp Leu Val Ile
              35             40             45

aag gaa gtt gga gat ggc aat ctc aac ttc gtt ttc atc gtt gtc gga     191
Lys Glu Val Gly Asp Gly Asn Leu Asn Phe Val Phe Ile Val Val Gly
              50             55             60

tcc act ggc tca ctc gtc atc aaa cag gcg ctt ccg tat ata cgt tgt     239
Ser Thr Gly Ser Leu Val Ile Lys Gln Ala Leu Pro Tyr Ile Arg Cys
              65             70             75

att gga gaa tca tgg cca atg acg aaa gaa aga gct tac ttt gaa gca     287
Ile Gly Glu Ser Trp Pro Met Thr Lys Glu Arg Ala Tyr Phe Glu Ala
              80             85             90             95

aca act ctg aga aag cac ggt ggt ttg tct ccg gat cat gtt cct gaa     335
Thr Thr Leu Arg Lys His Gly Gly Leu Ser Pro Asp His Val Pro Glu
              100            105            110

gtc tac cat ttt gac aga acc atg gct ttg att gga atg aga tac ctc     383
Val Tyr His Phe Asp Arg Thr Met Ala Leu Ile Gly Met Arg Tyr Leu
              115            120            125

gag cct cct cac atc atc ctc cgc aaa gga ct                          415
Glu Pro Pro His Ile Ile Leu Arg Lys Gly
              130            135

```

&lt;210&gt; 64

&lt;211&gt; 137

&lt;212&gt; PRT

&lt;213&gt; Brassica napus

&lt;400&gt; 64

```

Val Asp Asp Phe Val Leu Arg Ala Lys Glu Met Ser Phe Asp Glu Phe
  1             5             10             15

Lys Pro Leu Asn Glu Lys Ser Leu Val Glu Tyr Ile Lys Ala Thr Pro
  20             25             30

Ala Leu Ser Ser Arg Leu Gly Asp Lys Tyr Asp Asp Leu Val Ile Lys
  35             40             45

Glu Val Gly Asp Gly Asn Leu Asn Phe Val Phe Ile Val Val Gly Ser
  50             55             60

Thr Gly Ser Leu Val Ile Lys Gln Ala Leu Pro Tyr Ile Arg Cys Ile
  65             70             75             80

Gly Glu Ser Trp Pro Met Thr Lys Glu Arg Ala Tyr Phe Glu Ala Thr
  85             90             95

```

PF 53790

90

Thr Leu Arg Lys His Gly Gly Leu Ser Pro Asp His Val Pro Glu Val  
 100 105 110  
 Tyr His Phe Asp Arg Thr Met Ala Leu Ile Gly Met Arg Tyr Leu Glu  
 115 120 125  
 Pro Pro His Ile Ile Leu Arg Lys Gly  
 130 135

&lt;210&gt; 65

&lt;211&gt; 424

&lt;212&gt; DNA

&lt;213&gt; Oryza sativa

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (3)..(422)

&lt;223&gt; coding for 5-methylthioribose kinase

&lt;400&gt; 65

cc ctt ctc tac aac tcc acc act gat cac aag aaa gga gtt gct cag 47  
 Leu Leu Tyr Asn Ser Thr Thr Asp His Lys Lys Gly Val Ala Gln  
 1 5 10 15  
 tac tgc gat aat gtg gag atg tgt agg ctc aca gag caa gtc gtg ttc 95  
 Tyr Cys Asp Asn Val Glu Met Cys Arg Leu Thr Glu Gln Val Val Phe  
 20 25 30  
 tca gac cca tac atg ctc gcc aaa tac aat cgt tgc aca tca ccc ttc 143  
 Ser Asp Pro Tyr Met Leu Ala Lys Tyr Asn Arg Cys Thr Ser Pro Phe  
 35 40 45  
 cta gat aat gat gct gca gcg gtt cga gag gat gct gag ctt aaa ttg 191  
 Leu Asp Asn Asp Ala Ala Ala Val Arg Glu Asp Ala Glu Leu Lys Leu  
 50 55 60  
 gag att gct gaa ttg aaa tca atg ttt att gag aga gca cag gct ctt 239  
 Glu Ile Ala Glu Leu Lys Ser Met Phe Ile Glu Arg Ala Gln Ala Leu  
 65 70 75  
 ctt cat gga gat ctc cac act ggt tcc atc atg gtg aca cca gat tct 287  
 Leu His Gly Asp Leu His Thr Gly Ser Ile Met Val Thr Pro Asp Ser  
 80 85 90 95  
 act caa gtg att gat cca gaa ttt gct ttc tat ggc cca atg ggt tac 335  
 Thr Gln Val Ile Asp Pro Glu Phe Ala Phe Tyr Gly Pro Met Gly Tyr  
 100 105 110  
 gac att ggg gcc ttc ctg ggg aac ttg att ttg gca tat ttt tca caa 383  
 Asp Ile Gly Ala Phe Leu Gly Asn Leu Ile Leu Ala Tyr Phe Ser Gln  
 115 120 125  
 gat gga cac gct gat caa gca aat gat cgt aag gct tac aa 424  
 Asp Gly His Ala Asp Gln Ala Asn Asp Arg Lys Ala Tyr  
 130 135 140

&lt;210&gt; 66

&lt;211&gt; 140

&lt;212&gt; PRT

&lt;213&gt; Oryza sativa

&lt;400&gt; 66

Leu Leu Tyr Asn Ser Thr Thr Asp His Lys Lys Gly Val Ala Gln Tyr  
 1 5 10 15

PF 53790

## 91

Cys Asp Asn Val Glu Met Cys Arg Leu Thr Glu Gln Val Val Phe Ser  
                   20                                  25                                  30  
 Asp Pro Tyr Met Leu Ala Lys Tyr Asn Arg Cys Thr Ser Pro Phe Leu  
                   35                                  40                                  45  
 Asp Asn Asp Ala Ala Ala Val Arg Glu Asp Ala Glu Leu Lys Leu Glu  
                   50                                  55                                  60  
 Ile Ala Glu Leu Lys Ser Met Phe Ile Glu Arg Ala Gln Ala Leu Leu  
                   65                                  70                                  75                                  80  
 His Gly Asp Leu His Thr Gly Ser Ile Met Val Thr Pro Asp Ser Thr  
                                   85                                  90                                  95  
 Gln Val Ile Asp Pro Glu Phe Ala Phe Tyr Gly Pro Met Gly Tyr Asp  
                                   100                                  105                                  110  
 Ile Gly Ala Phe Leu Gly Asn Leu Ile Leu Ala Tyr Phe Ser Gln Asp  
                   115                                  120                                  125  
 Gly His Ala Asp Gln Ala Asn Asp Arg Lys Ala Tyr  
                   130                                  135                                  140

&lt;210&gt; 67

&lt;211&gt; 404

&lt;212&gt; DNA

&lt;213&gt; Glycine max

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (3)..(404)

&lt;223&gt; coding for 5-methylthioribose kinase

&lt;400&gt; 67

ta atc ccc gaa cat gtt cct gaa gtg tat cac ttt gac cgt acc atg 47  
   Ile Pro Glu His Val Pro Glu Val Tyr His Phe Asp Arg Thr Met  
     1                                  5                                  10                                  15  
 tct ttg atc ggt atg cgt tac ttg gag ccc cca cat ata atc ctc ata 95  
   Ser Leu Ile Gly Met Arg Tyr Leu Glu Pro Pro His Ile Ile Leu Ile  
                   20                                  25                                  30  
 aaa ggg ttg att gct ggg att gag tac cct ttt ttg gct gaa cac atg 143  
   Lys Gly Leu Ile Ala Gly Ile Glu Tyr Pro Phe Leu Ala Glu His Met  
                   35                                  40                                  45  
 gct gat ttc atg gcg aag aca ctc ttc ttc acg tct ctg ctt ttc cgt 191  
   Ala Asp Phe Met Ala Lys Thr Leu Phe Phe Thr Ser Leu Leu Phe Arg  
                   50                                  55                                  60  
 tcc act gct gac cac aaa cgg gac gtt gcc gaa ttt tgt ggg aat gtg 239  
   Ser Thr Ala Asp His Lys Arg Asp Val Ala Glu Phe Cys Gly Asn Val  
                   65                                  70                                  75  
 gag tta tgc agg ctc act gaa cag gtc gtt ttc tct gac cct tat aaa 287  
   Glu Leu Cys Arg Leu Thr Glu Gln Val Val Phe Ser Asp Pro Tyr Lys  
                   80                                  85                                  90                                  95  
 gtt tct caa tat aat cgt tgg act tcc ccc tat ctt gat cgt gat gct 335  
   Val Ser Gln Tyr Asn Arg Trp Thr Ser Pro Tyr Leu Asp Arg Asp Ala  
                   100                                  105                                  110  
 gag gct gtt cgg gaa gac aat ctg ctg aag ctt gaa gtt gct gag ctg 383  
   Glu Ala Val Arg Glu Asp Asn Leu Leu Lys Leu Glu Val Ala Glu Leu  
                   115                                  120                                  125

PF 53790

92

aaa tcc aag ttc att gag agc  
 Lys Ser Lys Phe Ile Glu Ser  
 130

404

&lt;210&gt; 68

&lt;211&gt; 134

&lt;212&gt; PRT

&lt;213&gt; Glycine max

&lt;400&gt; 68

Ile Pro Glu His Val Pro Glu Val Tyr His Phe Asp Arg Thr Met Ser  
 1 5 10 15  
 Leu Ile Gly Met Arg Tyr Leu Glu Pro Pro His Ile Ile Leu Ile Lys  
 20 25 30  
 Gly Leu Ile Ala Gly Ile Glu Tyr Pro Phe Leu Ala Glu His Met Ala  
 35 40 45  
 Asp Phe Met Ala Lys Thr Leu Phe Phe Thr Ser Leu Leu Phe Arg Ser  
 50 55 60  
 Thr Ala Asp His Lys Arg Asp Val Ala Glu Phe Cys Gly Asn Val Glu  
 65 70 75 80  
 Leu Cys Arg Leu Thr Glu Gln Val Val Phe Ser Asp Pro Tyr Lys Val  
 85 90 95  
 Ser Gln Tyr Asn Arg Trp Thr Ser Pro Tyr Leu Asp Arg Asp Ala Glu  
 100 105 110  
 Ala Val Arg Glu Asp Asn Leu Leu Lys Leu Glu Val Ala Glu Leu Lys  
 115 120 125  
 Ser Lys Phe Ile Glu Ser  
 130

&lt;210&gt; 69

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence:  
 oligonucleotide primer

&lt;400&gt; 69

cgtgaatacgcggtggagtc g

21

&lt;210&gt; 70

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

<223> Description of the artificial sequence:  
 oligonucleotide primer

&lt;400&gt; 70

cggcaggata atcaggttg

20

&lt;210&gt; 71

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

PF 53790

93

<220>

<223> Description of the artificial sequence:  
oligonucleotide primer

<400> 71

gtcaacgtaa ccaaccctgc

20

We claim:

- 5 1. A process for preparing transformed plant cells or organisms,  
which comprises the following steps:
  - 10 a) transforming a population of plant cells, with the cells  
of said population containing at least one marker protein  
capable of causing directly or indirectly a toxic effect  
for said population, with at least one nucleic acid se-  
quence to be inserted in combination with at least one  
15 double-stranded marker protein ribonucleic acid sequence  
or an expression cassette or expression cassettes ensur-  
ing expression thereof capable of reducing the expression  
of at least one marker protein, and
  - 20 b) selecting transformed plant cells whose genome contains  
said nucleic acid sequence and which have a growth advan-  
tage over nontransformed cells, due to the action of said  
double-stranded marker protein ribonucleic acid sequence,  
from said population of plant cells, the selection being  
carried out under conditions under which the marker pro-  
tein can exert its toxic effect on the nontransformed  
25 cells.
2. The process as claimed in claim 1, wherein the marker protein  
is capable of converting directly or indirectly a substance X  
30 which is nontoxic for said population of plant cells into a  
substance Y which is toxic for said population, which process  
comprises the following steps:
  - 35 a) transforming the population of plant cells with at least  
one nucleic acid sequence to be inserted in combination  
with at least one double-stranded marker protein ribonu-  
cleic acid sequence or an expression cassette or expres-  
sion cassettes ensuring expression thereof capable of re-  
ducing the expression of at least one marker protein, and
  - 40 b) treating said population of plant cells with the sub-  
stance X at a concentration which causes a toxic effect  
for nontransformed cells, due to the conversion by the  
marker protein, and
  - 45 c) selecting transformed plant cells whose genome contains  
said nucleic acid sequence and which have a growth advan-  
tage over nontransformed cells, due to the action of said

2

double-stranded marker protein ribonucleic acid sequence, from said population of plant cells, the selection being carried out under conditions under which the marker protein can exert its toxic effect on the nontransformed cells.

- 5
3. The process as claimed in claim 2, wherein the nontoxic substance X is a substance which does not naturally occur in plant cells or organisms or occurs naturally therein only at a concentration which can essentially not cause any toxic effect.
- 10
4. The process as claimed in claim 2 or 3, wherein the substance X is a substance selected from the group consisting of pro-herbicides, proantibiotics, nucleoside analogs, 5-fluorocytosine, auxinamide compounds, naphthalacetamide, dihaloalkanes, Acyclovir, Ganciclovir, 1,2-deoxy-2-fluoro-b-D-arabinofuranosil-5-iodouracil, 6-thioxanthine, allopurinol, 6-methylpurine deoxyribonucleoside, 4-aminopyrazolopyrimidine, 2-amino-4-methoxybutanoic acid, 5-(trifluoromethyl)thioribose and allyl alcohol.
- 20
5. The process as claimed in any of claims 1 to 4, wherein the marker protein is selected from the group consisting of cytosine deaminases, cytochrome P-450 enzymes, indoleacetic acid hydrolases, haloalkane dehalogenases, thymidine kinases, guanine phosphoribosyl transferases, hypoxanthine phosphoribosyl transferases, xanthine guanine phosphoribosyl transferases, purine nucleoside phosphorylases, phosphonate monoester hydrolases, indoleacetamide synthases, indoleacetamide hydrolases, adenine phosphoribosyl transferases, methoxinine dehydrogenases, rhizobitoxin synthases, 5-methylthioribose kinases and alcohol dehydrogenases.
- 25
- 30
- 35
6. The process as claimed in any of claims 1 to 5, wherein the marker protein is encoded by
- 40
- a) a sequence described by the GenBank accession number S56903, M32238, NC003308, AE009419, AB016260, NC002147, M26950, J02224, V00470, V00467, U10247, M13422, X00221, M60917, U44852, M61151, AF039169, AB025110, AF212863, AC079674, X77943, M12196, AF172282, X04049 or AF253472
- 45
- b) a sequence according to SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40,

42, 44, 46 or 48.

7. The process as claimed in any of claims 1 to 6, wherein a sequence coding for a resistance to at least one toxin, antibiotic or herbicide is introduced together with the nucleic acid sequence to be inserted and selection is carried out additionally using the toxin, antibiotic or herbicide.
8. The process as claimed in any of claims 1 to 7, wherein the nucleic acid sequence to be inserted into the genome of the plant cell or of the plant organism comprises at least one expression cassette capable of expressing, under the control of a promoter functional in plant cells or in plant organisms, an RNA and/or a protein which does not cause the expression, amount, activity and/or function of a marker protein to be reduced.
9. The process as claimed in any of claims 1 to 8, wherein the plant cell is part of a plant organism or of a tissue, part, organ, cell culture or propagation material derived therefrom.
10. The process as claimed in any of claims 1 to 9 for preparing transformed plant cells or organisms, which comprises the following steps:
- a) transforming a population of plant cells which comprises at least one non-endogenous (preferably non-plant) marker protein capable of converting directly or indirectly a substance X which is nontoxic for said population of plant cells into a substance Y which is toxic for said population, with at least one nucleic acid sequence to be inserted in combination with at least one nucleic acid sequence coding for a double-stranded marker protein ribonucleic acid sequence or an expression cassette or expression cassettes ensuring expression thereof ribonucleic acid sequence capable of reducing the expression, amount, activity and/or function of said marker protein, and
- b) treating said population of plant cells with the substance X at a concentration which causes a toxic effect for nontransformed cells, due to the conversion by the marker protein, and



4

- 5 c) selecting transformed plant cells (and/or populations of plant cells, such as plant tissues or plants) whose genome contains said nucleic acid sequence and which have a growth advantage over nontransformed cells, due to the action of said double-stranded marker protein ribonucleic acid sequence, from said population of plant cells, the selection being carried out under conditions under which the marker protein can exert its toxic effect on the non-transformed cells, and
- 10 d) regenerating fertile plants, and
- 15 e) eliminating by crossing the nucleic acid sequence coding for the marker protein and isolating fertile plants whose genome contains said nucleic acid sequence but does not contain any longer the sequence coding for the marker protein.
- 20 11. An amino acid sequence coding for a plant 5-methylthioribose kinase, wherein said amino acid sequence contains at least one sequence selected from the group consisting of SEQ ID NO: 60, 62, 64, 66 or 68.
- 25 12. A nucleic acid sequence coding for a plant 5-methylthioribose kinase, wherein said nucleic acid sequence contains at least one sequence selected from the group consisting of SEQ ID NO: 59, 61, 63, 65 or 67.
- 30 13. A double-stranded RNA molecule, comprising
- 35 a) a "sense" RNA strand comprising at least one ribonucleotide sequence which is essentially identical to at least a part of the "sense" RNA transcript of a nucleic acid sequence coding for a marker protein, and
- 40 b) an "antisense" RNA strand which is essentially, preferably fully, complementary to the RNA sense strand under a).
- 45 14. The double-stranded RNA molecule as claimed in claim 13, wherein the marker protein is defined as in any of claims 2 to 6.
15. The double-stranded RNA molecule as claimed in either of claims 13 and 14, wherein the "sense" RNA strand and the "an-

## 5

tisense" RNA strand are covalently linked to one another in the form of an inverted repeat.

- 5 16. A transgenic expression cassette, comprising a nucleic acid sequence which codes for a double-stranded RNA molecule as claimed in any of claims 13 to 15 and which is functionally linked to a promoter functional in plant organisms.
- 10 17. A transgenic vector, comprising a transgenic expression cassette as claimed in claim 16.
- 15 18. A transgenic plant organism, comprising a double-stranded RNA molecule as claimed in any of claims 13 to 15, a transgenic expression cassette as claimed in claim 16 or a transgenic vector as claimed in claim 17.
- 20 19. The transgenic plant organism as claimed in claim 18, selected from the group of plants, consisting of wheat, oats, millet, barley, rye, corn, rice, buckwheat, sorghum, triticale, spelt, linseed, sugar cane, oilseed rape, cress, arabis, dopsis, cabbage species, soybean, alfalfa, pea, bean plants, peanut, potato, tobacco, tomato, eggplant, paprika, sunflower, tagetes, lettuce, calendula, melon, pumpkin and zucchini.
- 25 20. A tissue, an organ, a part, a cell, a cell culture or propagation material, derived from a transgenic plant organism as claimed in either of claims 18 and 19.

30

35

40

45

1/11




Fig. 1

2/11




Fig. 2-A

3/11




Fig. 2-B

4/11




Fig. 3

5/11




Fig. 4

6/11




Fig. 5



7/11




Fig. 6

8/11





Fig. 7

9/11



pSUN1-codA-RNAi-At.Act.-2-At.A/s-R-ocsT

Fig. 8

10/11

	1	50
Klebsiella pneumoniae	(1) -----MSQYHTFTAHDVAVYAAQ	
Clostridium tetani.	(1) -----MSRFDSHFRMETEDAILYAKE	
Zea mays	(1) ARALLSSPLAGASPCQSASAMAEQGFRLDESSLAYIKATPALAS	
A.thaliana	(1) -----MSFEEFTPLNEKSLVDYIKSTPALSS	
Brassica napus-2	(1) -----VDDFVLRAKEMSFDEFKPLNEKSLVEYIKATPALSS	
Soy-1	(1) -----	
Oryza sativa-1	(1) -----	
Consensus	(1) -----	L V A
	51	100
Klebsiella pneumoniae	(19) FAGIDNPSELVSAQEVGDGNLNLVFKVFDROGVSRAIVKQALPYVRCVGE	
Clostridium tetani.	(22) KLGIFDEHAKLQAEIIGDGNINIVFKVWDVNTKKSVIKHADIFLRSSGR	
Zea mays	(51) RLGGSGLSDSIEIKEVGDGNLNFVYIVQSEAGA--IVVKQALPYVRCVGD	
A.thaliana	(27) KIGADKSDDDLVIKEVGDGNLNFVIVVGSSGS--LVIKQALPYIRCIGE	
Brassica napus -2	(37) RLGDKY--DDLVIKEVGDGNLNFVIVVGSTGS--LVIKQALPYIRCIGE	
Soy -1	(1) -----	
Oryza sativa -1	(1) -----	
Consensus	(51) KLG D L EVGDGNLNFV V G LVIKQALPYIRCIGE	
	101	150
Klebsiella pneumoniae	(69) SWPLTLDRARLEAQTILVAHYQHSPQHTVKIHHFDP ELAVMV MEDLS-DHR	
Clostridium tetani.	(72) --ELDVDRNR IEAEV LMLQGILAPGLVPKVYKYSVMCNLSMEDIS-DHR	
Zea mays	(99) SWPMTTRERAYFEASTLREHGR LCP EHTPEVYHFDRTL SLMGMRYIEP PHI	
A.thaliana	(75) SWPMTKERAYFEATTLRKHG NLSPDHVPEVYHFDRTMALIGMRYLEP PHI	
Brassica napus -2	(83) SWPMTKERAYFEATTLRKHG GLSPDHVPEVYHFDRTMALIGMRYLEP PHI	
Soy -1	(1) -----IPEHVPEVYHFDRTMSLIGMRYLEP PHI	
sativa -1	(1) -----	
Consensus	(101) SWPMT ERA EA TL HG LSPDHVPEVYHFDRTMALIGMRYLEP PHI	
	151	200
Klebsiella pneumoniae	(118) IWRGELIANVYYPQAARQLGDYLAQVLFHTSDFY LHPHEKKAQVAQFIN-	
Clostridium tetani.	(119) NLRKELLKRNTFPSFAEHITTFIVDTLLPTTDLVMDSGEKKDNVKKYIN-	
Zea mays	(149) ILRKGLVAGVEYPLLADHMSDYMAKTLFFTSLLYNTTDDHKNGVAKYSAN	
A.thaliana	(125) ILRKGLIAGIEYPFLADHMSDYMAKTLFFTSLLYHDTTEHRRAVTEFCGN	
Brassica napus -2	(133) ILRKG-----	
Soy -1	(29) ILIKGLIAGIEYPFLAEHMA DFM AKTLFFTSLLFRSTADHKRDVAEFCGN	
Oryza sativa -1	(1) -----LLYNSTTDHKKGVAQYCDN	
Consensus	(151) ILRKGLIA I YP ADHM DYMA TLF TSLLY T DHK VA F N	
	201	250
Klebsiella pneumoniae	(167) PAMCEITEDLFNDPYQIHERN--NYP AELEADVAALRDDAQLKLAVAAL	
Clostridium tetani.	(168) KDLCKISEDLVFTEPFIDYKSRNTVLEENIEFVKRQLYEDKELILEAGKL	
Zea mays	(199) VEMCRLTEQVVFSDPYRVSKFNR-WTSPYLDKDAEAVREDDELKLEVAGL	
A.thaliana	(175) VELCRLTEQVVFSDPYRVSTFNR-WTSPYLD DDAKAVREDSALKLEIAEL	
Brassica napus -2	(138) -----	
Soy -1	(79) VELCRLTEQVVFSDPYKVSQYNR-WTSPYLD RDAEAVREDNLLKLEVAEL	
Oryza sativa -1	(20) VEMCRLTEQVVFSDPYMLAKYNR-CTSPFLDNDAAVREDAELKLEIAEL	
Consensus	(201) VELCRLTEQVVFSDPY VS FNR TSPYLD DA AVRED LKLEVA L	

Fig. 9a

11/11

251	300
Klebsiella pneumoniae	(215) KHRFFAHAEALLHGDIHSGSIFVAEGSLKAIDAEFGYFGPIGFDIGTAIG
Clostridium tetani.	(218) KNNFMNNSQALIHGDLHSGSIFVNEESTKILDPEFAFYGPIGYDLGNVIG
Zea mays	(248) KSMFIERAQALIHGDLHTGSIMVTEVQLKSLIQNLGSMGPMGFDIGSLPW
A.thaliana	(224) KSMFCERAQALIHGDLHTGSVMVTQDSTQVIDPEFSFYGPMGFDIGAYLG
Brassica napus -2	(138) -----
Soy -1	(128) KSKFIES-----
Oryza sativa -1	(69) KSMFIERAQUALIHGDLHTGSIMVTPDSTQVIDPEFAFYGPMGYDIGAFLG
Consensus	(251) KS FIE AQALIHGDLHTGSI V S ID EFAFYGPMGFDIG IG
	301 350
Klebsiella pneumoniae	(265) NLLNLYCGLPGQLGIRDAAAAREQRLNDIHQLWTTFAERFQALAAEKTRD
Clostridium tetani.	(268) NLFFAWANAYVTEDGKEVEEFTIWIEKTIENILELFKEKFIKKYKEIVTD
Zea mays	(298) KPDFGHTMHRMGMLIKRMIVRLTRMDLEDN-----
A.thaliana	(274) NLILAFFAQDGHATQENDRKEYKQWILRTIEQTWNLFNKRFIALWDQNKD
Brassica napus -2	(138) -----
Soy -1	(135) -----
Oryza sativa -1	(119) NLILAYFSQDGHADQANDRKAY-----
Consensus	(301) NL AY
	351 400
Klebsiella pneumoniae	(315) AALAYPGYASAFLLKKVWADAVGFCGSELIRRSVGLSHVADIDTIQDDAMR
Clostridium tetani.	(318) VMAKEEYYMNWYLSILSDTAGQVGLIIRRVVGDSKVLDITSITDINKR
Zea mays	(328) -----
A.thaliana	(324) GPGEAYLADIYNNTEVLKQVQENYMRNLLHDSLGFGAAMIRRIVGVAHV
Brassica napus -2	(138) -----
Soy -1	(135) -----
Oryza sativa -1	(141) -----
Consensus	(351) -----
	401 447
Klebsiella pneumoniae	(365) HECLRHAITLGRALIVLAERIDSVDELLARVRQYS-----
Clostridium tetani.	(368) VKAERILILSAKTFIKNRHKIKTGKRYVEIFNSNMY-----
Zea mays	(328) -----
A.thaliana	(374) EDFESIEEDKRRRAICERSALEFAKMLLKERRKFKSIGEVVSAIQQOS
Brassica napus -2	(138) -----
Soy -1	(135) -----
Oryza sativa -1	(141) -----
Consensus	(401) -----

Fig. 9b

**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

**THIS PAGE BLANK (USPTO)**